

**CHIRONOMUS NEWSLETTER ON CHIRONOMIDAE  
RESEARCH**

Co-Editors:  
Ruth CONTRERAS-LICHTENBERG, Naturhistorisches Museum Wien,  
Burgring 7, A-1014 WIEN, Austria  
Peter H. LANGTON, 5 Kylebeg Avenue, Mountsandel Colerane,  
Co. LONDONDERRY, Northern Ireland

Bibliography:  
Odwin HOFFRICHTER, Institut Für Biologie I, Albert-Ludwigsuniversität,  
Freiburg, Hauptstrasse 1, D-79104 Germany  
Treasurer: Trond ANDERSEN, Museum of Zoology, University of Bergen,  
Museplass 3, N-5007 Bergen-Norway

ISSN 0172-1941

No. 17

October 2004

**CONTENTS**


Memorial Meeting of the Russian Chironomidologists on centenary of Aleksey A. Chernovsky .....	1
The chironomid collection of the Hydrobiological Station Falkau/Black Forest .....	3
Theses .....	4
Short - Communications .....	6
List of regional representatives 2004 .....	9
Current Bibliography .....	12



**Zoological Institute, Russian Academy of Sciences (RAS),  
St. Petersburg, Russia  
Institute of Biology and Soil Sciences, Far East Branch RAS,  
Vladivostok, Russia  
Russian Dipterological School, St. Petersburg, Russia  
Russian Entomological Society**



**MEMORIAL MEETING OF THE RUSSIAN CHIRONOMIDOLOGISTS ON  
CENTENARY OF ALEKSEY A. CHERNOVSKY**



(15.03.1904 – 11.02.1942)

‘During two days, March 15-16<sup>th</sup>, 2004, the Memorial meeting of Russian chironomists on the centenary of the birth of Aleksey Alekseevitch Chernovsky took place at the Zoological Institute of the Russian Academy of Sciences.

A.A. Chernovsky had a short life but he was a talented freshwater hydrobiologist and chironomid taxonomist, a very kind and real man. All Russian scientists began the study of chironomids after his famous Key book on chironomid larvae of the USSR (1949). Now Russian chironomists continue investigations of chironomids which were started by A.A. Chernovsky. In support of it, 18 scientists from many Russian Institutes situated in Europe, Siberia and ‘the Far East presented lectures at the conference. Most of these lectures will be published in a special issue of the “EUROASIAN ENTOMOLOGICAL JOURNAL” in December 2004.

List of papers of the Special Issue (Number 4)

1. BALUSHKINA E.V. Changes in the structure of benthic animal communities under anthropogenic impact on aquatic ecosystems (by the example of small lakes of Leningrad Region).
2. ERBAEVA E.A., SAFRONOV G.P. Chironomidae of the Angara river and its reservoirs.
3. ERMOLAEVA O. V. Comparative karyological analysis of chironomids from Diamesinae and Prodiamesinae subfamilies (Diptera, Chironomidae).
4. ISTOMINA A.G., KIKNADZE I.I. Chironomus bernensis in West Siberia. Karyotype and chromosome polymorphism.
5. KIKNADZE I.I., GUNDERINA L.I. ISTOMINA., A.G, GUSEV V.D., NEMYTIKOVA L. A. Reconstruction of chromosomal evolution in the genus *Chironomus*.
6. KLISHKO O.K. About the new finding of *Paratanytarsus baicalensis* (Tshern.) (Diptera, Chironomidae) in the Upper Amur basin.
7. KRAVTSOVA L.S. Spatial distribution of chironomids in different habitats (Lake Baikal and its tributaries).
8. MAKARCHENKO E.A., MAKARCHENKO M.A. Chironomids of the genus *Rheocricotopus* Thienemann et Harnisch, 1932 (Diptera, Chironomidae, Orthoclaadiinae) of the Russian Far East.
9. MOROZOVA E.E. Karyotype and chromosome polymorphism of *Cryptochironomus obreptans* Walk. (Diptera, Chironomidae) from the Volga.
10. NARTSHUK E.P. Chironomid midges – as the most adapted group of Diptera to aquatic conditions.
11. NAZAROVA L.B., BROOKS S.J. Chironomids in paleoclimatic investigations.
12. PETROVA N.A., MICHAILOVA P.V., CHUBAREVA L.A., SHOBANOV N.A., ZELENTSOV N.I. The system of A.A. Chernovsky [1949] as a base of the cytotaxonomy of the family Chironomidae.
13. POLUKONOVA N.V. To specification of subgenus structure of a genus *Chironomus* Meigen, 1803 (Diptera, Chironomidae) with description of new subgenus *Halochironomus* n. subg.
14. SERGEEVA I.V. The revision of the Russian Tanypodinae of the genus *Clinotanypus* Kieffer, 1913 (Diptera, Chironomidae, Tanypodinae).
15. ZORINA O.V. Chironomids of the genus *Paratendipes* Kieffer (Diptera, Chironomidae, Chironominae) of the Russian Far East.

You can subscribe this Special Chironomid Issue by following address:

K.G. Mikhailov

Fax +7 (095) 203–2717

E–mail: [kmk2000@online.ru](mailto:kmk2000@online.ru)

You can find more detail information about the subscription and the Journal here –

<http://euroentjourn.narod.ru/indexengs.html>

**This Information was prepared by Eugenyi Makarchenko**

## The chironomid collection of the Hydrobiological Station Falkau/Black Forest.

By Wolfgang Wuelker

In May 1952 I began working as scientific assistant of the Hydrobiological Station Falkau (later the Limnological Institute of the University of Freiburg, Walter Schlienz-Institut). I was advised to investigate the fauna of the lakes, in the middle of which the Hydrobiological Station was situated. For chironomids, at that time this was possible only with the most appreciated help of Prof. Thienemann and his crew in Ploen.

Previously, members and guests of the Institute (Prof.Dr.H.J.Elster, Frau Dr.H.Hauer-Eichhardt, Dr.J.Lundbeck) had collected over the years 1948-52 as many as 2200 samples, mainly from 7 lakes (Key numbers in brackets): Feldsee near Feldberg, 1109m NN (I), Windgfaellwiher near Altglashuetten, 966m NN (II), Schluchsee, 930m NN (III), Titisee, 846m NN (IV), Ursee near Lenzkirch, 837m NN (V), Mathisleweiher near Hinterzarten, 1000m NN (VI) and Schluechtsee near Rothaus, 914m NN (VII).


During the following years (1952-57) the number of samples in the Black Forest increased to nearly 3000, moreover the collection had been enriched by material from my research trips to Spain (1954), Sudan (1963), USA and Canada (1964), Alps (different years), Kenya (1975), Fennoscandinavia (1956, 1980, 1985), Hungary (1981) and Russia, especially Siberia (between 1989 and 1999).

Today, the material is concentrated on chromosome preparations of the genera *Chironomus* and *Sergentia* (about 100 slide boxes), but contains also permanent preparations of larvae, pupal exuviae and adults, examples of parasitic intersexuality and of the structure of imaginal discs, as well as bottled alcohol material (no pinned specimens!). Protocols are present, but not for 1948-1952.

The collection is at present in my private home, but there is the provision for it to be incorporated in the Zoologische Staatssammlung Munich.

I left Falkau in 1962 in favour of a career as zoologist in the University of Freiburg; the Hydrobiological Station in 1971 turned into the Limnological Institute of the University of Konstanz. Nevertheless, I am trying to make the chironomid collection of Falkau ready for use for all interested chironomists.

Wolfgang Wuelker, Professor of Zoology and Limnology  
Kalkackerweg 28, D 79111 Freiburg, Germany  
Tel. xx49 0761 4764492, Fax xx49 0761 4702813  
e.mail [Wolfgang.Wuelker@biologie.uni.freiburg.de](mailto:Wolfgang.Wuelker@biologie.uni.freiburg.de)



## **THESES**

### **PHD-THESIS ON “CHIRONOMIDS (DIPTERA, NEMATOCERA) OF TEMPORARY POOLS – AN ECOLOGICAL CASE STUDY” (2004) (IN ENGLISH)**

By P.-M. Andreas Dettinger-Klemm

This thesis (371pp, 96 figures and 74 tables) is downloadable at <http://archiv.ub.uni-marburg.de/diss/z2004/0076>. It is a synopsis of field work (three pools of different habitat permanence were investigated for 5-7 years) and laboratory investigations (main aspects: growth and development in relation to photoperiod and temperature, drought tolerance, parthenogenesis) of the four principal species – *Chironomus dorsalis* sensu STRENZKE (1959) and KEYL & KEYL (1959), *Polypedilum tritum* (WALKER, 1856), *Limnophyes asquamatus* ANDERSEN, 1937 and *Paralimnophyes hydrophilus* (GOETGHEBUER, 1921) – and summarizes our present knowledge of the topic. Taxonomic and morphological aspects of the four typical pool species are also presented (including the description of the larvae and pupae of *L. asquamatus*, *L. minimus* sensu stricto (MEIGEN, 1818) and *L. natalensis* (KIEFFER, 1914), the definition of three ecotypes in *L. asquamatus* and indications that *Polypedilum tritum* and *P. uncinatum* GOETGHEBUER, 1921, may be synonyms).

An abstract (English and German) of the thesis is available at the above web address. The PDF-document is separated into 14 documents, which can be downloaded separately: document 1 “Contents”, document 2 “Preface & Acknowledgements”, document 3 “Introduction”, document 4 “Materials & Methods”, document 5 “Results/The Habitat”, document 6 “Results/The Chironomid Community”, document 7 “Results/Morphology & Taxonomy”, document 8 “Results/ Laboratory Studies”, document 9 “Results/Field Study”, document 10 “Discussion”, document 11 “Summary”, document 12 “Zusammenfassung”, document 13 “References” and document 14 “Appendix”.

*Hard copies or a CD-version of the thesis also can be ordered from Andreas Dettinger-Klemm, Plattenhof, D-64560, Riedstadt-Erfelden, Germany ([dettingerklemm@aol.com](mailto:dettingerklemm@aol.com)).*

### **A STUDY OF THE FRESHWATER MACROINVERTEBRATE FAUNA OF MADEIRA AND THEIR APPLICATION IN A REGIONAL ECOLOGICAL ASSESSMENT SYSTEM.**

Samantha Jane Hughes

Thesis submitted to King’s College, University of London, in partial fulfillment requirements of the degree of Doctor of Philosophy; March 2003

#### **Abstract**

This is the first comprehensive overview and baseline study of the physicochemical properties and benthic macroinvertebrate fauna of the lotic habitats of the island of Madeira. The findings provide the essential basis for developing monitoring systems for the island’s surface waters, in compliance with the Water Framework Directive (2000/60/CE).

Factors shaping local hydrological resources such as geology, orography, climate and vegetation are described as well as the hydrology and physical aspects of the surface waters. Benthic macroinvertebrate-based ecological assessment methods are reviewed together with their limitations when applied to an impoverished island fauna.

Physicochemical and biological hand net samples were taken along physical and pollution gradients at a total of 45 sites in 10 catchments in 1991 and 1997. Classification and ordination revealed physicochemical gradients related to altitude, season and anthropogenic input. A preliminary physicochemical classification system is proposed.

A revised checklist reveals over 25% of the macroinvertebrate fauna to be endemic, illustrating the importance of implementing conservation measures. Samples exhibited low abundance and diversity with several predominant taxa, emphasising the “disharmonic” nature of the macroinvertebrate fauna. Classification and ordination revealed distinct faunal assemblages at sites with good ecological quality and at impacted sites. Sites of intermediate quality at lower elevations are difficult to separate due to seasonal variation, a lack of strong unimodal indicator taxa, the predominance of taxa normally associated with low ecological quality and the restricted distribution of pollution intolerant taxa. These factors limit the direct comparison of chemical quality to ecological quality and also the application of 20 well-established metrics. The Madeiran Biotic Score II, an improved ecological assessment system is proposed which better represents ecological quality at low altitude sites. The Chironomidae is the most diverse macroinvertebrate group on Madeira. Analyses of amalgamated samples of pupal exuviae indicate that unpolluted sites are more diverse and with a greater proportion of intolerant taxa than polluted sites. In the light of these findings, comprehensive multimetric monitoring techniques are proposed to include a suite of biological elements.

<b>Chapter 6. Results: The Chironomidae</b>	224
<b>6.1</b> Introduction	224
<b>6.2</b> A brief introduction to the Chironomidae (Diptera: Nematocera) and CPET	224
<b>6.3</b> A review of the Chironomidae of Madeira and other Atlantic Islands	229
<b>6.3.1</b> The biogeography of the Chironomidae of Madeira	233
<b>6.4</b> Global analysis of the collections	235
<b>6.4.1</b> Relative abundance	235
<b>6.4.2</b> Taxon richness and sampling effort	237
<b>6.4.3</b> Taxa accretion and rank abundance	239
<b>6.4.4</b> Relative and rank abundance of individuals	241
<b>6.4.5</b> Frequency of occurrence, distribution over sampling sites and voltinism	246
<b>6.4.6</b> Species diversity	254
<b>6.4.7</b> Tolerance ranges and trophic groups	257
<b>6.4.8</b> -Assessing ecological quality using CPET	261
a Single samples	261
b Amalgamated samples	265
<b>6.4.9</b> Classification and Ordination	266
<b>6.4.10</b> Classification using cluster analysis	267
<b>6.4.11</b> Ordination using Principal Components Analysis	274
<b>6.4.12</b> Conclusions from multivariate analyses	

## **“STUDY OF PHOTOBHAVIOUR IN CHIRONOMUS RAMOSUS”**

**(M.Phil Thesis by Mr. Anand A. Babrekar)**

The ontogeny of photosensitivity in a holometabolous insect midge *Chironomus ramosus* was studied. Extracellular electrical activity was recorded from larval and adult photoreceptor organs. We found a progressive increase in photosensitivity, as the development proceeds from larva to adult stage. This is a first report of its kind where developmental profile of photosensitivity in any insect has been described from an ecological context. Chironomid midges have been chosen for this study since developmental stages show ecological transitions. Aquatic bottom dwelling tubicolous larvae metamorphose to a transient pupal stage and subsequently, eclose to terrestrial low-flying adult midges. Unlike larvae, adults were found to be positive phototactic.

We have fabricated special devices and designed novel assays to study photobehavioural responses in different developmental stages. We have also formulated phototactic index (P.I.) for quantitative analysis of photobehaviour. Moreover, the study aimed at finding whether response to intensity and wavelength of light varies in different developmental stages. Interestingly, our study showed a developmental shift in photobehavioural response during metamorphosis. Early and late larval instars showed variable pattern of photoresponse under shorter and longer wavelength of visible spectrum and the findings have been correlated to their ecological transitions from pelagic to benthic life style. Similarly, P.I. values shifted from positive to negative and vice-versa throughout the life-cycle of *Chironomus*. Behavioural data has been corroborated with ERG data. (collaborative work with Dr. Gauri Kulkarni, Biophysics Unit, Dept. of Physics, University of Pune). Our electrophysiological data link sensitivity of developmental stage specific photoreceptor organs of larvae and adults to its ecological adaptations.

*Chironomus* larvae are known as pests and adults are known for creating a nuisance in human habitats. Adult midges are also medically known for allergic reactions to human. Therefore, we believe that our findings will help in designing ‘light-trap’ as an effective pest control strategy based on developmental stage specific photosensitivity

## ***SHORT – COMMUNICATIONS***

### **NEWS FROM INDIA**

**The study of DIPTERAN TAXONOMY  
under TAXONOMIC CAPACITY BUILDING PROJECT of All India Coordinated project on Insect  
Taxonomy  
funded by the Ministry of Environment & Forests, Govt. of India  
Drs. P.K.Chaudhuri and A.Mazumdar  
Department of Zoology, University of Burdwan**

This project is progressing satisfactorily. During the past few years, a good number of ecosystems of the Indian subregion have been explored. A large number of genera and species with life stages have been recorded for the first time in the country. Identification of the insects is still going on. In addition, assessment of riverine water quality using chironomids has been initiated in the laboratory. Chironomid material /publications of

workers around the world are being solicited for the purpose of building a Center for Dipteran Taxonomy .

**Dr. Mazumdar** has also started an investigation into the cytotoxic categorization of the Indian species of *Chironomus* Meigen and *Glyptotendipes* Kieffer. Suggestions / advice or literature on the subject of cytotoxicity of chironomids will be gratefully acknowledged.

#### CHIRONOMID RESEARCH AT PUNE UNIVERSITY

Anand A. Babrekar received Prof. V.C.Shah best poster presentation award for the paper entitled “Structural & functional analysis of photoreceptor cells of *Chironomus ramosus*” at the XXVII All India Cell Biology Conference & International Symposium, Jan 7-10,2004 at Pune, India.

He received his M.Phil in Zoology at Pune University, India in 2003 and worked under the supervision of Dr. B.B.Nath, Reader Department of Zoology, University of Pune, Pune, India. Exerpts of his M.Phil thesis is as follows :

#### Research papers published in 2003-2004.

1. **Hazra Niladri, Mazumdar, A. & P.K. Chaudhuri, 2003. Life stages of four species of *Polypedilum* Kieffer (Diptera: Chironomidae) with a new species of the genus from Darjeeling-Sikkim Himalayas.** Belgian Entomology Journal 5 (1-2): 125-137 (2003).
2. **Hazra, Niladri, Nath, Saswati & P.K.Chaudhuri, 2003. Genus *Corynoneura* Winnertz (Diptera: Chironomidae) from Darjeeling-Sikkim Himalayas of India with description of three new species.** Ent. Monthly Mag., 139 : 69-82 .

#### Addresses of workers on Indian Chironomids

1. Prof. P.K.Chaudhuri  
Department of Zoology, University of  
Burdwan, Burdwan 713 104.  
Telephone: +91- 342 – 2559884  
(Lab.), Fax: +91 – 342 – 2530452 (request)  
Email: [chaudhuri\\_pk@yahoo.co.in](mailto:chaudhuri_pk@yahoo.co.in)  
(Interest- Biosystematics, Biogeography &  
Cladistics of Indian chironomids)
2. Dr. J.R.B.Alfred, Director, Zoological Survey  
of India  
535,M Block, Alipore, Calcutta 700 053.  
Email: [enviszsi@cal.vsnl.net.in](mailto:enviszsi@cal.vsnl.net.in)  
(Ecology of lotic chironomids).
3. Dr. Bimalendu Nath, Reader, Department of  
Zoology,  
University of Pune,  
Pune 412 007.  
Telefax +91-20-2569-0087.  
Email: [<bbnath@unipune.ernet.in>](mailto:bbnath@unipune.ernet.in)  
(Karyosystematics & Electrophysiology of  
chironomids)
4. Dr. Sailesh Chattopadhyay, Associate  
Professor,  
Department of Entomology, Birsa Agricultural  
University  
Kanke Road, Ranchi 830 006.  
Telephone +91- 651-3092144
5. Dr. Dipak K. Som and Mr. U.K. Mandal  
Department of Zoology, Hooghly Mohsin  
College (University of Burdwan),  
Chinsurah-712 101, India  
Telephone : +91-09830179230  
Email: “Dipak Som” [<dsom@vsnl.net.in>](mailto:dsom@vsnl.net.in)  
(Lotic and lentic chironomids)
6. Mr. U. Majumdar,  
Department of Zoology, University of  
Burdwan, Burdwan 713 104, India  
Email: [uttaran11@rediffmail.com](mailto:uttaran11@rediffmail.com)  
(Biosystematics & Cladistics analysis )

7. Mr. Nirmalya Das,  
Lecturer, Department of Zoology,  
Kalimpong College (University of North  
Bengal),  
Kalimpong 734 301, Dt. Darjeeling.  
Email: dnirmalya@yahoo.com  
(Interest : Biosystematics of Orthoclad)
8. Dr. A.K. Sadhu,  
Department of Zoology, Burdwan Raj College,  
Burdwan 713 104, India  
Telephone +91-342-2568956  
Email: <arupsadhu@rediffmail.com>  
(chironomids as fish food)
9. Dr. Pargya Khanna  
Department of Zoology, University of Jammu,  
Jammu 180 006, J & K.  
Tel.: +91-191-2435030  
Email: <Pragyajamu@yahoo.co.in>  
( Biosystematics and Ecology of Chironomids)
10. Dr. Rishi Das,  
Department of Zoology, Tinsukia College,  
Tinsukia 786 125, Assam, India.  
( Ecology and Biochemistry of haemoglobin of  
Chironomids).

## **RED LIST OF ENDANGERED CHIRONOMID SPECIES OF BAVARIA (GERMANY)**

By Drs. Claus Orendt & Nicola Reiff

It was with considerable astonishment that I received a copy of this work through e.mail. Not that receiving it by e.mail was an astonishment, nor that the authors had so kindly sent it me, but the very thought that there was anywhere in the world so advanced in the recording of its chironomids and habitat evaluation that chironomid species could be listed to any degree of confidence as endangered was....but then, this list is about Bavaria, a state where both data sets are far advanced (perhaps only for the Black Forest Region of Germany and for Holland is there a similar data base from which to generate such a list). Claus Orendt and Nicola Reiff are to be commended on submitting this work to the scrutiny of the chironomid fraternity: anyone who has collected in Bavaria and has information that will be of use to the authors is encouraged to send it to them. Freshwater habitats are particularly under threat by an expanding human population and industry; data like these may save more of them from destruction. It is worth looking at, not just to see what can be done, but also to check out the list – you will find species there that you have not encountered (perhaps not even heard of!) and some that will surprise you, as did to me the listing of *Metriocnemus eurynotus* as endangered (in Britain it is the most widespread and frequent midge of winter-time).

**Copies may be obtained (in German or English) on <http://www.orendt-hydro.de/roteli.htm>**

Editorial comment by Peter H. Langton

### **Pink *Chironomus exuviae*!**

By Peter H. Langton

In the English language the colour pink is used to describe a number of character states: physiological (“in the pink”, i.e. in good health); sociological (“pink with embarrassment”), psychological (“seeing the world through rose tinted spectacles”) and is the colour favoured for young girls. All quite interesting, perhaps, and nothing really to do with the report I am about to begin.

Earlier this year I was on a day’s outing with the family by the sea in Donegal (Ireland). The sun shone brightly and there was little air movement – caps, factor 16, sunglasses – in fact a great day to be out in the open by the sea. Yes, this is also a bit of a digression, but the weather here in the north of Ireland gets a bad press and it’s nice to record just how pleasant it can be. Being a bit past the youthful enthusiasm for building sand castles, I happily kept overall watch from a distance turning over stones and other discarded objects from the building plot nearby. Amongst these artefacts I came across a large plastic bucket (diameter about 40cm.) half-full of water (about 20cm deep). The water



was clear, there were a few small leaf fragments in the bottom and some chironomid larval galleries. Floating at the edge was a number of Chironomus exuviae, bright pink in colour. I have some considerable experience of Chironomus pupal exuviae and never encountered bright pink ones before: at the very least, surely, a species new to Ireland...perhaps even to science... As soon as I was able after returning home, I had one under the microscope: strange...the exuviae weren't pink at all - nothing other than Chironomus piger! The pink colour was due to the exuviae being full of the alga Haematococcus pluvialis. I refer you back to the state of the water: transparent, uncoloured. All explanations that have occurred to me seem inadequate: surely chironomid exuviae cannot be such efficient nets; they frequently contain small animals and algae, but I have always considered this fortuitous. The time exuviae remain afloat is but one or two days: it seems unlikely that the conditions inside the exuviae are such that an algal cell or two would generate such a large population in the time. My best suggestion is that the algae are trapped in the meniscus at the edge of the water where the exuviae congregate and here the algae are scooped up when the water is agitated by a breeze...any better suggestion?

## REGIONAL REPRESENTATIVES 2004

### AFRICA

AMAKYE, Josef S., Institute of Aquatic Biology (C.S.I.R.), P.O.Box 38, Achimota - Accra, Ghana. **Regional representative for West Africa**

### AMERICAS

MASAFERRO, Julieta, Department of Entomology, Natural History Museum, Cromwell Road, London SW7 5BD, U.K. Tel: +44 (0) 207 942 5198. Fax: +44 (0) 207 942 5229. Email: [J.Massafferro@nhm.ac.uk](mailto:J.Massafferro@nhm.ac.uk) **Regional representative for Argentina**

CALLISTO, Marcos, Universidade Federal de Minas Gerais, ICB, Depto. Biologia Geral, Lab. Limnologia/Ecologia de Bentos, CP. 486, CEP. 30.161-970, Belo Horizonte, MG, Brazil. Tel: +55 31 499 25 95. Fax: +55 31 499 25 67. Email : [callisto@mono.icb.ufmg.br](mailto:callisto@mono.icb.ufmg.br) **Regional representative for Brazil**

WALKER, Ian R., Science Building, Room 161 Department of Biology North Kelowna Campus Okanagan University College 3333 University Way Kelowna, British Columbia CANADA V1V 1V7 E-mail: [iwalker@ouc.bc.ca](mailto:iwalker@ouc.bc.ca) Tel. Office: (250) 762-5445 local 7559 Fax: (250) 470-6004 **Regional representative for Canada**

DE LA ROSA, Carlos, Program Director, The Nature Conservancy's Disney Wilderness Preserve, 2700 Scrub Jay Trail, Kissimmee, FL 34759, U.S. Tel.: ( 407) 935-0002 ext. 103.

Fax: ( 407) 935-0005. Email: [cdelarosa@tnc.org](mailto:cdelarosa@tnc.org) **Regional representative for Central America**

ALCOCER, Javier, UNAM Campus Iztacala, Limnology Lab, Environmental Conservation & Improvement Project, Universidad Nacional Autonoma de Mexico, Calle 15, #51 San Pedro de los Pinos, Mexico City, D.F. 03800, Mexico. Fax: (525) 277-1829. Email: [jalcocer@unamvm1.dgsca.unam.mx](mailto:jalcocer@unamvm1.dgsca.unam.mx) **Regional representative for Mexico**

BURGOS, Arnoldine, Bureau voor Openbare Gezondheidszorg, Centraal laboratorium, Rode Kruislaan 13 Postbus 1911, Paramaribo, Suriname. **Regional representative for Suriname**

FERRINGTON Jr., L.C., Department of Entomology, Hodson Hall, 1980 Folwell Avenue, University of Minnesota, St. Paul, MN 55108, U.S. Tel.: (612) 624-3265. Email: [ferri016@umn.edu](mailto:ferri016@umn.edu) **Regional representative for the U.S.A.**

### ASIA

WANG, Xinhua, Department of Biology, Nankai University, Tianjin, 300071 China. Phone: +86 22 23508389 (Office), +86 22 23501450 (Home), Fax: +86 22 23508800. Email: [xhwang@nankai.edu.cn](mailto:xhwang@nankai.edu.cn) **Regional representative for the P. R. of China**

MAZUMDAR, Abhijit, Dept of Zoology, University of Burdwan, Burdwan 713 104, W.B., India Email: [abhijitau@rediffmail.com](mailto:abhijitau@rediffmail.com) **Regional representative for India**

MOUSAYI, Seyed Karim, University of Tromsø, Institute of Clinical Medicine (IKM), KK Lab, Brevika Centre, N-9037 Tromsø, Norway. Tel: +47 77 64 48 48. Email:

[karimm@fagmed.uit.no](mailto:karimm@fagmed.uit.no) **Regional representative for Iran**

KUGLER, Jehoshua, Dept of Zoology, Tel-Aviv University, Tel-Aviv 69978, Israel.

**Regional representative for Israel**

IWAKUMA, Toshio, Hokkaido University, Kita-Jujo-Nishi 5, Kita-ku, Sapporo, Hokkaido, 060 Japan. Email: [iwakuma@ees.hokudai.ac.jp](mailto:iwakuma@ees.hokudai.ac.jp)

**Regional representative for Japan**

ISMAIL, A.R., Jabatan Biologie, University Pertanian Malaysia, 43400 UPM Serdang, Selangor, Malaysia. **Regional representative for Malaysia**

MAKARCHENKO, Eugenyi A., Laboratory of Freshwater Hydrobiology, Institute of Biology and Soil Sciences, Far Eastern Branch of the Russian Academy of Sciences, 690022 Vladivostok - 22, Russia. Fax: +7 (4232) 310193. Email: [emakarchenko@mail.ru](mailto:emakarchenko@mail.ru) (home) and [emakar@ibss.dvo.ru](mailto:emakar@ibss.dvo.ru) (work)

WWW: [www.tendipes.febras.ru](http://www.tendipes.febras.ru) **Regional representative for the Far East of Russia**

ERBAEVA, Engelsina, Institute of Biology, Irkutsk State University, P.O. Box 24, Lenin street 3, 664033 Irkutsk, Russia. **Regional representative for Lake Baikal and River Angara, South Siberia**

BYEONG-JIN, Youn, Pusan National University, Department of Biology, #30 Changjeon-dong Kumjeong-Ku, Pusan 609-735, South Korea. **Regional representative for South Korea**

## EUROPE

CONTRERAS, Ruth, Naturhistorisches Museum, 2. Zoologische Abt., Burgring 7 (Box 417), A-1014 Wien, Austria. Tel: ++43 01 521 77/317. Fax: ++43 01 523 52 54.

Email: [ruth.contreras@aon.at](mailto:ruth.contreras@aon.at) **Regional representative for Austria**

GODDEERIS, Boudewijn, Koninklijk Belgisch Instituut voor Natuurwetenschappen, Afdeling Zoetwaterbiologie, Vautierstraat 29, B-1040 Brussel, Belgium. Tel: ++32 2 627 43 14. Fax: ++ 32 2 646 44 33. Email:

[goddeeri@kbinirnsnb.be](mailto:goddeeri@kbinirnsnb.be) **Regional representative for Belgium**

MICHAILOVA, Parasaleva, Institute of Zoology, boul. Rouski 1, Bulgarian Academy of Sciences, Institute of Zoology, Sofia 1000, Bulgaria. Email: [parmich@mail.bol.bg](mailto:parmich@mail.bol.bg)

**Regional representative for Bulgaria**

CHVOJKA, Pavel, National Museum, Department of Entomology, Kunratice 1, 14800 Praha, Czechia. **Regional representative for Czech Republic**

LINDEGAARD, Claus, Freshwater Biological Laboratory, University of Copenhagen, 51 Helsingorsgade, DK-3400, Hillerød, Denmark. Fax: +45 48241476. Email:

[clindegaard@zi.ku.dk](mailto:clindegaard@zi.ku.dk) **Regional representative for Denmark**

KANGUR, Andu & Kulli, Vorstjarv Limnological station, EE2454 Rannu, Estonia. Fax: +372 343 3472. Email:

[andu@lim.tartu.ee](mailto:andu@lim.tartu.ee) **Regional representative for Estonia** andu@lim.tartu.ee

KOSKENNIEMI, Esa, West Finland Regional Environment Centre, P.O. Box 262, FIN-65101 Vaasa, Finland. Tel: + 358-6-3675453. Fax: + 358-6-3675451. Email:

[Esa.Koskenniemi@environment.fi](mailto:Esa.Koskenniemi@environment.fi) **Regional representative for Finland**

DELETTRE, Yannick R., C.N.R.S. (U.M.R. 6553 "ECOBIO") - Université de Rennes I, Station Biologique, F-35380 Paimpont, France. Tel: (+33) 02.99.61.81.73. Fax: (+33) 02.99.61.81.87 (from abroad don't dial the zero). Email: [yannick.Delettre@univ-rennes1.fr](mailto:yannick.Delettre@univ-rennes1.fr) WWW: <http://ecobio.univ-rennes1.fr/Fiches%20perso/YDelettre/YDelett>

[re.htm](http://ecobio.univ-rennes1.fr/Fiches%20perso/YDelettre/YDelett) **Regional representative for France**

SPIES, Martin, Schraemelstr. 151, D-81247 München, Germany. Fax : + 49 89 8107 300.

Email : [spies@zi.biologie.uni-muenchen.de](mailto:spies@zi.biologie.uni-muenchen.de) **Regional representative for Germany**

DÉVAI, György, L. Kossuth University, Ecological Institute, H-4010 Debrecen, Hungary. <http://www.ecol.klte.hu/yellow.html> **Regional representative for Hungary**

OLAFSSON, Jon Sigurdur, University of Iceland, Institute of Biology, Grensasvegi 12, 108 Reykjavik, Iceland. Email: [jsol@rhi.hi.is](mailto:jsol@rhi.hi.is) **Regional representative for Iceland**

MURRAY, Declan, Dept of Zoology, University College Dublin, Belfield, Stillorgan Road, Dublin 4, Ireland. Tel: +353-1-706-

2336. Fax: +353-1-7061152. Email:  
[declan.murray@ucd.ie](mailto:declan.murray@ucd.ie) WWW:  
<http://www.ucd.ie/~zoology/zoology.html>  
**Regional representative for Ireland**

ROSSARO, Bruno, Univ. of Milano, Dept. of  
Biology, Sect. Ecology, via Celoria 26, I-  
20133 Milano, Italy Fax: +39 2 26604361.  
Email: [rossaro@mailserver.unimi.it](mailto:rossaro@mailserver.unimi.it) **Regional  
representative for Italy**

LESLIE, Heather A., Department of Aquatic  
Ecology and Ecotoxicology, IBED, Faculty of  
Science, University of Amsterdam, Kruislaan  
320, P.O. Box 94084, 1090 GB, Amsterdam,  
The Netherlands Tel. +31 (0)20 525-7712.  
Fax: +31 (0)20 525-7716. Email:  
[leslie@science.uva.nl](mailto:leslie@science.uva.nl) **Regional  
representative for the Netherlands**

WILLASSEN, Endre, University of Bergen,  
Institute of Zoology, Museum of Zoology,  
Muséplass 3, N-5007 Bergen, Norway. Phone:  
+47 55582901. Fax: +47 55589677. Email:  
[Endre.Willassen@zmb.uib.no](mailto:Endre.Willassen@zmb.uib.no) **Regional  
representative for Norway**

KOWNACKI, Andrzej, Institute of Freshwater  
Biology, Academy of Sciences, Ul.  
Slawkowska 17, PL-31016 Krakow, Poland.  
Tel: +48 12 222115 **Regional representative  
for Poland**

HUGHES, Samantha, Laboratório Regionalde  
Engenharia Civil, Departamento de Recursos  
Naturais e de Hidaúlica, Caminho do  
Esmeraldo, 9000-264 Funchal, Portugal; and  
Centro de Estudos da Macaronesia,  
Universidade da Madeira, Campus da  
Penteada, 9000 Funchal, Portugal. E-mail:  
[samjhughes@hotmail.com](mailto:samjhughes@hotmail.com) **Regional  
representative for Portugal**

TUDORANCEA, Maria-Monica, Department  
of Ecology-Genetics, Faculty of Biology and

## **PACIFIC**

CRANSTON, Peter S., Department of  
Entomology, University of California, One  
Shields Avenue, Davis, CA 95616, US Email:  
[pscranston@ucdavis.edu](mailto:pscranston@ucdavis.edu) WWW:  
<http://entomology.ucdavis.edu/faculty/cranston.html> **Regional representative for Australia**

BOOTHROYD, Ian K.G., Kingett Mitchell &  
Associates, Level 2, ASDA Plaza, 4 Fred  
Thomas Drive, P.O. Box 33849, Takapuna,  
Auckland, New Zealand Phone: ++64 9

Geology, "Babes-Bolyai" University, 1 M.  
Kogalniceanu Str., Cluj-Napoca, 3400  
ROMANIA Fax: (40) (64) 191906 or 40 64  
431858. Email: [mtudor@biolog.ubbcluj.ro](mailto:mtudor@biolog.ubbcluj.ro)  
**Regional representative for Romania**

ZINCHENCO, Tatiana, Institute of Ecology of  
the Volga River Bassin, Russian Academy of  
Sciences, Togliatti 445003, Russia. Interests:  
Volga River Catchments and basin. Email:  
[tdz@mailru.com](mailto:tdz@mailru.com) **Regional representative for  
Togliatti**

BITUSIK, Peter, Faculty of Ecology and  
Environmental Sciences, Technical University,  
Kolpasska 9, SK-969 01 Banska Stiavnica,  
Slovakia. Email: [bitusik@fee.tuzvo.sk](mailto:bitusik@fee.tuzvo.sk)  
**Regional representative for Slovakia**

PRAT, Narcis, Dept Ecologia, Diagonal, 645  
Universitat de Barcelona, ES-08028  
Barcelona, Spain. Fax: (3)4111438. Email:  
[narcis@porthos.bio.ub.es](mailto:narcis@porthos.bio.ub.es) **Regional  
representative for Spain**

JOHNSON, Richard K., University of  
Agricultural Sciences, Box 7050, S 75007  
Uppsala, Sweden. Fax: +46 18 673156. Email:  
[Richard.Johnson@ma.slu.se](mailto:Richard.Johnson@ma.slu.se) **Regional  
representative for Sweden**

LODS-CROZET, Brigitte, Service des Eaux,  
Sols et Assainissement, Chemin des  
Boveresses 155, CH-1066 Epalinges,  
Switzerland. Tel: (+41 21) 316 71 87; Fax:  
(+41 21) 316 71 82. Email: [brigitte.lods-crozet@sesa.vd.ch](mailto:brigitte.lods-crozet@sesa.vd.ch) **Regional representative  
for Switzerland**

LANGTON, Peter H., 5 Kylebeg Avenue,  
Mountsandel, Coleraine, Co. Londonderry,  
Northern Ireland, BT52 1JN - Northern Ireland  
Email: [PHLangton@kylebegave.fsnet.co.uk](mailto:PHLangton@kylebegave.fsnet.co.uk)  
**Regional representative for UK**

4885033. Fax: ++64 9 4868072. Email: (work)  
[i Boothroyd@kma.co.nz](mailto:i Boothroyd@kma.co.nz) (home)  
[Ian.b@xtra.co.nz](mailto:Ian.b@xtra.co.nz) **Regional representative for  
New Zealand**

CATALAN, Zenaida Batac, Institute of  
Environmental Science and Management, The  
University of the Philippines at Los Banos,  
College, Laguna, Philippines. **Regional  
representative for Philippines**

## Current Bibliography: 1 Jan. 2003 - 31 Dec. 2003

by Odwin Hoffrichter

[© 2004 by CHIRONOMUS, and O. Hoffrichter - Associated Editor]

This listing is compiled, as usual, from many sources: databases, tables of contents of journals, references and citations of papers, autopsy of many periodicals, lists provided by authors (thanks to you!). Because titles of a particular year are not fully retrieved the following year, the current titles are preceded by supplements to the two preceding years. Only printed titles are reported here. Online publications should be retrieved elsewhere, in particular, check the chironomid home page for eventual references.

### Supplement to 2001 Current Bibliography: additions & corrections

- Albertoni, E. F., Palma-Silva, C. and Esteves, F. A. 2001a. Macroinvertebrates associated with *Chara* in a tropical coastal lagoon (Imboassica lagoon, Rio de Janeiro, Brazil). - *Hydrobiologia* 457: 215-224.
- Arrascue, A., Iannaccone, J., Alvarino, L., Basilio, S. y Lazcano, C. 2001a. El insecto *Chironomus calligraphus* Goeldi y la bacteria *Escherichia coli* como ensayos ecotoxicológicos para evaluar sedimentos elutriados dulceacuícolas. - *Revta per. Ent.* 42: 159-173.
- Bíró, K. és Specziár, A. 2001a. Adatok a Balaton árvaszúnyog (Diptera: Chironomidae) faunájához. (Contribution to the knowledge of Lake Balaton's chironomid fauna (Diptera, Chironomidae).) - *Hidrol. Közl.* 81: 322-325.
- Boyero, L. and Bailey, R. C. 2001a. Organization of macroinvertebrate communities at a hierarchy of spatial scales in a tropical stream. - *Hydrobiologia* 464: 219-225.
- Carter, J. L. and Fend, S. V. 2001a. Inter-annual changes in the benthic community structure of riffles and pools in reaches of contrasting gradient. - *Hydrobiologia* 459: 187-200.
- Czerniawska-Kusza, I. 2001a. Zmiany w zespole fauny dennej dolnego biegu Nysy Kłodzkiej pod wpływem czynników abiotycznych i antropogenicznych. (Changes in the bottom fauna community of the lower course of Nysa Kłodzka River under the influence of abiotic and anthropogenic factors.) - *Zesz. przyr. Opolsk. Tow. przyjac. Nauk* 35: 72-84.
- Dalton, C. P. D., Battarbee, R. W., Birks, H. J. B., Brooks, S. J., Cameron, N. G., Derrick, S., Evershed, R. P., McGovern, A., Peglar, S. M., Scott, J. A. and Thompson, R. 2001a. Climate history through the Holocene at Lochnagar, Scotland. - *Biol. Envir.* 101: 159.
- Funder, S., Bennike, O., Böcher, J., Israelson, C., Petersen, K. S. and Simonarson, L. A. 2001a. Late Pliocene Greenland - The Kap København Formation in North Greenland. - *Bull. geol. Soc. Denm.* 48: 117-134.
- Kaller, M. D., Hartman, K. J. and Angradi, T. R. 2001a. Experimental determination of benthic macroinvertebrate metric sensitivity to fine sediment in Appalachian streams. - *Proc. a. Conf. SEast. Ass. Fish Wildl. Ag.* 55:105-115.
- Lardicci, C., Como, S., Corti, S. and Rossi, F. 2001a. Recovery of the macrozoobenthic community after severe dystrophic crises in a Mediterranean coastal lagoon (Orbetello, Italy). - *Mar. Pollut. Bull.* 42: 202-214.
- Mastrantuono L. 2001b. Biomonitoring in the lake littoral using invertebrate fauna associated with sandy sediments: a short synthesis. - *Riv. Idrobiol.* 40: 115-128.
- Merritt, R. W. and Wallace, J. R. 2001a. The role of aquatic insects in forensic investigations. - In: Byrd, J. H. and Castner, J. L. (eds.): *Forensic entomology. The utility of arthropods in legal investigations*, pp. 177-222. CRC Pr., Boca Raton, Lond., N. Y., Wash.
- Nessimian, J. L. 2001a. Seasonal variation in the zoobenthic community of a sand dune marsh in Rio de Janeiro State, Brazil. - *Bolm Mus. nac., nov. Sér., Zool.* 447: 1-15.

- Österling, M. and Pihl, L. 2001a. Effects of filamentous green algal mats on benthic macrofaunal functional feeding groups. - *J. exp. mar. Biol. Ecol.* 263: 159-183.
- Ordóñez, A. J. A. 2001a. *Distribution of macrozoobenthos in Lake Scharmützel - a repetition of Wundsch's (1940) zoobenthos campaign of 1935.* - B. Sc. Thes., Tech. Univ. Cottbus. 41 pp.
- Pandourski, I. 2001a. Recherches hydrobiologiques des zones humides de la côte Bulgare de la mer Noire. I. Le lac de Vaja. - *Riv. Idrobiol.* 40: 321-334.
- Smith, H., Wood, P. J. and Gunn, J. 2001a. The macroinvertebrate communities of limestone springs in the Wye Valley, Derbyshire Peak District, UK. - *Cave Karst Sci.* 28: 67-78.
- Valdovinos, C. 2001a. Procesamiento de detritus ripariano por macroinvertebrados bentónicos en un estero boscoso de Chile central. - *Revta chil. Hist. nat.* 74: 445-453.
- Supplement to 2002 Current Bibliography:  
additions & corrections
- Afanas'ev, S. A. 2002a. Reaktsiya bioty gornykh rek na zalpovyе zagryazneniya. (Reaction of biota of mountain rivers on volley contamination.) - *Gidrobiol. Zh.* 38, 2: 42-50.
- Beaty, B. J., Mackie, R. S., Mattingly, K. S., Carlson, J. O. and Rayms-Keller, A. 2002a. The midgut epithelium of aquatic arthropods: A critical target organ in environmental toxicology. - *Envir. Hlth Perspect.* 110, Suppl. 6: 911-914.
- Boyero, L. and Bosch, J. 2002a. Spatial and temporal variation in macroinvertebrate drift in two neotropical streams. - *Biotropica* 34: 567-574.
- Camus, P. A. y Barahona, R. M. 2002a. Insectos del intermareal de Concepción, Chile: perspectivas para la investigación ecológica. - *Revta chil. Hist. nat.* 75: 793-803.
- Cobo, F., Soriano, O. y Báez, M. 2002a. Chironomidae Eretmopteridae Tendipedidae. - In: Carles-Tolrá, M. (ed.): *Catálogo de los Díptera de España, Portugal y Andorra (Insecta). Monogr. Soc. ent. aragon.* 8: 35-44.
- Cobo, F., Soriano, O. y González, M. A. (2001) 2002a. Inventario de los Quironómidos (Diptera Chironomidae) de Portugal. - *Nova Acta cient. compostelana Biol.* 11: 225-248.
- Corbi, J. J. and Trivinho-Strixino, S. 2002a. Spatial and bathymetric distribution of the macrobenthic fauna of the Ribeirão das Anhumas reservoir (Américo Brasiliense - SP, Brazil). - *Acta limnol. bras.* 14: 35-42.
- Dobson, M., Magana, A., Mathooko, J. M. and Ndegwa, F. K. 2002a. Detritivores in Kenyan highland streams: more evidence for the paucity of shredders in the tropics? - *Freshwat. Biol.* 47: 909-919.
- Edema, C. U., Ayeni, J. O. and Aruoture, A. 2002a. Some observations on the zooplankton and macrobenthos of the Okhuo River, Nigeria. - *J. aquat. Sci.* 17: 145-149.
- Eitam, A., Blaustein, L. and Mangel, M. 2002a. Effects of *Anisops sardea* (Hemiptera : Notonectidae) on oviposition habitat selection by mosquitoes and other dipterans and on community structure in artificial pools. - *Hydrobiologia* 485: 183-189.
- Elexová, E. 2002a. Changes in benthic communities along the Slovak stretch of the River Danube due to flow velocity gradient. - *Verh. int. Verein. Limnol.* 27: 279-282.
- Fleituch, T., Soszka, H., Kownacki, A. and Kudelska, D. 2002a. Sensitivity of macroinvertebrate metrics to detect pollution stress in Polish rivers. - In: Kownacki, A., Soszka, H., Fleituch, T. and Kudelska, D. (eds.): *River biomonitoring and benthic invertebrate communities*, pp. 63-70. Inst. Envir. Protect., Warsaw; Inst. Bot., Pol. Acad. Sci., Kraków.
- Franceschini, A. and Lencioni, V. 2002a. Lo zoobenthos (Capitolo 6). - In: Cantonati, M., Tolotti, M. e Lazzara, M. (eds.): *I laghi del Parco Adamello – Brenta. Ricerche limnologiche su laghi d'alta quota del settore siliceo del Parco.* - *Documenti del Parco* 14: 245-266.
- Gardarsson, A. and Einarsson, Á. 2002a. The food relations of waterbirds of Myvatn, Iceland. - *Verh. int. Verein. Limnol.* 28: 754-763.

- Harrison, A. D. (2001) 2002a. Chironomidae (Diptera) in the Albany Museum part 1. - *Ann. east. Cape Mus.* 2: 9-18.
- Hensel, E. C. K., Kirschbaum, F., Williot, P., Wirth, M. and Gessner, J. 2002a. Restoration of the European sturgeon, *Acipenser sturio* L., 1758 in Germany: Effect of different feed items on specific growth rates of large juvenile fish. - *Int. Rev. Hydrobiol.* 87: 539-551.
- Hillman, T. J. and Quinn, G. P. 2002a. Temporal changes in macroinvertebrate assemblages following experimental flooding in permanent and temporary wetlands in an Australian floodplain forest. - *River Res. Applic.* 18: 137-154.
- Hirabayashi, K. and Nakamoto, N. 2002a. Effects of sound traps on capture of chironomid midges near a hyper-eutrophic lake in urban area in Japan. - In: Jones, S. C., Zhai, J. and Robinson, W. H. (eds.): *Proc. 4<sup>th</sup> Int. Conf. Urban Pests*: 235-241. Coll. Charleston.
- Ingvason, H. R. 2002a. *The food-ecology of Tanytarsus gracilentus in Lake Myvatn.* - M. Sc. Thes., Univ. Iceland. 68 pp.
- Jónsson, I. R. og Malmquist, H. J. 2002a. Rannsóknir á Þrírhyrningsvatni 1998. (Investigations in Þrírhyrningsvatn 1998.) - *VMST-R/0202. Veiðimálastofnun* (= Institute of Freshwater Fisheries), Reykjavík. 17 pp.
- Karlson, K., Rosenberg, R. and Bonsdorff, E. 2002a. Temporal and spatial large-scale effects of eutrophication and oxygen deficiency on benthic fauna in Scandinavian and Baltic waters - a review. - *Oceanogr. mar. Biol. a. Rev.* 40: 427-489.
- Kim, C. K. and Cha, E. Y. 2002a. Applications of wavelet transform and artificial neural networks to pattern recognition for environmental monitoring. - In: McKay, B. and Slaney, J. (eds.): *AI 2002: Advances in Artificial Intelligence. Proc. 15<sup>th</sup> Aust. Jt Conf. artif. Intell. - Lect. Notes Comput. Sci.* 2557: 385-394.
- Klink, A. H., Moller Pillot, H. en Vallenduuk, H. 2002a. *Determinatiesleutel voor de larven van de in Nederland voorkomende soorten Polypedilum. (Identification key for the larvae of Polypedilum species occurring in the Netherlands.)* - StoWa, Utrecht. 17 pp.
- Kornijów, R. and Lachowska, G. 2002a. Effect of treated sewage on benthic invertebrate communities in the upland Bystrzyca Lubelska River (eastern Poland). - In: Kownacki, A., Soszka, H., Fleituch, T. and Kudelska, D. (eds.): *River biomonitoring and benthic invertebrate communities*, pp. 45-52. Inst. Envir. Protect., Warsaw; Inst. Bot., Pol. Acad. Sci., Kraków.
- Kownacki, A., Fleituch, T. and Dumnicka, E. 2002a. The effect of treated wastes on benthic invertebrate communities in the mountain zone of the Dunajec River (southern Poland). - In: Kownacki, A., Soszka, H., Fleituch, T. and Kudelska, D. (eds.): *River biomonitoring and benthic invertebrate communities*, pp. 29-43. Inst. Envir. Protect., Warsaw; Inst. Bot., Pol. Acad. Sci., Kraków.
- Kownacki, A., Soszka, H., Fleituch, T. and Kudelska, D. 2002a. The ecological assessment of river quality in Poland on the basis of communities of benthic invertebrates. - In: Kownacki, A., Soszka, H., Fleituch, T. and Kudelska, D. (eds.): *River biomonitoring and benthic invertebrate communities*, pp. 71-88. Inst. Envir. Protect., Warsaw; Inst. Bot., Pol. Acad. Sci., Kraków.
- Kuz'mina, Ya. S. 2002a. Raspredelenie lichinok khironomid v donnykh gruntakh rek zony Timanskogo kryazha (Respublika Komi). (Distribution of larvae of Chironomidae in bottom grounds of Timan ridge zone (Republic of Komi).) - *Gidrobiol. Zh.* 38, 4: 21-29.
- Lencioni, V., 2002a. I Chironomidi. - In: Lencioni, V. e Maiolini, B. (eds.): *L'ecologia di un ecosistema acquatico alpino (Val de la Mare, Parco Nazionale dello Stelvio)*. *Natura alpina* 54: 69-71.
- Lencioni, V., Maiolini, B., Zuccati, S. and Corradini, F. (2001) 2002a. Zoobenthos drift in two high mountain streams in the de la Mare glacial system (Stelvio National Park, Trentino, Italy). - In: Lencioni, V. and Maiolini, B. (eds.): *Atti del XV Convegno Gadio, Trento. Studi trent. Sci. nat., Acta biol.* 78/1: 49-57.

- Linevich, A. A., Makarchenko, E. A. i Aleksandrov, V. N. 2002a. *Khironomidy Baikala i Pribaikal'ya: Podonominae, Tanypodinae, Diamesinae, Prodiamesinae, Orthocladiinae: Annotirovannyi spisok vidov i form. (Chironomids of Lake Baikal and Pribaikal: Podonominae, Tanypodinae, Diamesinae, Prodiamesinae, Orthocladiinae: Annotated list of species and forms.)* - Nauka, Novosibirsk. 136 pp.
- Lyashenko, A. V. 2002a. Makrozoobentos urbanizirovannoi reki. (Macrozoobenthos of an urbanized river.) - *Gidrobiol. Zh.* 38, 6: 21-32.
- MacKenzie, R. A. and Kaster, J. L. 2002a. A preservative-free emergent trap for the isotopic and elemental analysis of emergent insects from a wetland system. - *Gt Lakes Ent.* 35: 47-51.
- Malmquist, H. J., Inginarsson, F., Jóhannsdóttir, E. E., Ólafsson, J. S. and Gíslason, G. M. 2002a. Zoobenthos in the littoral and profundal uzones of four Faroese lakes. - *Fróðskaparit* 50: 81-95.
- McKie, B. G. L. 2002a. *Multiscale abiotic, biotic and biogeographic influences on the ecology and distribution of lotic Chironomidae (Diptera) in the Australian Wet Tropics.* - Ph. D. Thes., James Cook Univ., Townsville.
- Merz, J. E. 2003b. Seasonal feeding habits, growth, and movement of steelhead trout in the lower Mokelumne River, California. - *Calif. Fish Game* 88: 95-111.
- Olander, H. 2002a. *Subfossil Chironomidae (Insecta: Diptera) as indicators of limnological and climatological conditions in Finland with particular focus on subarctic lakes.* - Ph. D. Thes., Univ. Helsinki.
- Percipalle, P., Jonsson, A., Nashchekin, D., Karlsson, C., Bergman, T., Guialis, A. and Daneholt, B. 2002a. Nuclear actin is associated with a specific subset of hnRNP A/B-type proteins. - *Nucleic Acids Res.* 30: 1725-1734.
- Porinchu, D. F. and MacDonald, G. M. (2001) 2002a. The modern distribution of midge flies (Chironomidae; Insecta: Diptera) in eastern Sierra Nevada, California, lakes: potential for paleoclimatic reconstruction. - In: West, G. J. and Buffaloe, L. D. (eds.): *Climate variability of the eastern North Pacific and Western North America. PACLIM. Proc. 18<sup>th</sup> a. Pacif. Clim. Workshop 2001. Tech. Rep. Interag. ecol. Progm S. Francisco Estuary*: 61-69.
- Pusch, M., Michels, U., Feld, C. K., Berger, T., Garcia, X.-F., Grünert, U. und Klausnitzer, B. 2002a. Benthische Wirbeltiere. - In: Köhler, J., Gelbrecht, J. und Pusch, M. (eds.): *Die Spree. Zustand, Probleme, Entwicklungsmöglichkeiten. Limnologie aktuell 10*: 166-185; (338)348-350.
- Rossaro, B., Lencioni, V. e Casalegno, C. (2001) 2002a. Relazioni tra specie di Ditteri Chironomidi e fattori ambientali esaminate con un data base relazionale. - In: Lencioni, V. e Maiolini, B. (eds.): *Atti del XV Convegno Gadio, Trento, 5-7 maggio 2001. Studi trent. Sci. nat., Acta biol.* 78/1: 201-206.
- Shevtsova, L. V., Yavorskii, V. Yu. i Omel'yanenko, D. I. 2002a. Zoobentos reki Desny, ust'evykh uchastkov nekotorykh ee pritokov i otsenka ikh ekologicheskogo sostoyania na territorii Ukrainy. (Zoobenthos of River Desna, mouth areas of some of its tributaries and assessment of their ecological state in the territory of the Ukraine.) - *Gidrobiol. Zh.* 38, 5: 3-16.
- Silaeva, A. A. i Protasov, A. A. 2002a. Zoobentos vodoema-okhladitelya Khmel'nitskoi AES i otsenka kachestva vody po donnym bespozvonochnym. (Zoobenthos of the cooling reservoir of the Khmel'nitsk NPP and evaluation of water quality according to bottom invertebrates.) - *Gidrobiol. Zh.* 38, 6: 46-59.
- Stańczykowska, A., Korycińska, M. and Królak, E. 2002a. The effect of treated wastewater on benthic invertebrate communities in the lowland Liwiec River (central Poland). - In: Kownacki, A., Soszka, H., Fleituch, T. and Kudelska, D. (eds.): *River biomonitoring and benthic invertebrate communities*, pp. 53-62. Inst. Envir. Protect., Warsaw; Inst. Bot., Pol. Acad. Sci., Kraków.
- Thiéry, A. and Puente, L. 2002a. Crustacean assemblage and environmental characteristics of a man-made solar saltwork in southern France, with emphasis on anostracan (Branchiopoda) population dynamics. - *Hydrobiologia* 486: 91-200.



- Wolff, R. H., Brasher, A. M. and Richards, A. B. 2002a. New generic records of Hawaiian Chironomidae (Diptera). - *Bishop Mus. occ. Pap.* 69: 31-33.
- World Health Organization 2002a. Review of: Deltamethrin 25% WG & WP and Agnique MMF. - *Report of the sixth WHOPEs [=WHO Pesticide Evaluation Scheme] working group meeting, WHO/HQ, Geneva 6-7 November 2002. WHO/CDS/WHOPEs/2002.6.* 56 pp.
- Yalynskaya, N. S., Oleksiv, I. T., Andrushchishin, O. P., Dumich, O. Ya., Edynak, O. P., Savitskaya, O. N. i Zabytivskii, Yu. M. 2002a. Hidrobiologicheskie indikatory toksifikatsii prudov Zapadnogo regiona Ukrainy. (Hydrobiological indicators of toxification of ponds of the Western Ukraine area.) - *Gidrobiol. Zh.* 38, 4: 57-70.
- Current Bibliography 2003**
- Åkerblom, N. and Goedkoop, W. 2003a. Stable isotopes and fatty acids reveal that *Chironomus riparius* feeds selectively on added food in standardized toxicity tests. - *Envir. Toxic. Chem.* 22: 1473-1480.
- Adams, C. E., Woltering, C. and Alexander, G. 2003a. Epigenetic regulation of trophic morphology through feeding behaviour in Arctic charr, *Salvelinus alpinus*. - *Biol. J. Linn. Soc.* 78: 43-49.
- Aguiaro, T., Branco, C. W. C., Verani, J. R. and Caramaschi, E. P. 2003a. Diet of the clupeid fish *Platanichthys platana* (Regan, 1917) in two different Brazilian coastal lagoons. - *Braz. Archs Biol. Technol.* 46: 215-222.
- Albertoni, E. F., Palma-Silva, C. and Esteves, F. A. 2003a. Natural diet of three species of shrimp in a tropical coastal lagoon. - *Braz. Archs Biol. Technol.* 46: 395-403.
- Alencar, Y. B., Ríos-Velásquez, C. M., Lichtwardt, R. W. and Hamada, N. 2003a. Trichomycetes (Zygomycota) in the digestive tract of arthropods in Amazonas, Brazil. - *Mems Inst. Oswaldo Cruz* 98: 799-810.
- Aleporou-Marinou, V., Marinou, H. and Patargias, T. 2003a. A mini review of the high mobility group proteins of insects. - *Biochem. Genet.* 41: 291-304.
- Amweg, E. L., Stuart, D. L. and Weston, D. P. 2003a. Comparative bioavailability of selenium to aquatic organisms after biological treatment of agricultural drainage water. - *Aquat. Toxic.* 63: 13-25.
- Andersson, J. 2003a. Effects of diet-induced resource polymorphism on performance in Arctic charr (*Salvelinus alpinus*). - *Evolut. Ecol. Res.* 5: 213-228.
- Andreev, A. A., Tarasov, P. E., Schwamborn, G., Ilyashuk, B. P., Ilyashuk, E. A., Bobrov, A. A., Stachura-Suchoples, K. and Hubberten, H.-W. 2003a. Holocene multi-proxy records from Nikolay Lake, Lena River Delta, arctic Russia. - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp., Espoo*: 149.
- Anwand, K., Valentin, M. and Mehner, T. 2003a. Species composition, growth and feeding ecology of fish community in Lake Stechlin - an overview. - *Arch. Hydrobiol., Spec. Iss. Advanc. Limnol.* 58: 237-246.
- Araújo-Coutinho, C. J. P. C., Cunha, A. B. P. V., Serra-Freire, N. M. and Mello, R. P. 2003a. Evaluation of the impact of *Bacillus thuringiensis* serovar *israelensis* and temephos, used for the control of *Simulium (Chirostilbia) pertinax* Kollar, 1832 (Diptera, Simuliidae) on the associated entomofauna, Paraty, State of Rio de Janeiro, Brazil. - *Mems Inst. Oswaldo Cruz* 98: 697-702.
- Armitage, P. D., Szoszkiewicz, K., Blackburn, J. H. and Nesbitt, I. 2003a. Ditch communities: a major contributor to floodplain biodiversity. - *Aquat. Conserv. mar. Freshwat. Ecosyst.* 13: 165-185.
- Arcott, D. B., Tockner, K. and Ward, J. V. 2003a. Spatio-temporal patterns of benthic invertebrates along the continuum of a braided Alpine river. - *Arch. Hydrobiol.* 158: 431-460.
- Artola-Garicano, E., Sinnige, T. L., van-Holsteijn, I., Vaes, W. H. J. and Hermens, J. L. M. 2003a. Bioconcentration and acute toxicity of polycyclic musks in two benthic organisms (*Chironomus riparius* and *Lumbriculus variegatus*). - *Envir. Toxic. Chem.* 22: 1086-1092.
- Back, R. C., Gorski, P. R., Cleckner, L. B. and Hurley, J. P. 2003a. Mercury content and speciation in the plankton and benthos of Lake Superior. - *Sci. tot. Envir.* 304: 349-354.



- Balci, P. and Kennedy, J. H. 2003a. Comparison of chironomids and other macroinvertebrates associated with *Myriophyllum spicatum* and *Heteranthera dubia*. - *J. Freshwat Ecol.* 18: 235-247.
- Balushkina, E. V. and Finogenova, N. P. 2003a. Changes in the benthic community structure and assessment of the water quality and the state of the ecosystems of Neva Bay and the eastern Gulf of Finland in 1994-2001. - *Proc. Eston. Acad. Sci. Biol. Ecol.* 52: 365-377.
- Barrett, S. D. and de Carvalho, C. R. 2003a. A software tool to straighten curved chromosome images. - *Chromosome Res.* 11: 83-88.
- Baumgärtner, D. and Rothaupt, K.-O. 2003a. Predictive length-dry mass regressions for freshwater invertebrates in a pre-Alpine lake littoral. - *Int. Rev. Hydrobiol.* 88: 453-483.
- Bazzanti, M., Della Bella, V. and Seminara, M. 2003a. Factors affecting macroinvertebrate communities in astatic ponds in central Italy. - *J. Freshwat. Ecol.* 18: 537-548.
- Bear, D. G., Fomproix, N., Soop, T., Björkroth, B., Masich, S. and Daneholt, B. 2003a. Nuclear poly(A)-binding protein PABPN1 is associated with RNA polymerase II during transcription and accompanies the released transcript to the nuclear pore. - *Expl Cell Res.* 286: 332-344.
- Beard, C. E. and Adler, P. H. 2003a. Zygospores of selected Trichomycetes in larval Diptera of the families Chironomidae and Simuliidae. - *Mycologia* 95: 317-320.
- Beckenbach, A. T. and Borkent, A. 2003a. Molecular analysis of the biting midges (Diptera: Ceratopogonidae), based on mitochondrial cytochrome oxidase subunit 2. - *Molec. Phylogenet. Evol.* 27: 21-35.
- Beier, S. and Traunspurger, W. 2003a. Temporal dynamics of meiofauna communities in two small submountain carbonate streams with different grain size. - *Hydrobiologia* 498: 107-131.
- Benbow, M. E., Burky, A. J. and Way, C. M. 2003a. Life cycle of a torrenticolous Hawaiian chironomid (*Telmatogeton torrenticola*): stream flow and microhabitat effects. - *Annls Limnol.* 39: 103-114.
- Berezina, N. A. 2003a. Rezistentnost' presnovodnykh bespozvonochnykh k izmeneniyu mineralizatsii vody. (Tolerance of freshwater invertebrates to changes in water salinity.) - *Ekologiya* 2003, 4: 296-301. [also published in: *Russ. J. Ecol.* 34: 261-266.]
- Bervoets, L., De Bruyn, L., Van Ginneken, L. and Blust, R. 2003a. Accumulation of <sup>137</sup>Cs by larvae of the midge *Chironomus riparius* from sediment: effect of potassium. - *Envir. Toxic. Chem.* 22: 1589-1596.
- Bettinetti, R., Giarei, C. and Provini, A. 2003a. Chemical analysis and sediment toxicity bioassays to assess the contamination of the River Lambro (Northern Italy). - *Archs envir. Contam. Toxic.* 45: 72-78.
- Bhattacharyya, S., Klerks, P. L. and Nyman, J. A. 2003a. Toxicity to freshwater organisms from oils and oil spill chemical treatments in laboratory microcosms. - *Envir. Pollut.* 122: 205-215.
- Biessmann, H. and Mason, J. M. 2003a. Telomerase-independent mechanisms of telomere elongation. - *Cell. molec. Life Sci.* 60: 2325-2333.
- Bigler, C., Grahn, E., Larocque, I., Jeziorski, A. and Hall, R. 2003a. Holocene environmental change at Lake Njulla (999 m a.s.l.), northern Sweden: a comparison with four small nearby lakes along an altitudinal gradient. - *J. Paleolimnol.* 29: 13-29.
- Bigrigg, J. U., Binkowski, F. P. and Bruch, R. 2003a. Food habit study of Lake Sturgeon, *Acipenser fulvescens*. - *Abstr. Aquat. Sci. Meet. 2003, Salt Lake City*, p. 27. Am. Soc. Limnol. Oceanogr.
- Björk, P., Baurén, G., Gelius, B., Wrangé, Ö. and Wieslander, L. 2003a. The *Chironomus tentans* translation initiation factor eIF4H is present in the nucleus but does not bind to mRNA until the mRNA reaches the cytoplasmic perinuclear region. - *J. Cell Sci.* 116: 4521-4532.
- Bojsen, B. H. and Jacobsen, D. 2003a. Effects of deforestation on macroinvertebrate diversity and assemblage structure in Ecuadorian Amazon streams. - *Arch. Hydrobiol.* 158: 317-342.
- Bond, N. R. and Downes, B. J. 2003a. The independent and interactive effects of fine sediment and flow on benthic invertebrate communities characteristic of small upland streams. - *Freshwat. Biol.* 48: 455-465.
- Bonsdorff, E., Laine, A. O., Hänninen, J., Vuorinen, I. & Norkko, A. 2003a. Zoobenthos

- of the outer archipelago waters (N. Baltic Sea) - the importance of local conditions for spatial distribution patterns. - *Boreal Envir. Res.* 8: 135-145.
- Boulton, A. J., Dole-Olivier, M.-J. and Marmonier, P. 2003a. Optimizing a sampling strategy for assessing hyporheic invertebrate biodiversity using the Bou-Rouch method: Within-site replication and sample volume. - *Arch. Hydrobiol.* 156: 431-456.
- Bowen, M. D. and Nelson, S. M. 2003a. Environmental variables associated with a chinook salmon redd in Deer Creek, California. - *Calif. Fish Game* 89: 176-186.
- Boyero, L. 2003a. Multiscale patterns of spatial variation in stream macroinvertebrate communities. - *Ecol. Res.* 18: 365-379.
- Boyero, L. 2003b. The effect of substrate texture on colonization by stream macroinvertebrates. - *Annls Limnol.* 39: 211-218.
- Boyle, T. P. and Fraleigh, H. D. Jr. 2003a. Natural and anthropogenic factors affecting the structure of the benthic macroinvertebrate community in an effluent-dominated reach of the Santa Cruz River, AZ. - *Ecol. Indicators* 3: 93-117.
- Brackenbury, J. 2003a. Swimming kinematics and wake elements in a worm-like insect: the larva of the midge *Chironomus plumosus* (Diptera). - *J. Zool.* 260: 195-201.
- Brady, L. and Cowell, B. C. 2003a. Colonization of fine particulate organic matter by invertebrates in a central Florida stream. - *Invert. Biol.* 122: 83-92.
- Brennan, A., Learner, M. A., Randerson, P. F. and Turk, R. 2003a. Chironomid (Diptera: Chironomidae) species assemblages and their ecology in the Wye river system. - *Arch. Hydrobiol. Suppl.* 139: 513-561.
- Brodersen, K. P. 2003a. Species-specific oxy-regulatory capacity of chironomids (Diptera) from a West Greenland training set. - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp., Espoo*: 9.
- Brodersen, K. P. and Bennike, O. 2003a. Interglacial Chironomidae (Diptera) from Thule, Northwest Greenland: matching modern analogues to fossil assemblages. - *Boreas* 32: 560-565.
- Brooks, B. W., Dzialowski, M. N., Turner, P. K., Stanley, J. K. and Glidewell, E. A. 2003a. Freshwater invertebrate responses to pharmaceuticals. - *Abstr. Aquat. Sci. Meet. 2003, Salt Lake City*, p. 31. Am. Soc. Limnol. Oceanogr.
- Brooks, B. W., Turner, P. K., Stanley, J. K., Weston, J. J., Glidewell, E. A., Foran, C. M., Slattery, M., La Point, T. W. and Huggett, D. B. 2003a. Waterborne and sediment toxicity of fluoxetine to select organisms. - *Chemosphere* 52: 135-142.
- Brooks, S. J. 2003a. Chironomid analysis to interpret and quantify Holocene climate change. - In: Mackay, A. W., Battarbee, R. W., Birks, H. J. B. and Oldfield, F. (eds.): *Global change in the Holocene*, pp. 328-341. Arnold, Lond.
- Brooks, S. J., Spiro, B. and Weiss, D. 2003a. Using chironomids to detect the impact of copper smelting on lakes in the southern Ural Mountains, Russia. - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp., Espoo*: 124.
- Brown, B. L. 2003a. Spatial heterogeneity reduces temporal variability in stream insect communities. - *Ecol. Lett.* 6: 316-325.
- Broza, M., Halpern, M., Gahanma, L. and Inbar, M. 2003a. Nuisance chironomids in waste water stabilization ponds: monitoring and action threshold assessment based on public complaints. - *J. Vector Ecol.* 28: 31-36.
- Burgherr, P., Klein, B., Robinson, C. T. and Tockner, K. 2003a. Surface zoobenthos. - In: Ward, J. V. and Uehlinger, U. (eds.): *Ecology of a glacial flood plain*, pp. 153-174. Kluwer Acad. Publs, Dordrecht, Boston, Lond.
- Burmeister, E.-G. und Reiss, F. † 2003a. *Rheotanytarsus reissi*, eine zweite tuffbildende Wasserinsektenart (Diptera, Chironomidae). - *Lauterbornia* 48: 77-88.
- Burt, J., Ciborowski, J. J. H. and Reynoldson, T. B. 2003a. Baseline incidence of mouthpart deformities in Chironomidae (Diptera) from the Laurentian Great Lakes, Canada. - *J. Gt Lakes Res.* 29: 172-180.
- Butcher, J. T., Stewart, P. M. and Simon, T. P. 2003a. A Benthic Community Index for streams in the Northern Lakes and Forests Ecoregion. - *Ecol. Indicators* 3: 181-193.

- Carew, M. E., Pettigrove, V. and Hoffmann, A. A. 2003a. Identifying chironomids (Diptera: Chironomidae) for biological monitoring with PCR-RFLP. - *Bull. ent. Res.* 93: 483-490.
- Caseldine, C., Geirsdóttir, Á. and Langdon, P. 2003a. Efstadalsvatn - a multi-proxy study of a Holocene lacustrine sequence from NW Iceland. - *J. Paleolimnol.* 30: 55-73.
- Castro, B. B., Guilhermino, L. and Ribeiro, R. 2003a. In situ bioassay chambers and procedures for assessment of sediment toxicity with *Chironomus riparius*. - *Envir. Pollut.* 125: 325-335.
- Cavalli, L., Pech, N. and Chappaz, R. 2003a. Diet and growth of the endangered *Zingel asper* in the Durance River. - *J. Fish Biol.* 63: 460-471.
- Chadwick, M. A. and Huryn, A. D. 2003a. Effect of a whole-catchment N addition on stream detritus processing. - *J. N. Am. benthol. Soc.* 22: 194-206.
- Chessman, B. C. 2003a. New sensitivity grades for Australian river macroinvertebrates. - *Mar. Freshwat. Res.* 54: 95-103.
- Chial, Z. B., Persoone, G. and Blaise, C. 2003a. Cyst-based toxicity tests XVI - sensitivity comparison of the solid phase *Heterocypris incongruens* microbioassay with the *Hyalella azteca* and *Chironomus riparius* contact assays on freshwater sediments from Peninsula Harbour (Ontario, Canada). - *Chemosphere* 52: 95-101.
- Chick, J. H., Cosgriff, R. J. and Gittinger, L. S. 2003a. Fish as potential dispersal agents for floodplain plants: first evidence in North America. - *Can. J. Fish. aquat. Sci.* 60: 1437-1439.
- Collier, K. J. and Bowman, E. J. 2003a. Role of wood in pumice-bed streams: I: Impacts of post-harvest management on water quality, habitat and benthic invertebrates. - *Forest Ecol. Mgmt* 177: 243-259.
- Collier, K. J. and Quinn, J. M. 2003a. Land-use influences macroinvertebrate community response following a pulse disturbance. - *Freshwat. Biol.* 48: 1462-1481.
- Collier, K. J. and Smith, B. J. 2003a. Role of wood in pumice-bed streams: II: Breakdown and colonisation. - *Forest Ecol. Mgmt* 177: 261-276.
- Contreras-Lichtenberg, R. 2003a. Berichtigung zur "Revision der westpaläarktischen Arten des Genus *Glyptotendipes* KIEFFER, 1913 (Insecta: Diptera, Nematocera, Chironomidae), Teil 2: Sg. *Glyptotendipes* s. str. KIEFFER, 1913 und Sg. *Trichotendipes* HEYN, 1993". - *Annl. naturhist. Mus. Wien* 104 B: 339-340.
- Cranston, P. S. 2003a. The oriental genus *Shangomyia* Sæther & Wang (Chironomidae: Diptera): immature stages, biology, putative relationships and the evolution of wood mining in chironomid larvae. - *Raffles Bull. Zool.* 51: 179-186.
- Crompton, B., Thomason, J. C. and McLachlan, A. 2003a. Mating in a viscous universe: the race is to the agile, not to the swift. - *Proc. R. Soc. Lond. B* 270: 1991-1995.
- Cross, W. F., Benstead, J. P., Rosemond, A. D. and Wallace, J. B. 2003a. Consumer-resource stoichiometry in detritus-based streams. - *Ecol. Letters* 6: 721-732.
- Croteau, M.-N., Hare, L. and Marcoux, P. 2003a. Feeding patterns of migratory and non-migratory fourth instar larvae of two coexisting *Chaoborus* species in an acidic and metal contaminated lake: Importance of prey ingestion rate in predicting metal bioaccumulation. - *Arch. Hydrobiol.* 158: 57-74.
- Culp, J. M., Cash, K. J., Glozier, N. E. and Brua, R. B. 2003a. Effects of pulp mill effluent on benthic assemblages in mesocosms along the Saint John River, Canada. - *Envir. Toxic. Chem.* 22: 2916-2925.
- Czerniawska-Kusza, I. 2003a. Biological water quality assessment of the Ścinawa Niemodlińska River based on the Biotic Index method. - *Arch. Ochr. Środ. [Arch. Envir. Protect.]* 29: 39-46.
- Davis, S., Golladay, S. W., Vellidis, G. and Pringle, C. M. 2003a. Macroinvertebrate biomonitoring in intermittent coastal plain streams impacted by animal agriculture. - *J. envir. Qual.* 32: 1036-1043.
- Death, R. G. 2003a. Spatial patterns in lotic invertebrate community composition: Is-substrate disturbance actually important? - *Can. J. Fish. aquat. Sci.* 60: 603-611.
- Death, R. G., Baillie, B. and Fransen, P. 2003a. Effect of *Pinus radiata* logging on stream invertebrate communities in Hawke's bay, New

- Zealand. - *N. Z. Jl mar. Freshwat. Res.* 37: 507-520.
- De Biasi, A. M., Benedetti-Cecchi, L., Pacciardi, L., Maggi, E., Vaselli, S. and Bertocci, I. 2003a. Spatial heterogeneity in the distribution of plants and benthic invertebrates in the lagoon of Orbetello (Italy). - *Oceanol. Acta* 26: 39-46.
- deBruyn, A. M. H., Marcogliese, D. J. and Rasmussen, J. B. 2003a. The role of sewage in a large river food web. - *Can. J. Fish. Aquat. Sci.* 60: 1332-1344.
- DeFoe, D. L. and Ankley, G. T. 2003a. Evaluation of time-to-effects as a basis for quantifying the toxicity of contaminated sediments. - *Chemosphere* 51: 1-5.
- Delettre, Y. R., Frenot, Y., Vernon, P. and Chown, S. L. 2003a. First record of *Telmatogeton* sp. (Diptera: Chironomidae) at Heard Island. - *Polar Biol.* 26: 423-426.
- Den Besten, P. J., Naber, A., Grootelaar, E. M. M. and van de Guchte, C. 2003a. In situ bioassays with *Chironomus riparius*: Laboratory-field comparisons of sediment toxicity and effects during wintering. - *Aquat. Ecosyst. Hlth Mgmt* 6: 217-228.
- Dettinger-Klemm, P.-M. A. 2003a. *Chironomids (Diptera, Nematocera) of temporary pools - an ecological case study.* - Ph. D. Thes., Univ. Marburg. 371 pp.
- Devin, S., Piscart, C., Beisel, J. N. and Moreteau, J. C. 2003a. Ecological traits of the amphipod invader *Dikerogammarus villosus* on a mesohabitat scale. - *Arch. Hydrobiol.* 158: 43-56.
- D'heygere, T., Goethals, P. L. M. and De Pauw, N. 2003a. Use of genetic algorithms to select input variables in decision tree models for the prediction of benthic macroinvertebrates. - *Ecol. Model.* 160: 291-300.
- Di Sabatino, A., Cicolani, B. and Gerecke, R. 2003a. Biodiversity and distribution of water mites (Acari, Hydrachnidia) in spring habitats. - *Freshwat. Biol.* 48: 2163-2173.
- Dörner, H., Berg, S., Jacobsen, L., Hülsmann, S., Brojerg, M. and Wagner, A. 2003a. The feeding behaviour of large perch *Perca fluviatilis* (L.) in relation to food availability: a comparative study. - *Hydrobiologia* 506-509: 427-434.
- Donald, D. B. and Anderson, R. S. 2003a. Resistance of the prey-to-predator ratio to environmental gradients and to biomanipulations. - *Ecology* 84: 2387-2394.
- Donohue, I., Verheyen, E. and Irvine, K. 2003a. *In situ* experiments on the effects of increased sediment loads on littoral rocky shore communities in Lake Tanganyika, East Africa. - *Freshwat. Biol.* 48: 1603-1616.
- Douglas, M. M. and O'Connor, R. A. 2003a. Effects of the exotic macrophyte, para grass (*Urochloa mutica*), on benthic and epiphytic macroinvertebrates of a tropical floodplain. - *Freshwat. Biol.* 48: 962-971.
- Drago, E. C., Ezcurra de Drago, I., Oliveros, O. B. and Paira, A. R. 2003a. Aquatic habitats, fish and invertebrate assemblages of the Middle Paraná River. - *Amazoniana* 17: 291-341.
- Duinen, G.-J. A. van, Brock, A. M.T., Kuper, J. T., Leuven, R. S. E. W., Peeters, T. M. J., Roelofs, J. G. M., Velde, G. van der, Verberk, W. C. E. P. and Esselink, H. 2003a. Do restoration measures rehabilitate fauna diversity in raised bogs? A comparative study on aquatic macroinvertebrates. - *Wetlands Ecol. Mgmt* 11: 447-459.
- Duxbury, C. 2003a. The use of previously colonized multiplate artificial substrates in experimental microcosms. - *J. Freshwat. Ecol.* 18: 459-464.
- Dyer, S. D., Peng, C., McAvoy, D. C., Fendinger, N. J., Masscheleyn, P., Castillo, L. V. and Lim, J. M. U. 2003a. The influence of untreated wastewater to aquatic communities in the Balatun River, The Philippines. - *Chemosphere* 52: 43-53.
- Edwards, D. D. and Smith, H. G. 2003a. Host sex preferences and transmission success by the water mite *Unionicola foili* (Acari: Unionicolidae) parasitic on the midge *Chironomus tentans* (Diptera : Chironomidae). - *J. Parasit.* 89: 681-685.
- Eggermont, H and Verschuren, D. 2003a. Impact of soil erosion in disturbed tributary drainages on the benthic invertebrate fauna of Lake Tanganyika, East Africa. - *Biol. Conserv.* 113: 99-109.
- Eggermont, H and Verschuren, D. 2003b. Subfossil Chironomidae from Lake Tanganyika, East Africa I. Tanypodinae and Orthocladiinae. - *J. Paleolimnol.* 29: 31-48.

- Eggermont, H and Verschuren, D. 2003c. Subfossil Chironomidae from Lake Tanganyika, East Africa. 2. Chironominae Chironomini and Tanytarsini). - *J. Paleolimnol.* 29: 423-457.
- Ekrem, T. 2003a. Towards a phylogeny of *Tanytarsus* van der Wulp (Diptera: Chironomidae). Is morphology alone sufficient to reconstruct the genealogical relationship? - *Insect Syst. Evol.* 34: 199-219.
- Ekrem, T., Sublette, M. F. and Sublette, J. E. 2003a. North American *Tanytarsus* I. Descriptions and keys to species in the *eminulus*, *gregarius*, *lugens* and *mendax* species groups (Diptera: Chironomidae). - *Ann. ent. Soc. Am.* 96: 265-328.
- Elexová, K. and Némethová, D. 2003a. The effect of abiotic environmental variables on the Danube macrozoobenthic communities. - *Limnologica* 33: 340-354.
- Elliott, J. M. 2003a. A comparative study of the functional response of four species of carnivorous stoneflies. - *Freshwat. Biol.* 48: 191-202.
- Erickson, B. E. 2003a. Small fiber estimates body residues. - *Envir. Sci. Technol.* 37: 56A-57A.
- Eriksson, L. 2003a. Dark/light side of *Heterotrissocladius*. - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp., Espoo*: 206.
- Falk, R., Agaton, C., Kiesler, E., Jin, S., Wieslander, L., Visa, N., Hober, S. and Ståhl, S. 2003a. An improved dual-expression concept, generating high-quality antibodies for proteomics research. - *Biotechnol. appl. Biochem.* 38: 231-239.
- Fargasova, A. 2003a. Cd, Cu, Zn, Al and their binary combinations acute toxicity for *Chironomus plumosus* larvae. - *Fresenius envir. Bull.* 12: 830-834.
- Ferrarese, U. 2003a. Chironomidi (Diptera: Chironomidae) di alcuni torrenti, laghi e zone umide del Trentino, con segnalazione di specie nuove per l'Italia. - *Studi trent. Sci. nat., Acta biol.* 79: 203-211.
- Ferrarese, U. and Lencioni, V. 2003a. *Acamptocladius reissi* Cranston & Sæther, 1982 (Diptera, Chironomidae): the first Italian records. - *Lav. Soc. venez. Sci. nat.* 28: 77-78.
- Ferreira-Peruquetti, P. e Trivinho-Strixino, S. 2003a. Notas sobre relações foréticas entre espécies de Chironomidae e Odonata do Estado de São Paulo, Brasil. - *Entomotropica* 18: 149-151.
- Fielding, N. J., MacNeil, C., Dick, J. T. A., Elwood, R. W., Riddell, G. E. and Dunn, A. M. 2003a. Effects of the acanthocephalan parasite *Echinorhynchus truttae* on the feeding ecology of *Gammarus pulex* (Crustacea: Amphipoda). - *J. Zool.* 261: 321-325.
- Fisher, T., Crane, M. and Callaghan, A. 2003a. Induction of cytochrome P-450 activity in individual *Chironomus riparius* Meigen larvae exposed to xenobiotics. - *Ecotoxic. envir. Saf.* 54: 1-6.
- Fleituch, T. 2003a. Structure and functional organization of benthic invertebrates in a regulated stream. - *Int. Rev. Hydrobiol.* 88: 332-344.
- Fochetti, R., Amici, I. and Argano, R. 2003a. Seasonal changes and selectivity in the diet of brown trout in the River Nera (Central Italy). - *J. Freshwat. Ecol.* 18: 437-444.
- Fore, L. S. 2003a. *Development and testing of invertebrate biomonitoring tools for Florida streams*. - Draft Rep., Fla Dep. envir. Protect., Tallahassee. 71 pp.
- Fowlkes, M. D., Michael, J. L., Crisman, T. L. and Prenger, J. P. 2003a. Effects of the herbicide imazapyr on benthic macroinvertebrates in a logged pond cypress dome. - *Envir. Toxic. Chem.* 22: 900-907.
- Foy, R. H., Lennox, S. D. and Gibson, C. E. 2003a. Changing perspectives on the importance of urban phosphorus inputs as the cause of nutrient enrichment in Lough Neagh. - *Sci. tot. Envir.* 310: 87-99.
- Frost Nerbonne, J. and Vondracek, B. 2003a. Volunteer macroinvertebrate monitoring: assessing training needs through examining error and bias in untrained volunteers. - *J. N. Am. benthol. Soc.* 22: 152-163.
- Frost, P. C., Tank, S. E., Turner, M. A. and Elser, J. J. 2003a. Elemental composition of littoral invertebrates from oligotrophic and eutrophic Canadian lakes. - *J. N. Am. benthol. Soc.* 22: 51-62.
- Fyodorova, M. V. and Azovsky, A. I. 2003a. Interactions between swarming *Chironomus annularius* (Diptera: Chironomidae) males:

- role of acoustic behavior. - *J. Insect Behav.* 16: 295-306.
- Galas, J. and Dumnicka, E. 2003a. Organic matter dynamics and invertebrate functional groups in a mountain stream in the West Tatra Mountains, Poland. - *Int. Rev. Hydrobiol.* 88: 362-371.
- Gaston, K. J., Jones, A. G., Hänel, C. and Chown, S. L. 2003a. Rates of species introduction to a remote oceanic island. - *Proc. R. Soc. Lond., Ser. B: Biol. Sci.* 270: 1091-1098.
- Gayraud, S. and Philippe, M. 2003a. Influence of bed-sediment features on the interstitial habitat available for macroinvertebrates in 15 French streams. - *Int. Rev. Hydrobiol.* 88: 77-93.
- Gido, K. B. 2003a. Effects of gizzard shad on benthic communities in reservoirs. - *J. Fish Biol.* 62: 1392-1404.
- Gilchrist, S. J. L., Brooks, S. J., Massaferro, J. and Metcalfe, S. E. 2003a. Late Quaternary palaeoclimatic reconstruction in Patagonia using chironomid analysis. - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp., Espoo*: 101.
- Gillespie, G. J. and Fox, M. G. 2003a. Morphological and life-history differentiation between littoral and pelagic forms of pumpkinseed. - *J. Fish Biol.* 62: 1099-1115.
- Girgin, S., N. Kazancı, N. and Dügel, M. 2003a. Ordination and classification of macroinvertebrates and environmental data of a stream in Turkey. - *Wat. Sci. Technol.* 47: 133-139.
- Goedkoop, W. and Peterson, M. 2003a. The fate, distribution, and toxicity of lindane in tests with *Chironomus riparius*: effects of bioturbation and sediment organic matter content. - *Envir. Toxic. Chem.* 22: 67-76.
- Golygina, V. V., Martin, J., Kiknadze, I. I., Siirin, M., Ivanchenko, O. V. and Makarchenko, E. A. 2003a. *Chironomus suwai*, a new species of the *plumosus* group (Diptera, Chironomidae) from Japan. - *Aquat. Insects* 25: 177-189.
- Gonçalves, J. F. Jr., Esteves, F. A. and Callisto, M. 2003a. Chironomids [sic!] colonization in *Nymphaea ampla* L. detritus during a degradative ecological succession experiment in a Brazilian coastal lagoon. - *Acta limnol. bras.* 15: 21-27.
- González Sagrario, M. A. and Balseiro, E. 2003a. Indirect enhancement of large zooplankton by consumption of predacious macroinvertebrates by littoral fish. - *Arch. Hydrobiol.* 158: 431-460.
- Gorab, E. 2003a. Reverse transcriptase-related proteins in telomeres and in certain chromosomal loci of *Rhynchosciara* (Diptera: Sciaridae). - *Chromosoma* 111: 445-454.
- Grasser, U. und Graf, W. 2003a. Erste Erfahrungen mit dem Potamon-Typie-Index (Schöll und Haybach 2001) in Österreich. - *Lauterbornia* 47: 153-172.
- Green, A. J. and Sánchez, M. I. 2003a. Spatial and temporal variation in the diet of Marbled Teal *Marmaronetta angustirostris* in the western Mediterranean: This globally threatened species is less dependent on invertebrates and more dependent on seeds than other ducks. - *Bird Study* 50: 153-160.
- Griffith, M. B., Husby, P., Hall, R. K., Kaufmann, P. R. and Hill, B. H. 2003a. Analysis of macroinvertebrate assemblages in relation to environmental gradients among lotic habitats of California's Central Valley. - *Envir. Monit. Assess.* 82: 281-309.
- Gunderina, L. I. i Salina, E. A. 2003a. Polimorfizm i divergentsiya multilokusnykh markerov DNK u vidov-dvoinikov *Chironomus riparius* Meigen i *Chironomus piger* Strenzke (Diptera, Chironomidae). - *Genetika* 39: 1059-1065. [also published as: Polymorphism and divergence of multilocus DNA markers in sibling species *Chironomus riparius* Meigen and *Chironomus piger* Strenzke (Diptera, Chironomidae).] - *Russ. J. Genet.* 39: 890-895.]
- Haapala, A., Muotka, T. and Laasonen, P. 2003a. Distribution of benthic macroinvertebrates and leaf litter in relation to streambed retentivity: implications for headwater stream restoration. - *Boreal Envir. Res.* 8: 19-30.
- Hämäläinen, H., Luotonen, H., Koskenniemi, E. and Liljaniemi, P. 2003a. Inter-annual variation in macroinvertebrate communities in a shallow forest lake in eastern Finland during 1990–2001. - *Hydrobiologia* 506-509: 389-397.
- Hall, D. L., Bergthold, B. S. and Sites, R. W. 2003a. The influence of adjacent land uses on macroinvertebrate communities of prairie

- streams in Missouri. - *J. Freshwat. Ecol.* 18: 55-68.
- Halpern, M., Gancz, H., Broza, M. and Kashi, Y. 2003a. *Vibrio cholerae* hemagglutinin/protease degrades chironomid egg masses. - *Appl. envir. Microbiol.* 69: 4200-4204.
- Hamada, N. and Oliveira S. J. de 2003a. Food items of larvae of *Rimanella arcana* (Needham, 1933) (Odonata: Amphipterygidae) in Central Amazonia, Brazil. - *Entomotropica* 18: 153-155.
- Harding, J. S. 2003a. Historic deforestation and the fate of endemic invertebrate species in streams. - *N. Z. J. mar. Freshwat. Res.* 37: 333-345.
- Harper, D. M., Childress, R. B., Harper, M. M., Boar, R. R., Hickley, P., Mills, S. C., Otieno, N., Drane, T., Vareschi, E., Nasirwa, O., Mwatha, W. E., Darlington, J. P. E. C. and Escuté-Gasulla, X. 2003a. Aquatic biodiversity and saline lakes: Lake Bogoria National Reserve, Kenya. - *Hydrobiologia* 500: 259-276.
- Hart, E. A. and Lovvorn, J. R. 2003a. Algal vs. macrophyte inputs to food webs of inland saline wetlands. - *Ecology* 84: 3317-3326.
- Hazra, N., Mazumdar, A. and Chaudhuri, P. K. 2003a. Life stages of four species of *Polypedilum* Kieffer (Diptera: Chironomidae) with a new species of the genus from Darjeeling-Sikkim Himalayas. - *Belg. J. Ent.* 5: 125-137.
- Hazra, N., Nath, S. and Chaudhuri, P. K. 2003a. The genus *Corynoneura* Winnertz (Dipt., Chironomidae) from the Darjeeling -Sikkim Himalayas of India, with descriptions of three new species. - *Entomologist's mon. Mag.* 139: 69-82.
- Heider, V. C. and Scharf, B. W. 2003a. Lake level fluctuations and temperature reconstruction: comparison of a deep and a shallow lake in southern Sweden - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp.*, Espoo: 190.
- Heino, J., Muotka, T., Paavola, R. and Paasivirta, L. 2003a. Among-taxon congruence in biodiversity patterns: can stream insect diversity be predicted using single taxonomic groups? - *Can. J. Fish. aquat. Sci.* 60: 1039-1049.
- Heiri, O. 2003a. Early human activity and chironomid-based temperature reconstruction: detecting anthropogenic artifacts in the records. - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp.*, Espoo: 19.
- Heiri, O. and Lotter, A. F. 2003a. 9000 years of chironomid assemblage dynamics in an Alpine lake: long-term trends, sensitivity to disturbance, and resilience of the fauna. - *J. Paleolimnol.* 30: 273-289.
- Heiri, O., Birks, H. J. B., Brooks, S. J., Velle, G. and Willassen, E. 2003a. Effects of within-lake variability of fossil assemblages on quantitative chironomid-inferred temperature reconstruction. - *Palaeogeogr. Palaeoclim. Palaeoecol.* 199: 95-106.
- Heiri, O., Lotter, A. F., Hausmann, S. and Kienast, F. 2003a. A chironomid-based Holocene summer air temperature reconstruction from the Swiss Alps. - *Holocene* 13: 477-484.
- Heiri, O., Wick, L., van Leeuwen, J. F. N., van der Knaap, W. O. and Lotter, A. F. 2003a. Holocene tree immigration and the chironomid fauna of a small Swiss subalpine lake (Hinterburgsee, 1515 m asl). - *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 189: 35-53.
- Henriques-Oliveira, A. L., Dorvillé, L. F. M. and Nessimian, J. L. 2003a. Distribution of Chironomidae larvae fauna (Insecta: Diptera) on different substrates in a stream at Floresta da Tijuca, RJ, Brazil. - *Acta limnol. bras.* 15: 69-84.
- Henriques Oliveira, A. L., Nessimian, J. L. and Dorvillé, L. F. M. 2003a. Feeding habits of chironomid larvae (Insecta: Diptera) from a stream in the Floresta da Tijuca, Rio de Janeiro, Brazil. - *Braz. J. Biol.* 63: 269-281.
- Hieber, M., Robinson, C. T. and Uehlinger, U. 2003a. Seasonal and diel patterns of invertebrate drift in different alpine stream types. - *Freshwat. Biol.* 48: 1078-1092.
- Hirabayashi, K., Hanazato, T. and Nakamoto, N. 2003a. Population dynamics of *Prosilocerus akamusi* and *Chironomus plumosus* (Diptera: Chironomidae) in Lake Suwa in relation to changes in the lake's environment. - *Hydrobiologia* 506-509: 381-388.
- Hirabayashi, K., Hanazato, T., Ogawara, M., Sakuma, M. and Nakamoto, N. 2003a. Long-term investigation of *Prosilocerus akamusi*

- (Tokunaga) (Diptera, Chironomidae) from a shallow eutrophic lake, Suwa, in Central Japan: An attempt to forecast the massive emergence of adult midges. - *Med. Ent. Zool.* 54: 89-96.
- Hirabayashi, K., Yamamoto, M., Takeda, M., Hanazato, T., and Nakamoto, N. 2003a. (Flight behavior of chironomid midges on the bank of Lake Suwa, Nagano Prefecture.) - *Pesutorogi Gatsukaishi [= Pest Control Res.]* 18: 91-101.
- Hirabayashi, K., Yamamoto, M., Yoshida, N. and Sasa, M. 2003a. Newly identified chironomids at Fuji Five Lakes in the fall season. - *Med. Ent. Zool.* 54: 389-394.
- Hoffsten, P.-O. 2003a. Effects of an extraordinarily harsh winter on macroinvertebrates and fish in boreal streams. - *Arch. Hydrobiol.* 157: 505-523.
- Holmen, J., Olsen, E. M. and Vøllestad, L. A. 2003a. Interspecific competition between stream-dwelling brown trout and Alpine bullhead. - *J. Fish Biol.* 62: 1312-1325.
- Holopainen, I. J., Holopainen, A.-L., Hämäläinen, H., Rahkola-Sorsa, M., Tkatcheva, V. and Viljanen, M. 2003a. Effects of mining industry waste waters on a shallow lake ecosystem in Karelia, north-west Russia. - *Hydrobiologia* 506-509: 111-119.
- Hooper, H. L., Sibly, R. M., Maund, S. J. and Hutchinson, T. H. 2003a. The joint effects of larval density and <sup>14</sup>C-cypermethrin on the life history and population growth rate of the midge *Chironomus riparius*. - *J. appl. Ecol.* 40: 1049-1059.
- Hooper, H. L., Sibly, R. M., Hutchinson, T. H. and Maund, S. J. 2003a. The influence of larval density, food availability and habitat longevity on the life history and population growth rate of the midge *Chironomus riparius*. - *Oikos* 102: 515-524.
- Hose, G. C., Lim, R. P., Hyne, R. V. and Pablo, F. 2003a. Short-term exposure to aqueous endosulfan affects macroinvertebrate assemblages. - *Ecotoxic. envir. Saf.* 56: 282-294.
- Hughes, S. J. 2003a. *A study of the freshwater macroinvertebrate fauna of Madeira and their application in a regional ecological monitoring system.* - Ph. D. Thes., King's Coll., Univ. London. 400 pp.
- Hunt, G. W. and Stanley, E. S. 2003a. Environmental factors influencing the composition and distribution of the hyporheic fauna in Oklahoma streams: Variation across ecoregions. - *Arch. Hydrobiol.* 158: 1-23.
- Hunter, J. G., Burger, J. and Cooper, K. R. 2003a. Use of an integrated mercury food web model for ecological risk assessment. - *J. envir. Sci. Hlth, Pt A: toxic/hazard. Subst. envir. Engng* 38: 1201-1214.
- Hwang, H., Fisher, S. W., Kim, K., Landrum, P. F., Larson, R. J. and Versteeg, D. J. 2003a. Assessing the toxicity of dodecylbenzene sulfonate to the midge *Chironomus riparius* using body residues as the dose metric. - *Envir. Toxic. Chem.* 22: 302-312.
- Hynynen, J. and Meriläinen, J. J. 2003a. Biological recovery from acidification inferred from palaeochironomid data and littoral benthos in forest lakes. - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp., Espoo*: 129.
- Iliopoulou-Georgudaki, J., Kantzaris, V., Katharios, P., Kaspiris, P., Georgiadis, T. and Montesantou, B. 2003a. An application of different bioindicators for assessing water quality: a case study in the rivers Alfeios and Pineios (Peloponnisos, Greece). - *Ecol. Indicators* 2: 345-360.
- Ilyashuk, B., Ilyashuk, E. and Dauvalter, V. 2003a. Chironomid responses to long-term metal contamination: a paleolimnological study in two bays of Lake Imandra, Kola Peninsula, northern Russia. - *J. Paleolimnol.* 30: 217-230.
- Ilyashuk, B. P., Ilyashuk, E. A., Andreev, A. A. B. and Hubberten, H.-W. 2003a. Post-glacial climate dynamics in East Siberian Arctic: palaeoclimatic and palaeolimnologic interpretation based on chironomid records. - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp., Espoo*: 58.
- Ilyashuk, E. A., Ilyashuk, B. P., Bennett, K. D. and Hammarlund, D. 2003a. Postglacial environmental changes inferred from midge (Diptera: Chironomidae, Chaoboridae and Ceratopogonidae) records in a lake on the southern Kola Peninsula, Northern Russia. - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp., Espoo*: 91.



- Iwakuma, T. and Gumiri, S. 2003a. The fate of free-swimming chironomid larvae in a humic, tropical oxbow lake. - *Verh. int. Verein. Limnol.* 28: 1494-1498.
- Jabłońska-Baran, I. i Łuczynski, M. 2003a. *Chironomus* f. l. *plumosus*, *Chironomus* sp. czy *Chironomus* spp. (*Chironomus* f. l. *plumosus*, *Chironomus* sp. or *Chironomus* spp.) - *Idee ekol.* 15, Ser. Szkice 8: 61.
- Jacobsen, D. 2003a. Altitudinal changes in diversity of macroinvertebrates from small streams in the Ecuadorian Andes. - *Arch. Hydrobiol.* 158: 145-167.
- Jakob, C., Robinson, C. T. and Uehlinger, U. 2003a. Longitudinal effects of experimental floods on stream benthos downstream from a large dam. - *Aquat. Sci.* 65: 223-231.
- Janssens de Bisthoven, L. and Gerhardt, A. 2003a. Chironomidae (Diptera, Nematocera) fauna in three small streams of Skania, Sweden. - *Envir. Monit. Assess.* 83: 89-102.
- Johnson, A. A. and Kleve, M. G. 2003a. *Strelkovimermis papillosus* n. sp. (Nematoda: Mermithidae), a parasite of chironomid (Insecta: Diptera) adults from the headwaters of the Mississippi River in Northern Minnesota. - *J. Parasit.* 89: 1186-1190.
- Johnson, B. R., Cross, W. F. and Wallace, J. B. 2003a. Long-term resource limitation reduces insect detritivore growth in a headwater stream. - *J. N. Am. benthol. Soc.* 22: 565-574.
- Johnson, L. B., Breneman, D. H. and Richards, C. 2003a. Macroinvertebrate community structure and function associated with large wood in low gradient streams. - *River Res. Appls* 19: 199-218.
- Johnson, R. K. 2003a. Development of a prediction system for lake stony-bottom littoral macroinvertebrate communities. - *Arch. Hydrobiol.* 158: 517-540.
- Johnson, Z. B. and Kennedy, J. H. 2003a. Macroinvertebrate assemblages of submerged woody debris in the Elm Fork of the Trinity River, Texas. - *J. Freshwat. Ecol.* 18: 187-197.
- Jónasson, P. M. 2003a. Hypolimnetic eutrophication of the N-limited dimictic L. Esrom 1908-1988. Pelagic-benthic coupling effects between phytoplankton and profundal zoobenthos, its growth, respiration and survival. - *Arch. Hydrobiol. Suppl.* 139: 449-512.
- Jones, A. G., Chown, S. L., Webb, T. J. and Gaston, K. J. 2003a. The free-living pterygote insects of Gough Island, South Atlantic Ocean. - *Syst. Biodivers.* 1: 213-273.
- Jones, N. E., Tonn, W. M. and Scrimgeour, G. J. 2003a. Selective feeding of age-0 Arctic grayling in lake-outlet streams of the Northwest Territories, Canada. - *Envir. Biol. Fishes* 67: 169-178.
- Jüttner, F. and Wessel, H. P. 2003a. Isolation of di(hydroxymethyl)dihydroxypyrrrolidine from the cyanobacterial genus *Cylindrospermum* that effectively inhibits digestive glucosidases of aquatic insects and crustacean grazers. - *J. Phycol.* 39: 26-32.
- Kajak, Z. and Prus, P. 2003a. Seasonal and year-to-year variation of numbers of *Chironomus plumosus* L. and Tubificidae in a lowland reservoir: Regularities, causes, mechanisms. - *Pol. J. Ecol.* 51: 339-351.
- Kaliszewicz, A. 2003a. Sublethal predation on *Stylaria lacustris*: a study of regenerative capabilities. - *Hydrobiologia* 501: 83-92.
- Kamimura, S., Matsuoka, A., Imai, K. and Shikama, K. 2003a. The swinging movement of the distal histidine residue and the autoxidation reaction for midge larval hemoglobins. - *Eur. J. Biochem.* 270: 1424-1433.
- Kangur, P., Kangur, A., Kangur, K. and Möls, T. 2003a. Condition and growth of ruffe *Gymnocephalus cernuus* (L.) in two large shallow lakes with different fish fauna and food recourse. - *Hydrobiologia* 506-509: 435-441.
- Karouna-Renier, N. K. and Zehr, J. P. 2003a. Short-term exposures to chronically toxic copper concentrations induce HSP70 proteins in midge larvae (*Chironomus tentans*). - *Sci. tot. Envir.* 312: 267-272.
- Karouna-Renier, N. K., Yang, W.-J. and Rao, K. R. 2003a. Cloning and characterization of a 70kDa heat shock cognate gene (HSC70) from two species of *Chironomus*. - *Insect molec. Biol.* 12: 19-26.
- Katano, O., Hosoya, K., Iguchi, K., Yamaguchi, M., Aonuma, Y. and Kitano, S. 2003a. Species diversity and abundance of freshwater fishes in irrigation ditches around rice fields. - *Envir. Biol. Fishes* 66: 107-121.

- Kawai, K. and Imabayashi, H. 2003a. Differences in conditions for collecting fertilized eggs in the laboratory among some Japanese chironomid species. - *Med. Ent. Zool.* 54: 125-131.
- Kawai, K., Kawai, T. and Imabayashi, H. 2003a. A comparison of improvemental ability of water quality among five chironomid species of the genus *Chironomus*. - *Med. Ent. Zool.* 54: 37-42.
- Kefford, B. J., Papas, P. J. and Nuggeoda, D. 2003a. Relative salinity tolerance of macroinvertebrates from the Barwon River, Victoria, Australia. - *Mar. Freshwat. Res.* 12: 755-765.
- Kehl, S. and Dettner, K. 2003a. Predation by pioneer water beetles (Coleoptera, Dytiscidae) from sandpit ponds, based on crop-content analysis and laboratory experiments. - *Arch. Hydrobiol.* 158: 109-126.
- Kelly, D. J., Bothwell, M. L. and Schindler, D. W. 2003a. Effects of solar ultraviolet radiation on stream benthic communities: an intersite comparison. - *Ecology* 84: 2724-2740.
- Khan, T. A. 2003a. Dietary studies on exotic carp (*Cyprinus carpio* L.) from two lakes of western Victoria, Australia. - *Aquat. Sci.* 65: 272-286.
- Khan, T. A. 2003b. Limnology of four saline lakes in western Victoria, Australia. II. Biological parameters. - *Limnologica* 33: 327-339.
- Kiesler, E., Miralles, F., Östlund Farrants, A.-K. and Visa, N. 2003a. The Hrp65 self-interaction is mediated by an evolutionarily conserved domain and is required for nuclear import of Hrp65 isoforms that lack a nuclear localization signal. - *J. Cell Sci.* 116: 3949-3956.
- Kiffney, P. M., Richardson, J. S. and Bull, J. P. 2003a. Responses of periphyton and insects to experimental manipulation of riparian buffer width along forest streams. - *J. appl. Ecol.* 40: 1060-1076.
- Kiiski, A., Hämäläinen, H., Salo, S. and Verta, M. 2003a. A paleolimnological analysis of changes in midge community structure and incidence of larval deformities in River Kymijoki in response to industrial pollution. - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp., Espoo*: 127.
- Kiknadze, I. I., Istomina, A. G., Makarchenko, E. A., Katokhin, A. V. i Golygina, V. V. 2003a. Kariotip i khromosomnyi polimorfizm khironomidy *Chironomus yoshimatsui* (Diptera, Chironomidae). (Karyotype and chromosomal polymorphysm [sic!] in the midge *Chironomus yoshimatsui* (Diptera, Chironomidae [sic!])). - *Zool. Zh.* 82: 1215-1221.
- Kleeberg, A. 2003a. Re-assessment of Wundsch's (1940) 'H<sub>2</sub>S-Oscillatoria-Lake' type using the eutrophic Lake Scharmützel (Brandenburg, NE Germany) as an example. - *Hydrobiologia* 501: 1-5.
- Klemetsen, A., Knudsen, R., Staldvik, F. J. and Amundsen, P.-A. 2003a. Habitat, diet and food assimilation of Arctic charr under the winter ice in two subarctic lakes. - *J. Fish Biol.* 62: 1082-1098.
- Knispel, S. and Castella, E. 2003a. Disruption of a longitudinal pattern in environmental factors and benthic fauna by a glacial tributary. - *Freshwat. Biol.* 48: 604-618.
- Kobayashi, T. 2003a. New record of the genus in Tanypodinae (Chironomidae) from Japan, *Hayesomyia tripunctata* (Goetghebuer, 1922). - *Med. Ent. Zool.* 54: 173-176.
- Kobayashi, T., Ohtaka, A. and Takahashi, T. 2003a. The second record of ectoparasitic Chironomidae on Trichoptera from Japan, *Polypedilum (Cerobregma) kamotertium* Sasa, 1989 (Insecta, Diptera, Chironomidae, Chironomini). - *Spixiana* 26: 83-91.
- Koehler, P. G. 2003a. Blind mosquitoes (aquatic midges). - *ENY-231*. Univ. Fla Ext, Inst. Food agric. Sci., Gainesville. 5 pp.
- Kone, T. and Teugels, G. G. 2003a. Food habits of brackish water tilapia *Sarotherodon melanotheron* in riverine and lacustrine environments of a West African coastal basin. - *Hydrobiologia* 490: 75-85.
- Kornijów, R., Radwan, S., Tarkowska-Kukuryk, M. and Kahlan, G. 2003a. Zoobenthos of ecotonal zones in several lakes of different trophic status (the region Polesie Lubelskie, Eastern Poland). - *Pol. J. Ecol.* 51: 237-246.
- Kownacki, A. 2003a. Stan i perspektywy badań zoobentosu w rzekach. (Status and perspectives of zoobenthos investigations in rivers.) - *Idee ekol.* 15, Ser. Szkiec 8: 25-35.

- Kravtsova, L. S., Karabanov, E. B., Kamaltynov, R. M., Mekhanikova, I. V., Sitnikova, T. Ya., Rozhkova, N. A., Slugina, Z. V., Izhboldina, L. A., Weinberg, L. V., Akinshina, T. V., Krivonogov, S. K. i Shcherbakov, D. Yu. 2003a. Makrozoobentos subakval'nykh landshaftov melkovodnoi zony Yuzhnogo Baikala. 1. Lokal'noe raznoobrazie donnogo naseleniya i osobennosti ego prostranstvennogo raspredeleniya. (Macrozoobenthos of subaquatic landscapes in shoal of Southern Baikal. 1. Local diversity of bottom population and peculiarities of its spatial distribution.) - *Zool. Zh.* 82: 307-317.
- Krawchuk, M. A. and Taylor, P. D. 2003a. Changing importance of habitat structure across multiple spatial scales for three species of insects. - *Oikos* 103: 153-161.
- Kutschera, U. 2003a. The feeding strategies of the leech *Erpobdella octoculata* (L.): a laboratory study. - *Int. Rev. Hydrobiol.* 88: 94-101.
- Kuz'mina, Ya. S., Shilova, A. I. i Zelentsov, N. I. 2003a. Fauna khironomid (Diptera; Chironomidae) rek Timanskogo kryazha. (Midge (Diptera, Chironomidae) fauna of the Timanskii Range rivers.) - *Ent. Obozr.* 52: 590-597; 796.
- Lahr, J., Maas-Diepeveen, J. L., Stuijzand, S. C., Leonards, P. E. G., Drüke, J. M., Lücker, S., Espeldoorn, A., Kerkum, L. C. M., van Stee, L. L. P. and Hendriks, A. J. 2003a. Responses in sediment bioassays used in the Netherlands: can observed toxicity be explained by routinely monitored priority pollutants? - *Wat. Res.* 37: 1691-1710.
- Laine, A. O., Luodekari, K., Poikonen, M. and Viitasalo, M. 2003a. A comparison between 1928 and 2000 indicates major changes in the macrozoobenthos species composition and abundance on the SW coast of Finland (Baltic Sea). - *Proc. Eston. Acad. Sci. Biol. Ecol.* 52: 3-16.
- Lamy-Enrici, M.-H., Dondeyne, A. and Thybaud, E. 2003a. Influence of the organic matter on the bioavailability of phenanthrene for benthic organisms. - *Aquat. Ecosyst. Hlth Mgmt* 6: 391-396.
- Lang, B., Bedford, A. P., Richardson, N. and Brooks, S. J. 2003a. The use of ultra-sound in the preparation of carbonate and clay sediments for chironomid analysis. - *J. Paleolimnol.* 30: 451-460.
- Langdon, P. G., Barber, K. E. and Lomas-Clarke, S. H. 2003a. Reconstructing climate and environmental change in northern England: evidence from Talkin Tarn, Cumbria. - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp., Espoo*: 93.
- LaPerriere, J. D. 2003a. Limnology of Harding Lake, Alaska: a deep, subarctic lake. - *Lake Reservoir Mgmt* 19: 93-107.
- Larocque, I. and Hall, R. I. 2003a. Chironomids as quantitative indicators of mean July air temperature: validation by comparison with century-long meteorological records from northern Sweden. - *J. Paleolimnol.* 29: 475-493.
- Lastra, C. C. L., Mazzucchelli, M. G. and Dikgolz, V. 2003a. Temporal changes in the prevalence of three species of *Trichomyces* (Zygomycota : Zygomycotina) in Dipteran aquatic larvae from Argentina. - *Fungal Diversity* 14: 85-93.
- Legssyer, B., Chergui, H. and Maamri, A. 2003a. Invertebrate dynamics during the decomposition of dry and fresh willow leaves in Oued Zegzel (Eastern Morocco). - *Annlis Limnol.* 39: 27-33.
- Lepori, F., Barbieri, A. and Ormerod, S. J. 2003a. Effects of episodic acidification on macroinvertebrate assemblages in Swiss Alpine streams. - *Freshwat. Biol.* 48: 1873-1885.
- Leppä, M., Hämäläinen, H. and Karjalainen, J. 2003a. The response of benthic macroinvertebrates to whole-lake biomanipulation. - *Hydrobiologia* 498: 97-105.
- Leppänen, M. T., Landrum, P. F., Kukkonen, J. K. V., Greenberg, M. S., Burton, G. A Jr., Robinson, S. D. and Gossiaux, D. C. 2003a. Investigating the role of desorption on the bioavailability of sediment-associated 3,4,3,4-tetrachlorobiphenyl in benthic invertebrates. - *Envir. Toxic. Chem.* 22: 2861-2871.
- Lin, H.-J., Shao, K.-T., Chiou, W.-L., Maa, C.-J. W., Hsieh, H.-L., Wu, W.-L., Severinghaus, L. L. and Wang, Y.-T. 2003a. Biotic communities of freshwater marshes and mangroves in relation to saltwater incursions: implications for wetland regulation. - *Biodivers. Conserv.* 12: 647-665.

- López L., C., Mazzucchelli, M. G. and Dikgolz, V. 2003a. Temporal changes in the prevalence of three species of Trichomycetes (Zygomycota: Zygomycotina) in dipteran aquatic larvae from Argentina. - *Fungal Divers. 14*: 85-93.
- Lüder, B., Brooks, S. J. and Birks, H. J. B. 2003a. Quantitative Holocene temperature reconstruction from Lake Reiersdalsvatnet, southern Norway, based on fossil chironomids: preliminary results. - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp., Espoo*: 193.
- Lukin, A., Dauvalter, V., Kashulin, N., Yakovlev, V., Sharov, A. and Vandyshev O. 2003a. Assessment of copper-nickel industry impact on a subarctic lake ecosystem. - *Sci. tot. Envir. 306*: 73-83.
- Lyytikäinen, M., Rantalainen, A.-L., Mikkelsen, P., Hämäläinen, H., Paasivirta, J. and Kukkonen, J. V. K. 2003a. Similarities in bioaccumulation patterns of polychlorinated dibenzo-*p*-dioxins and furans and polychlorinated diphenyl ethers in laboratory-exposed oligochaetes and semipermeable membrane devices and in field-collected chironomids. - *Envir. Toxic. Chem. 22*: 2405-2415.
- MacNeil, C., Bigsby, E., Dick, J. T. A., Hynes, H. B. N., Hatcher, M. J. and Dunn, A. M. 2003a. Temporal changes in the distribution of native and introduced freshwater amphipods in Lough Neagh, Northern Ireland. - *Arch. Hydrobiol. 157*: 379-395.
- Mäenpää, K. A., Sormunen, A. J. and Kukkonen, J. V. K. 2003a. Bioaccumulation and toxicity of sediment associated herbicides (ioxynil, pendimethalin, and bentazone) in *Lumbriculus variegatus* (Oligochaeta) and *Chironomus riparius* (Insecta). - *Ecotoxic. envir. Saf. 56*: 398-410.
- Maazono, Y. and Miyashita, T. 2003a. Community-level impacts induced by introduced largemouth bass and bluegill in farm ponds in Japan. - *Biol. Conserv. 109*: 111-121.
- Makarchenko, E. A. 2003a. Novyi vid *Psectrocladius* Kieffer (Diptera, Chironomidae, Orthocladiinae) s yuga rossiiskogo Dal'nego Vostoka. (A new species of *Psectrocladius* Kieffer (Diptera, Chironomidae, Orthocladiinae) from the south of Russian Far East.) - *Evroaziat. ent. Zh. 2*: 61-66.
- Makarchenko, E. A. i Makarchenko, M. A. 2003a. Novyi i maloizvestnyi vidy *Stilocladius* Rossaro, 1979 (Diptera, Chironomidae, Orthocladiinae) s rossiiskogo Dal'nego Vostoka. (A new and little known species of *Stilocladius* Rossaro, 1979 (Diptera, Chironomidae, Orthocladiinae) from the Russian Far East.) - *Evroaziat. ent. Zh. 2*: 135-140.
- Makarchenko, E. A. i Makarchenko, M. A. 2003b. Novye i maloizvestnye vidy khironomid (Diptera, Chironomidae) rossiiskogo Dal'nego Vostoka. (A new and little known species of chironomids (Diptera, Chironomidae) from the Russian Far East.) - In: *Chteniya pamyati Vladimira Yakovlevicha Levanidova (Vladimir Ya. Levanidov's Biennial Memorial Meetings) 2*: 204-217. Dal'nauka, Vladivostok.
- Makarchenko, E. A. i Makarchenko, M. A. 2003c. Novyi i maloizvestnyi vidy komarov-zvontsov (Diptera, Chironomidae) iz Yuzhnogo Primor'ya (Dal'nii Vostok Rossii). (New and little known species of non-biting midges (Diptera, Chironomidae) from South Primorye (Russian Far East).) - *Evroaziat. ent. Zh. 2*: 215-219.
- Malard, F. 2003a. Interstitial fauna. - In: Ward, J. V. and Uehlinger, U. (eds.): *Ecology of a glacial flood plain*, pp. 175-198. Kluwer Acad. Publs, Dordrecht, Boston, Lond.
- Malard, F., Ferreira, D., Dolédec, S. and Ward, J. V. 2003a. Influence of groundwater upwelling on the distribution of the hyporheos in a headwater river flood plain. - *Arch. Hydrobiol. 157*: 89-116.
- Malard, F., Galassi, D., Lafont, M., Dolédec, S. and Ward, J. V. 2003a. Longitudinal patterns of invertebrates in the hyporheic zone of a glacial river. - *Freshwat. Biol. 48*: 1709-1725.
- March, J. G., Benstead, J. P., Pringle, C. M and Luckymis, M. 2003a. Benthic community structure and invertebrate drift in a Pacific island stream, Kosrae, Micronesia. - *Biotropica 35*: 125-130.
- Marcogliese, D. J., Rasmussen, J. B. and deBruyn, A. M. H. 2003a. The role of sewage in a large river food web. - *Can. J. Fish. aquat. Sci. 60*: 1332-1344.

- Marqués, M. J., Martínez-Conde, E. and Rovira, J. V. 2003a. Effects of zinc and lead mining on the benthic macroinvertebrates of a fluvial ecosystem. - *Wat. Air Soil Pollut.* 148: 363-388.
- Martin, J., Guryev, V., Macdonald, S. S., Blinov, A. and Edwards, D. H. D. 2003a. Phylogenetic relationships of *Archaeochlus* Brundin, *Austrochilus* Cranston and *Afrochilus* Freeman (Diptera: Chironomidae), basal genera with a Gondwanan connection. - *Cimbebasia* 19: 193-203.
- Martinez, E. A., Moore, B. C., Schaumloffel, J. and Dasgupta, N. 2003a. Morphological abnormalities in *Chironomus tentans* exposed to cadmium- and copper-spiked sediments. - *Ecotoxic. envir. Saf.* 55: 204-212.
- Massaferro, J., Brooks, S. J. and Haberle, S. 2003a. The dynamics of vegetation and chironomid assemblages during the Late Quaternary at Laguna Facil, Chonos Archipelago, southern Chile. - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp., Espoo*: 168.
- Mastrantuono, L. and Mancinelli, T. 2003a. Meio-macroinvertebrates associated with submerged plants in a *Chara*-lake (Lake Martignano, Italy): benthic structure and environmental quality. - *Atti Ass. Ital. Oceanol. Limnol.* 16: 187-201.
- Maximov, A. A. 2003a. Changes of the bottom macrofauna in the eastern Gulf of Finland in 1985-2002. - *Proc. Eston. Acad. Sci. Biol. Ecol.* 52: 378-393.
- Maxted, J. R., Evans, B. F. and Scarsbrook, M. R. 2003a. Development of standard protocols for macroinvertebrate assessment of soft-bottomed streams in New Zealand. - *N. Z. Jl mar. Freshwat. Res.* 37: 793-807.
- Maycock, D. S., Prenner, M. M., Kheir, R., Morris, S., Callaghan, A., Whitehouse, P., Morrill, D. and Crane, M. 2003a. Incorporation of in situ and biomarker assays in higher-tier assessment of the aquatic toxicity of insecticides. - *Wat. Res.* 37: 4180-4190.
- McCabe, D. J. and Gotelli, N. J. 2003a. Caddisfly diapause aggregations facilitate benthic invertebrate colonization. - *J. Anim. Ecol.* 72: 1015-1026.
- McCarthy, D. 2003a. The trans-Pacific zipper effect: disjunct sister taxa and matching geological outlines that link the Pacific margins. - *J. Biogeogr.* 30: 1545-1561.
- McLachlan, A., Ladle, R. and Crompton, B. 2003a. Predator-prey interactions on the wing: aerobatics and body size among dance flies and midges. - *Anim. Behav.* 66: 911-915.
- Mendes, H. F., Marcondes, C. B. and De Pinho, L. C. 2003a. A new phytotelmatic species of *Monopelopia* Fittkau, 1962 (Insecta: Diptera: Chironomidae: Tanypodinae) from South Brazil. - *Zootaxa* 262: 1-10.
- Menéndez, M., Hernández, O. and Comín, F. A. 2003a. Seasonal comparisons of leaf processing rates in two Mediterranean rivers with different nutrient availability. - *Hydrobiologia* 495: 159-169.
- Meriläinen, J. J., Hynynen, J., Palomäki, A., Mäntykoski, K. and Witick, A. 2003a. Environmental history of an urban lake: a palaeolimnological study of Lake Jyväsjärvi, Finland. - *J. Paleolimnol.* 30: 387-406.
- Mermillod-Blondin, F., Gaudet, J.-P., Gérino, M., Desrosiers, G. and Creuzé des Châtelliers, M. 2003a. Influence of macroinvertebrates on physico-chemical and microbial processes in hyporheic sediments. - *Hydrol. Proces.* 17: 779-794.
- Metcalf-Smith, J. L., Holtze, K. E., Sirota, G. R., Reid, J. J. and de Solla, S. R. 2003a. Toxicity of aqueous and sediment-associated fluoride to freshwater organisms. - *Envir. Toxic. Chem.* 22: 161-166.
- Metzeling, L., Chessman, B., Hardwick, R. and Wong, V. 2003a. Rapid assessment of rivers using macroinvertebrates: the role of experience, and comparisons with quantitative methods. - *Hydrobiologia* 510: 39-52.
- Meyer, E. I. and Poepperl, R. 2003a. Secondary production of invertebrates in a Central European mountain stream (Steina, Black Forest, Germany). - *Arch. Hydrobiol.* 158: 25-42.
- Michailova, P., Ilkova, J. and White, K. N. 2003a. Functional and structural rearrangements of salivary gland polytene chromosomes of *Chironomus riparius* Mg. (Diptera, Chironomidae) in response to freshly neutralized aluminium. - *Envir. Pollut.* 123: 193-207.
- Michailova, P. V., Ilkova, J. and White, K. 2003b. Cytogenetic alterations in

- Prodiamesinae species (Diptera, Chironomidae) from different polluted regions. - *Folia biol., Kraków* 51: 69-79.
- Michiels, S. 2003a. *Pseudorthocladius berthelemyi*, eine für Österreich neu nachgewiesene Zuckmückenart (Diptera, Chironomidae). - *Lauterbornia* 48: 89-90.
- Michiels, S. 2003b. *Eukiefferiella ancyla* Svensson 1986, eine für Deutschland neu nachgewiesene Zuckmückenart (Diptera: Chironomidae) aus dem Schwarzwald. - *Mitt. bad. Landesver. Naturk. Natursch. N. F.* 18: 221-222.
- Milani, D., Reynoldson, T. B., Borgmann, U. and Kolasa, J. 2003a. The relative sensitivity of four benthic invertebrates to metals in spiked-sediment exposures and application to contaminated field sediment. - *Envir. Toxic. Chem.* 22: 845-854.
- Millet, L. et Verneaux, V. 2003a. Evolution des assemblages de Chironomidae (Insecta: Diptera) pendant le Tardiglaciaire dans le Lac du Lautrey (Jura, France): essai de reconstruction paléoenvironnementale. - *Eclogae geol. Helv.* 96, Suppl.: S99-S107.
- Millet, L., Verneaux, V. and Magny, M. 2003a. Lateglacial paleoenvironmental reconstruction using subfossil chironomid assemblages from Lake Lautrey (Jura, France). - *Arch. Hydrobiol.* 156: 405-429.
- Miserendino, M. L. and Pizzolon, L. A. 2003a. Distribution of macroinvertebrate assemblages in the Azul-Quemquemtreu river basin, Patagonia, Argentina. - *N. Z. Jl mar. Freshwat. Res.* 37: 525-539.
- Mischke, S., Herzsuh, U. and Zhang, C. 2003a. A Central Asiatic chironomid and ostracod record of the past 45,000 years. - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp., Espoo*: 73.
- Mistak, J. L., Hayes, D. B. and Bremigan, M. T. 2003a. Food habits of coexisting salmonines above and below Stronach Dam in the Pine River, Michigan. - *Envir. Biol. Fishes* 67: 179-190.
- Miyake, Y., Hiura, T., Kuhara, N. and Nakano, S. 2003a. Succession in a stream invertebrate community: a transition in species dominance through colonization. - *Ecol. Res.* 18: 493-501.
- Mörtl, M. and Rothhaupt, K.-O. 2003a. Effects of adult *Dreissena polymorpha* on settling juveniles and associated macroinvertebrates. - *Int. Rev. Hydrobiol.* 88: 561-569.
- Moller Pillot, H. 2003a. *Hoe waterdieren zich handhaven in een dynamisch wereld. 10 jaar onderzoek in de Roodloop, een bovenloopje van de Reusel in Noord-Brabant. (The survival of aquatic animals in a dynamic world. A ten-year study [in] the Roodloop, a small tributary of the Reusel in the Dutch Province of Noord-Brabant.)* - Sticht. Noordbrabants Landschap, Haaren. 182 pp. (+ CD-ROM).
- Moore, B. C., Martinez, E., Gay, J. M. and Rice, D. H. 2003a. Survival of *Salmonella enterica* in freshwater and sediments and transmission by the aquatic midge *Chironomus tentans* (Chironomidae: Diptera). - *Appl. envir. Microbiol.* 69: 4556-4560.
- Moretti, M. S., Goulart, M. D. C. e Callisto, M. 2003a. Avaliação rápida da macrofauna associada a *Eichhornia azurea* (Swartz) Kunth, 1843 e *Pontederia lanceolata* Nutt., 1818 (Pontederiaceae) na Baía do Coqueiro, Pantanal de Poconé (MT/Brasil). - *Revta bras. Zoociênc.* 5: 7-21.
- Morse, C. C., Huryn, A. D. and Cronan, C. 2003a. Impervious surface area as a predictor of the effects of urbanization on stream insect communities in Maine, U.S.A. - *Envir. Monit. Assess.* 89: 95-127.
- Mosch, E. C. and Scharf, B. W. 2003a. Paleolimnology of Lake Tiefer See (Mecklenburg-Vorpommern, NE Germany) based on remains of Chironomidae (Diptera). - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp., Espoo*: 113.
- Mousavi, S. K., Primicerio, R. and Amundsen, P.-A. 2003a. Diversity and structure of Chironomidae (Diptera) communities along a gradient of heavy metal contamination in a subarctic watercourse. - *Sci. tot. Envir.* 307: 93-110.
- Mucha, A. P., Vasconcelos, M. T. S. D. and Bordalo, A. A. 2003a. Macrobenthic community in the Douro estuary: relations with trace metals and natural sediment characteristics. - *Envir. Pollut.* 121: 169-180.
- Muotka, T., Heino, J., Paavola, R. and Paasivirta, L. 2003a. Among-taxon congruence in biodiversity patterns: can stream insect

- diversity be predicted using single taxonomic groups? - *Can. J. Fish. aquat. Sci.* 60: 1039-1049.
- Murray, D. A and Murray, W. A. 2003a. A reassessment of Chironomidae (Diptera) of Clare Island, Co Mayo, with first records of *Acamptocladius reissi* Cranston & Sæther and *Limnophyes angelicae* Sæther (Orthocladinae) for the Irish faunal checklist. - *Bull. Ir. biogeogr. Soc.* 27: 255-269.
- Nalepa, T. F., Fanslow, D. L., Lansing, M. B. and Lang, G. A. 2003a. Trends in the benthic macroinvertebrate community of Saginaw Bay, Lake Huron, 1987 to 1996: responses to phosphorus abatement and the zebra mussel, *Dreissena polymorpha*. - *J. Gt Lakes Res.* 29: 14-33.
- Narita, T., Kondo, S., Sugimoto, A., Kiyashko, S. I. and Wada, E. 2003a. Chironomid larvae eating methane. - *Geochim. cosmochim. Acta* 67, 18, Suppl. 1: A332.
- Nazarova, L. B. and Brooks, S. 2003a. Response of Chironomidae to recent environmental change in Northern Russia. - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp., Espoo*: 151.
- Negishi, J. N. and Richardson, J. S. 2003a. Responses of organic matter and macroinvertebrates to placements of boulder clusters in a small stream of southwestern British Columbia, Canada. - *Can. J. Fish. aquat. Sci.* 60: 247-258.
- Nelson, S. M. and Roline, R. A. 2003a. Effects of multiple stressors on hyporheic invertebrates in a lotic system. - *Ecol. Indicators* 3: 65-79.
- Nessimian, J. L., Amorim, R. M., Henriques-Oliveira, A. L. e Sanseverino, A. M. 2003a. Chironomidae (Diptera) do Estado do Rio de Janeiro: levantamento dos gêneros e hábitos de ocorrência. - *Publ. avuls. Mus. nac., Rio de Janeiro* 98: 1-16.
- Neumann, M., Baumeister, J., Liess, M. and Schulz, R. 2003a. An expert system to estimate the pesticide contamination of small streams using benthic macroinvertebrates as bioindicators. II. The knowledge base of LIMPACT. - *Ecol. Indicators* 2: 391-401.
- Niinioja, R., Holopainen, A.-L., Hämäläinen, H., Heitto, L., Luotonen, H., Mononen, P. and Rämö, A. 2003a. State of Lake Sysmäjärvi, Eastern Finland, after loading with mine water and municipal waste water for several decades. - *Hydrobiologia* 506-509: 773-780.
- Niituma, H. 2003a. Redescription of the larva of *Larsia miyagasensis* (Diptera, Chironomidae). - *Jap. J. syst. Ent.* 9: 43-45.
- Niituma, H. 2003b. First record of *Brundiniella* (Insecta: Diptera: Chironomidae) from the Palaearctic Region, with the description of a new species. - *Species Diversity* 8: 293-300.
- Nilsson, T., Spörting, S. and Björklund, E. 2003a. Selective supercritical fluid extraction to estimate the fraction of PCB that is bioavailable to a benthic organism in a naturally contaminated sediment. - *Chemosphere* 53: 1049-1052.
- Nonnis Marzano, C., Scalera Liaci, L., Fianchini, A., Gravina, F., Mercurio, M. and Corriero, G. 2003a. Distribution, persistence and change in the macrobenthos of the lagoon of Lesina (Apulia, southern Adriatic Sea). - *Oceanol. Acta* 26: 57-66.
- Nyström, P., McIntosh, A. R. and Winterbourn, M. J. 2003a. Top-down and bottom-up processes in grassland and forested streams. - *Oecologia* 136: 596-608.
- O'Brien, K., Hänninen, J., Kanerva, T., Metsärinne, L. and Vuorinen, I. 2003a. Macrozoobenthic zonation in relation to major environmental factors across the Archipelago Sea, northern Baltic Sea. - *Boreal Envir. Res.* 8: 159-170.
- O'Connell, M. T. 2003a. Direct exploitation of prey on an inundated floodplain by cherryfin shiners (*Lythrurus roseipinnis*) in a low order, blackwater stream. - *Copeia* 2003: 635-645.
- Okolelov, A. Y. i Shubin, A. O. 2003a. (Effects of environmental-factors on numbers and distribution of waders (Charadriiformes, Charadrii) in anthropogenic landscapes of the Oka-Don plain.) - *Zool. Zh.* 82: 388-401.
- Olsen, A., Bale, J. S., Leadbeater, B. S. C., Callow, M. E. and Holden, J. B. 2003a. Developmental thresholds and day-degree requirements of *Paratanytarsus grimmii* and *Corynoneura scutellata* (Diptera: Chironomidae): two midges associated with potable water treatment. - *Physiol. Ent.* 28: 315-322.
- Olson, N. W., Paukert, C. P., Willis, D. W. and Klammer, J. A. 2003a. Prey selection and diets of bluegill *Lepomis macrochirus* with differing

- population characteristics in two Nebraska natural lakes. - *Fish. Mgmt Ecol.* 10: 31-40.
- Opsahl, R. W., Wellnitz, T. and Poff, N. L. 2003a. Current velocity and invertebrate grazing regulate stream algae: results of an *in situ* electrical exclusion. - *Hydrobiologia* 499: 135-145.
- Orendt, C. 2003a. A classification of semi-natural northern prealpine river stretches based on chironomid communities. - *Annls Limnol.* 39: 219-237.
- Orendt, C. und Reiff, N. 2003a. Rote Liste gefährdeter Zuckmücken (Diptera: Chironomidae) Bayerns. - In: Bayer. Landesamt Umweltsch. (ed.): *Rote Liste gefährdeter Tiere Bayerns. SchrReihe 166*: 301-304.
- Ortí, F., Rosell, L. and Anadón, P. 2003a. Deep to shallow lacustrine evaporites in the Libros Gypsum (southern Teruel Basin, Miocene, NE Spain): an occurrence of pelletal gypsum rhythmites. - *Sedimentology* 50: 361-386.
- Owens, R. W. and Dittman, D. E. 2003a. Shifts in the diets of slimy sculpin (*Cottus cognatus*) and lake whitefish (*Coregonus clupeaformis*) in Lake Ontario following the collapse of the burrowing amphipod *Diporeia*. - *Aquat. Ecosyst. Hlth Mgmt* 6: 311-323.
- Paggi, A. C. and Rodrigues Capítulo, A. 2003a. Chironomid composition from drift and bottom samples in a regulated north-Patagonian river (Rio Limay, Argentina). - *Verh. int. Verein. Limnol.* 28: 1229-1235.
- Palaczyk, A. i Klasa, A. 2003a. Muchówki (Diptera) masywu Babiej Góry. (Diptera of Babia Góra Massif.) - In: Woloszyn, B. W., Woloszyn, D. i Celary, D. (eds): *Monografia fauny Babiej Góry*, pp. 305-357. Kom. Ochr. Przyr. PAN, Kraków.
- Palmer, C. G., Williams, M. L. and Gordon, A. K. 2003a. Riverine macroinvertebrate responses to chlorine and chlorinated sewage effluents - Community structure in the Umsunduze and Umbilo Rivers, KwaZulu-Natal, South Africa. - *Water SA* 29: 473-481.
- Papoucheva, E., Proviz, V., Lambkin, C., Goddeeris, B. and Blinov, A. 2003a. Phylogeny of the endemic Baikalian *Sergentia* (Chironomidae, Diptera). - *Molec. Phylogenet. Evol.* 29: 120-125.
- Pascoal, C., Pinho, M., Cassio, F. and Gomes, P. 2003a. Assessing structural and functional ecosystem condition using leaf breakdown: studies on a polluted river. - *Freshwat. Biol.* 48: 2033-2044.
- Pennuto, C. M. 2003a. Population dynamics and intraspecific interactions of an ectosymbiotic midge in a river in southern Maine, USA. - *J. N. Am. benthol. Soc.* 22: 249-262.
- Percipalle, P., Fomproix, N., Kylberg, K., Miralles, F., Björkroth, B., Daneholt, B. and Visa, N. 2003a. An actin-ribonucleoprotein interaction is involved in transcription by RNA polymerase II. - *Proc. natn. Acad. Sci. U. S. Am.* 100: 6475-6480.
- Péry, A. R. R., Ducrot, V., Mons, R. and Garric, J. 2003a. Modelling toxicity and mode of action of chemicals to analyse growth and emergence tests with the midge *Chironomus riparius*. - *Aquat. Toxic.* 65: 281-292.
- Péry, A. R. R., Ducrot, V., Mons, R., Miège, C., Gahou, J., Gorini, D. and Garric, J. 2003a. Survival tests with *Chironomus riparius* exposed to spiked sediments can profit from DEBtox model. - *Wat. Res.* 37: 2691-2699.
- Péry, A. R.R., Sulmon, V. Mons, R., Flammarion, P., Lagadic, L. and Garric, J. 2003a. A model to understand the confounding effects of natural sediments in toxicity tests with *Chironomus riparius*. - *Envir. Toxic. Chem.* 22: 2476-2481.
- Perry, R. W., Bradford, M. J. and Grout, J. A. 2003a. Effects of disturbance on contribution of energy sources to growth of juvenile chinook salmon (*Oncorhynchus tshawytscha*) in boreal streams. - *Can. J. Fish. aquat. Sci.* 60: 390-400.
- Petänen, T., Lyytikäinen, M., Lappalainen, J., Romantschuk, M. and Kukkonen, J. V. K. 2003a. Assessing sediment toxicity and arsenite concentration with bacterial and traditional methods. - *Envir. Pollut.* 122: 407-415.
- Peters, A. J., Armitage, P. D., Everett, S. J. and House, W. A. 2003a. Control of nuisance chironomid midge swarms from a slow sand filter. - *J. Wat. Supply Res. Technol.* - *AQUA* 52: 109-118.
- Petersen, F. T. and Meier, R. 2003a. Testing species-richness estimation methods on single-sample collection data using the Danish Diptera. - *Biodivers. Conserv.* 12: 667-686.



- Petrova, N. A. i Rakisheva, A. 2003a. Kariotip i morfologiya *Chironomus anthracinus* Zett. (Diptera, Chironomidae) iz Vostochnogo Kazakhstana. (Karyotype and morphology of the larvae of *Chironomus anthracinus* Zett. (Diptera, Chironomidae) from eastern Kazakhstan.) - *Tsitologiya* 45: 428-433.
- Petrova, N. A., Zelentsov, N. I., Klishko, O. K. i Chubareva, L. A. 2003a. Pervoopisanie politennykh khromosom, morfologiya lichinok i biologiya dvukh vidov roda *Prosilocerus* (Diptera, Chironomidae, Orthoclaadiinae). (First description of polytene chromosomes, larval morphology and biology of two species of genus *Prosilocerus* (Diptera, Chironomidae, Orthoclaadiinae).) - *Trudy russk. ent. Obshch.* 74: 33-50.
- Phillips, E. C. 2003a. Habitat preference of aquatic macroinvertebrates in an east Texas sandy stream. - *J. Freshwat. Ecol.* 18: 1-11.
- Phillips, E. C., Washek, M. E., Hertel, A. W. and Niebel, B. M. 2003a. The round goby (*Neogobius melanostomus*) in Pennsylvania tributary streams of Lake Erie. - *J. Gt Lakes Res.* 29: 34-40.
- Plant, W., Ciborowski, J. J. H. and Corkum, L. D. 2003a. Do tube-dwelling midges inhibit the establishment of burrowing mayflies? - *J. Gt Lakes Res.* 29: 521-528.
- Poinar, G. O. Jr and Poinar, R. 2003a. Description and development of *Gastromermis anisotis* sp. n. (Nematoda: Mermithidae), a parasite in a quadritrophic system involving a cyanobacterium, midge and virus. - *Nematology* 5: 325-338.
- Polukonova, N. V. 2003a. Samki komarov-zvontsov roda *Chironomus* Mg. (Diptera, Chironomidae). II. *Chironomus balatonicus* Devai et al., *Ch. muratensis* Ryser et al., *Ch. nudiventris* Ryser et al. i *Ch. entis* Shobanov iz gruppy the *plumosus*. (Females of the midge genus *Chironomus* Mg. (Diptera, Chironomidae). II. *Chironomus balatonicus* Devai et al., *Ch. muratensis* Ryser et al., *Ch. nudiventris* Ryser et al., and *Ch. entis* Shobanov of the *plumosus* group.) - *Ent. Obozr.* 82: 487-499, 531.
- Ponte, I., Vila, R. and Suau, P. 2003a. Sequence complexity of histone H1 subtypes. - *Molec. Biol. Evol.* 20: 371-380.
- Porinchu, D. F. and MacDonald, G. M. 2003a. The use and application of freshwater midges (Chironomidae: Insecta: Diptera) in geographical research. - *Prog. phys. Geogr.* 27: 378-422.
- Porinchu, D. F., MacDonald, G. M., Bloom, A. M. and Moser, K. A. 2003a. Late Pleistocene and early Holocene climate and limnological changes in the Sierra Nevada, California, USA inferred from midges (Insecta: Diptera: Chironomidae). - *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 198: 403-422.
- Prchalová, M., Draštík, V., Kubečka, J., Sricharoendham, B., Schiemer, F. and Vijverberg, J. 2003a. Acoustic study of fish and invertebrate behavior in a tropical reservoir. - *Aquat. liv. Resourc.* 16: 325-331.
- Quinlan, R., Douglas, M. S. V. and Smol, J. P. 2003a. Recent climate warming in high Arctic (Canada) ponds and lakes: subfossil chironomid evidence. - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp., Espoo*: 65.
- Quinlan, R., Paterson, A. M., Hall, R. I., Dillon, P. J., Wilkinson, A. N., Cumming, B. F., Douglas, M. S. V. and Smol, J. P. 2003a. A landscape approach to examining spatial patterns of limnological variables and long-term environmental change in a southern Canadian lake district. - *Freshwat. Biol.* 48: 1676-1697.
- Quinn, M. R., Feng, X., L. Folt, C. L. and Chamberlain, C. P. 2003a. Analyzing trophic transfer of metals in stream food webs using nitrogen isotopes. - *Sci. tot. Envir.* 317: 73-89.
- Rand, G. M., Wheat, J. V., Carriger, J. F. and Lee, T. A. 2003a. An automated overlying water-renewal system for sediment toxicity studies. - *Envir. Pollut.* 122: 169-175.
- Reed, T. 2003a. Macroinvertebrate assemblage change in a small Eastern Oregon stream following disturbance by grazing cattle. - *J. Freshwat. Ecol.* 18: 315-319.
- Reich, P. and Downes, B. J. 2003a. The distribution of aquatic invertebrate egg masses in relation to physical characteristics of oviposition sites at two Victorian upland streams. - *Freshwat. Biol.* 48: 1497-1513.
- Reshetnikov, A. N. 2003a. The introduced fish, rotan (*Perccottus glenii*), depresses populations of aquatic animals (macroinvertebrates, amphibians, and a fish). - *Hydrobiologia* 510: 83-90.

- Rikardsen, A. H., Amundsen, P. A. and Bodin, P. J. 2003a. Growth and diet of anadromous Arctic charr after their return to freshwater. - *Ecol. Freshwat. Fish* 12: 74-80.
- Robinson, C. T. and Gessner, M. O. 2003a. Litter decomposition. - In: Ward, J. V. and Uehlinger, U. (eds.): *Ecology of a glacial flood plain*, pp. 217-230. Kluwer Acad. Publs, Dordrecht, Boston, Lond.
- Robinson, C. T., Uehlinger, U. and Monaghan, M. T. 2003a. Effects of a multi-year experimental flood regime on macroinvertebrates downstream of a reservoir. - *Aquat. Sci.* 65: 210-222.
- Rodrigues Capítulo, A. and Paggi, A. C. 2003a. Zoobenthic communities in relation to slope, substrate heterogeneity and urban disturbances in Pampean hill streams ( Argentina). - *Verh. int. Verein. Limnol.* 28: 1267-1273.
- Romanuk, T. N. and Levings, C. D. 2003a. Associations between arthropods and the supralittoral ecotone: dependence of aquatic and terrestrial taxa on riparian vegetation. - *Envir. Ent.* 32: 1343-1353.
- Roque, F. O. and Trivinho-Strixino, S. 2003a. *Guassutanypus oliveirai*, a new genus and species of Macropelopiini from Brazil (Insecta, Diptera, Chironomidae). - *Spixiana* 26: 159-164.
- Roque, F. O., Corbi, J. J. and Trivinho-Strixino, S. 2003a. Macroinvertebrates on different technosubstrates in a stream of an urban area of São Carlos - SP, Brazil. - *Multiciência* 5: 172-177.
- Roque, F. O., Pepinelli, M., Fragoso, E. N., Ferreira, W. A., Barillari, P. R., Yoshinaga, M. Y., Strixino, S. T., Verani, N. F. e Lima, M. I. S. 2003a. Ecología de macroinvertebrados, peixes e vegetação ripária de um córrego de primeira ordem em região de Cerrado do Estado de São Paulo (São Carlos, SP). - In: Raoul, H. (ed.): *Ecótonos nas interfaces dos ecossistemas aquáticos*, pp. 313-337. São Carlos, Rima.
- Roque, F. O., Trivinho-Strixino, S., Strixino, G., Agostinho, R. C. and Fogo, J. C. 2003a. Benthic macroinvertebrates in streams of the Jaragua State Park (Southeast of Brazil) considering multiple spatial scales. - *J. Insect Conserv.* 7: 63-72.
- Roque, F. O., Trivinho-Strixino, S., Yamada, M. V. and Penteado-Dias, A. M. 2003a. A preliminary survey of Chironomidae adults (Insecta: Diptera) in a heterogeneous landscape of Jaraguá State Park, São Paulo, Brazil. - *Biotemas* 16: 67-78.
- Rosén, P., Segerström, U., Eriksson, L. and Renberg, I. 2003a. Do diatom, chironomid, and pollen records consistently infer Holocene July air temperature? A comparison using sediment cores from four alpine lakes in northern Sweden. - *Arct. Antarct. alp. Res.* 35: 279-290.
- Ross, S. W. and Rohde, F. C. 2003a. Life history of the swampfish from a North Carolina stream. - *SEast. Nat.* 2: 105-120.
- Rossaro, B., Casalegno, C. e Lencioni, V. 2003a. Contributo alla revisione del sottogenere *Orthocladius* s. str. van der Wulp (Diptera: Chironomidae): distribuzione ed autoecologia delle specie note in Italia. - In: Baldacchini, G. N. e Sansoni, G. (eds.): *Atti Sem. Studi "Nuovi orizzonti dell'ecologia"*, pp. 329-333. Prov. Auton. Trento, Ag. Prov. Protez. Amb. Trento, Centro Ital. Studi Biol. Amb. Trento.
- Rossaro, B., Lencioni, V. and Casalegno, C. (2002) 2003a. Revision of West Palaearctic species of *Orthocladius* s. str. van der Wulp, 1874 (Diptera: Chironomidae: Orthoclaadiinae), with a new key to species. - *Studi trent. Sci. nat., Acta biol.* 79: 213-241.
- Rowe, D. K., Dean, T. L., Williams, E. and Smith, J. P. 2003a. Effects of turbidity on the ability of juvenile rainbow trout, *Oncorhynchus mykiss*, to feed on limnetic and benthic prey in laboratory tanks. - *N. Z. Jl mar. Freshwat. Res.* 37: 45-52.
- Roy, A. H., Rosemond, A. D., Leigh, D. S., Paul, M. J. and Wallace, J. B. 2003a. Habitat-specific responses of stream insects to land cover disturbance: biological consequences and monitoring implications. - *J. N. Am. benthol. Soc.* 22: 292-307.
- Roy, A. H., Rosemond, A. D., Paul, M. J., Leigh, D. S. and Wallace, J. B. 2003a. Stream macroinvertebrate response to catchment urbanisation (Georgia, U.S.A.). - *Freshwat. Biol.* 48: 329-346.
- Roy, B. and Dutta, B. K. 2003a. In vitro lethal efficacy of leaf extract of *Cannabis sativa* Linn on the larvae of *Chironomus samoensis* Edward: an insect of public health concern. - *Indian J. exp. Biol.* 41: 1338-1341.

- Ruetz, C. R. III, Hurford, A. L. and Vondracek, B. 2003a. Interspecific interactions between brown trout and slimy sculpin in stream enclosures. - *Trans. Am. Fish. Soc.* 132: 611-618.
- Ruiz, Z., Rippey, B. H. R. T., Carter, C. E. and Bennion, H. 2003a. Chironomid-dissolved oxygen inference models from Northern Irish lakes: comparisons with Canadian counterparts. - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp., Espoo*: 205.
- Sadaka, N. and Ponge, J.-F. 2003a. Soil animal communities in holm oak forests: influence of horizon, altitude and year. - *Eur. J. Soil Biol.* 39: 197-207.
- Sæther, O. A. 2003a. A review of *Orthocladius* subgen. *Symposiocladius* Cranston (Diptera: Chironomidae). - *Aquat. Insects* 25: 281-317.
- Sæther, O. A. and Andersen, T. 2003a. Redescription of *Rhinocladius* Edwards (Diptera: Chironomidae: Orthoclaadiinae). - *Zootaxa* 217: 1-20.
- Sæther, O. A. and Ekrem, T. 2003a. Biogeography of afrotropical Chironomidae (Diptera), with special reference to Gondwanaland. - *Cimbebasia* 19: 123-139.
- Sæther, O. A. and Ferrington, L. C. Jr.. 2003a. Nomenclature notes on some orthoclaids (Diptera: Chironomidae). - *Zootaxa* 322: 1-7.
- Samietz, R. 2003a. Beitrag zur Chironomidenfauna (Diptera: Chironomidae) der Hochmoore des Thüringer Waldes. - *Thüring. Faun. Abh.* 9: 227-232.
- Sanford, M. R., Keiper, J. B. and Walton, W. E. 2003a. The impact of wetland vegetation drying time on abundance of mosquitoes and other invertebrates. - *J. Am. Mosquito Control Ass.* 19: 361-366.
- Sanseverino, A. M., Wiedenbrug, S. and Fittkau, E. J. (2002) 2003a. *Marauia* group: a new species group in the genus *Tanytarsus* VAN DER WULP, 1874, from the Neotropics (Diptera, Chironomidae). - *Studia dipterol.* 9: 453-468.
- Sanzone, D. M., Meyer, J. L., Marti, E., Gardiner, E. P., Tank, J. L. and Grimm, N. B. 2003a. Carbon and nitrogen transfer from a desert stream to riparian predators. - *Oecologia* 134: 238-250.
- Scheibe, M. A. 2003a. Über den Einfluss von Straßenbeleuchtung auf aquatische Insekten (Ephemeroptera, Plecoptera, Trichoptera, Diptera: Simuliidae, Chironomidae, Empididae). - *Nat. Landsch.* 78: 264-267.
- Schleuter, M. und Haybach, A. 2003a. Das Makrozoobenthos des Mains in den Jahren 1992-2001 - Eine Artenliste. - *Lauterbornia* 48: 45-56.
- Schuler, L. J., Wheeler, M., Bailer, A. J. and Lydy, M. J. 2003a. Toxicokinetics of sediment-sorbed benzo[a]pyrene and hexachlorobiphenyl using the freshwater invertebrates *Hyaella azteca*, *Chironomus tentans*, and *Lumbriculus variegatus*. - *Envir. Toxic. Chem.* 22: 439-449.
- Schulz, R, Moore, M. T., Bennett, E. R., Farris, J. L., Smith, S. and Cooper, C. M. 2003a. Methyl parathion toxicity in vegetated and nonvegetated wetland mesocosms. - *Envir. Toxic. Chem.* 22: 1262-1268.
- Scrimgeour, G. J. and Kendall, S. 2003a. Effects of livestock grazing on benthic invertebrates from a native grassland ecosystem. - *Freshwat. Biol.* 48: 347-362.
- Segner, H., Caroll, K., Fenske, M., Janssen, C. R., Maack, G., Pascoe, D., Schäfers, C., Vandenberg, G. F., Watts, M. and Wenzel, A. 2003a. Identification of endocrine-disrupting effects in aquatic vertebrates and invertebrates: report from the European IDEA project. - *Ecotoxic. envir. Saf.* 54: 302-314.
- Seredszus, F. 2003a. *Wasserinsekten des Baltischen Bernsteins unter besonderer Berücksichtigung der Chironomiden. Grundlagen zum Verständnis von aquatischen Lebensräumen und Lebensgemeinschaften im eozänen Bernsteinwald.* - Diss., Univ. Köln. 177 pp.
- Seredszus, F. und Wichard, W. (2002) 2003a. Buchonomyiinae (Diptera, Chironomidae) im Baltischen Bernstein. - *Studia dipterol.* 9: 393-402.
- Sergeeva, I. V. 2003a. Khironomidy podsem. Tanypodinae (Diptera, Chironomidae) Kuril'skikh ostrovov. (Midges of the subfamily Tanypodinae (Diptera, Chironomidae) of Kuril Islands.) - *Ent. Obozr.* 82: 846-849; 946.
- Sharma, O. P, Tripathi, N. K. and Khanna, P. 2003a. Karyological study on *Cryptochironomus* sp. (Diptera:

- Chironomidae) from Jammu. - *Nucleus, Calcutta. Int. J. Cytol.* 46: 48-53.
- Sherfy, M. H. and Kirkpatrick, R. L. 2003a. Invertebrate response to snow goose herbivory on moist-soil vegetation. - *Wetlands* 23: 236-249.
- Shieh, S.-H., Ward, J. V. and Kondratieff, B. C. 2003a. Longitudinal changes in macroinvertebrate production in a stream affected by urban and agricultural activities. - *Arch. Hydrobiol.* 157: 483-503.
- Shitikov, V. K., Rozenberg, G. S. i Zinchenko, T. D. 2003a. *Kolichestvennaya gidroekologiya: metody sistemnoi identifikatsii. (Quantitative hydroecology: methods of system identification.)*. - Inst. Ekol. Volzh. Bass. Ross. Akad. Nauk, Tol'yatti. 463 pp.
- Shobanov, N. A. 2003a. Morfologicheskaya differentsiatsiya vidov roda *Chironomus* Mg. gruppy *plumosus* (Diptera, Chironomidae). Kukolki. (Morphological differentiation of species of the *plumosus* group, genus *Chironomus* Mg. (Diptera, Chironomidae). Pupae.) - *Ent. Obozr.* 82: 472-486; 531.
- Sierszen, M. E., McDonald, M. E. and Jensen, D. A. 2003a. Benthos as the basis for arctic lake food webs. - *Aquat. Ecol.* 37: 437-445.
- Siirin, M. T. 2003a. *Chironomus arcustylus* sp. n. iz gruppy *obtusidens* (Diptera, Chironomidae). (*Chironomus arcustylus* sp. n. of the group *obtusidens* (Diptera, Chironomidae).) - *Zool. Zh.* 82: 613-622.
- Siirin, M. T., Rubtsov, N. B., Karamysheva, T. V., Katokhin, A. V., Karagodin, D. A. i Kiknadze, I. I. 2003a. (Molecular-cytogenetic characteristics of B-chromosomes in chironomid (Diptera, Chironomidae).) - *Tsitologiya* 45: 582-589.
- Sindilariu, P.-D. and Freyhof, J. 2003a. Food overlap of benthic fishes in the Danube Delta, with special respect to two invasive gobiids (Teleostei: Gobiidae, Percidae, Cyprinidae). - *Lauterbornia* 46: 149-157.
- Smith, H., Wood, P. J. and Gunn, J. 2003a. The influence of habitat structure and flow permanence on invertebrate communities in karst spring systems. - *Hydrobiologia* 510: 53-66.
- Smolders, A. J. P., Lock, R. A. C., Van der Velde, G., Medina Hoyos, R. I. and Roelofs, J. G. M. 2003a. Effects of mining activities on heavy metal concentrations in water, sediment, and macroinvertebrates in different reaches of the Pilcomayo River, South America. - *Archs envir. Contam. Toxic.* 44: 314-323.
- Solimini, A. G., Ruggiero, A., Bernardini, V. and Carchini, G. 2003a. Temporal pattern of macroinvertebrate diversity and production in a new man made shallow lake. - *Hydrobiologia* 506-509: 373-379.
- Soop, T., Nashchekin, D., Zhao, J., Sun, X., Alzhanova-Ericsson, A. T., Björkroth, B., Ovchinnikov, L. and Daneholt, B. 2003a. A p50-like Y-box protein with a putative translational role becomes associated with pre-mRNA concomitant with transcription. - *J. Cell Sci.* 116: 1493-1503.
- Speziár, A. and Bíró, P. 2003a. Population structure and feeding characteristics of Volga pikeperch, *Sander volgensis* (Pisces, Percidae), in Lake Balaton. - *Hydrobiologia* 506-509: 503-510.
- Stead, T. K., Schmid-Araya, J. M. and Hildrew, A. G. 2003a. All creatures great and small: patterns in the stream benthos across a wide range of metazoan body size. - *Freshwat. Biol.* 48: 532-547.
- Stefanova, I., Ognjanova-Rumenova, N., Hofmann, W. and Ammann, B. 2003a. Late Glacial and Holocene environmental history of the Pirin Mountains (SW Bulgaria): a paleolimnological study of Lake Dalgoto (2310 m). - *J. Paleolimnol.* 30: 95-111.
- Stenroth, P. and Nyström, P. 2003a. Exotic crayfish in a brown water stream: effects on juvenile trout, invertebrates and algae. - *Freshwat. Biol.* 48: 466-475.
- Stevens, M. M., Warren, G. N. and Braysher, B. D. 2003a. Oviposition response of *Chironomus tepperi* to nitrogenous compounds and bioextracts in two-choice laboratory tests. - *J. chem. Ecol.* 29: 911-920.
- Stewart, T. W., Shumaker, T. L. and Radzio, T. A. 2003a. Linear and nonlinear effects of habitat structure on composition and abundance in the macroinvertebrate community of a large river. - *Am. Midl. Nat.* 149: 293-305.
- Stief, P., Altmann, D., Franke, U. and De Beer, D. 2003a. Probing the microenvironment of the sediment-dwelling insect *Chironomus riparius*. - *Abstr. Aquat. Sci. Meet. 2003, Salt Lake City*, p. 124. Am. Soc. Limnol. Oceanogr.

- Stocker, A. J. and Gorab, E. 2003a. Local enrichment with homopolymeric (dA/dT) DNA in genomes of some lower dipterans and *Drosophila melanogaster*. - *Chromosoma* 111: 455-460.
- Stoffels, R. J., Karbe, S. and Paterson, R. A. 2003a. Length-mass models for some common New Zealand littoral-benthic macroinvertebrates, with a note on within-taxon variability in parameter values among published models. - *N. Z. Jl mar. Freshwat. Res.* 37: 449-460.
- Stoichev, S. and Danova, E. 2003a. Hydrofaunistic investigation of the Mussalenski Ezera Glacial Lakes, Eastern Rila Mountains, Southwest Bulgaria. - *Acta zool. bulg.* 55: 75-80.
- Strayer, D. L., Lutz, C., Malcom, H. M., Munger, K. and Shaw, W. H. 2003a. Invertebrate communities associated with a native (*Vallisneria americana*) and an alien (*Trapa natans*) macrophyte in a large river. - *Freshwat. Biol.* 48: 1938-1949.
- Su, T. Y., Webb, J. R., Meyer, R. R. and Mulla, M. S. 2003a. Spatial and temporal distribution of mosquitoes in underground storm drain systems in Orange County, California. - *J. Vector Ecol.* 28: 79-89.
- Sushchik, N. N., Gladyshev, M. I., Moskvichova, A. V., Makhutova, O. N. and Kalachova, G. S. 2003a. Comparison of fatty acid composition in major lipid classes of the dominant benthic invertebrates of the Yenisei River. - *Comp. Biochem. Physiol. Pt B Biochem. molec. Biol.* 134B: 111-122.
- Suren, A. M., Biggs, B. J. F., Duncan, M. J., Bergey, L. and Lambert, P. 2003a. Benthic community dynamics during summer low-flows in two rivers of contrasting enrichment. 2. Invertebrates. - *N. Z. Jl mar. Freshwat. Res.* 37: 71-83.
- Sushchik, N. N., Gladysheva, M. I., Moskvichova, A. V., Makhutova, O. N. and Kalachova, G. S. 2003a. Comparison of fatty acid composition in major lipid classes of the dominant benthic invertebrates of the Yenisei River. - *Comp. Biochem. Physiol. B: Biochem. molec. Biol.* 134: 111-122.
- Tanaka, N., Hashizume, S., Onogawa, M. and Sasa, M. 2003a. Studies on chironomid midges in winter from the Tone River, Gunma Prefecture. - *Med. Ent. Zool.* 54: 299-303.
- Tanaka, N., Sasa, M. and Hashizume, S. 2003a. Studies on chironomid midges in a rice paddy area in the suburbs of Maebashi City, Gunma Prefecture. - *Med. Ent. Zool.* 54: 121-124.
- Tank, S. E., Schindler, D. W. and Arts, M. T. 2003a. Direct and indirect effects of UV radiation on benthic communities: epilithic food quality and invertebrate growth in four montane lakes. - *Oikos* 103: 651-667.
- Tancioni, L., Mariani, S., Maccaroni, A., Mariani, A., Massa, F., Scardi, M. and Cataudella, S. 2003a. Locality-specific variation in the feeding of *Sparus aurata* L.: evidence from two Mediterranean lagoon systems. - *Estuar. coast. Shelf Sci.* 57:469-474.
- Thapanya, D., Prapamontol, T., Jatisatienr, A. and Prakobvitayakit, O. 2003a. Impact of pesticides on chironomid cholinesterase activity and the macroinvertebrate community. - *Verh. int. Verein. Limnol.* 28: 1724-1726.
- Tillman, D. C., Moerke, A. H., Ziehl, C. L. and Lamberti, G. A. 2003a. Subsurface hydrology and degree of burial affect mass loss and invertebrate colonisation of leaves in a woodland stream. - *Freshwat. Biol.* 48: 98-107.
- Tockner, K., Robinson, C. T. and Burgherr, P. 2003a. Drift and colonization dynamics. - In: Ward, J. V. and Uehlinger, U. (eds.): *Ecology of a glacial flood plain*, pp. 243-257. Kluwer Acad. Publs, Dordrecht, Boston, Lond.
- Toft, J. D., Simenstad, C. A., Cordell, J. R. and Grimaldo, L. F. 2003a. The effects of introduced water hyacinth on habitat structure, invertebrate assemblages, and fish diet. - *Estuaries* 26: 746-758.
- Tolonen, K. T., Hämäläinen, H., Holopainen, I. J., Mikkonen, K. and Karjalainen, J. 2003a. Body size and substrate association of littoral insects in relation to vegetation structure. - *Hydrobiologia* 499: 179-190.
- Townsend, C. R., Dolédec, S., Norris, R., Peacock, K. and Arbuckle, C. 2003a. The influence of scale and geography on relationships between stream community composition and landscape variables: description and prediction. - *Freshwat. Biol.* 48: 768-785.
- Trivinho-Strixino, S. and Sanseverino, A. M. 2003a. *Tanytarsus rhabdomantis*: New combination for *Nimbocera rhabdomantis* Trivinho-Strixino & Strixino, 1991 (Diptera: Chironomidae). - *Zootaxa* 389: 1-10.

- Trivinho-Strixino, S. and Strixino, G. 2003a. The immature stages of two *Caladomyia* Sæwedal, 1981 species from São Paulo State, Brazil (Chironomidae, Chironominae, Tanytarsini). - *Revta bras. Ent.* 47: 597-602.
- Trzcinski, M. K., Walde, S. J. and Taylor, P. D. 2003a. Colonisation of pitcher plant leaves at several spatial scales. - *Ecol. Ent.* 28: 482-489.
- Turak, E. and Koop, K. 2003a. Use of rare macroinvertebrate taxa and multiple-year data to detect low-level impacts in rivers. - *Aquat. Ecosyst. Hlth Mgmt* 6: 167-175.
- Varga, I. 2003a. Structure and changes of macroinvertebrate community colonizing decomposing rhizome litter of common reed at Lake Fertő/Neusiedler See (Hungary). - *Hydrobiologia* 506-509: 413-420.
- Velásquez, S. M. and Miserendino, M. L. 2003a. Habitat type and macroinvertebrate assemblages in low order Patagonian streams. - *Arch. Hydrobiol.* 158: 461-483.
- Velle, G. and Brooks, S. J. 2003a. An evaluation of chironomids as a tool for inferring Holocene climate. - In: Sorvari, S., Salonen, V.-P., Korhola, A. and Ojala, A. (eds.): *Abstr. 9<sup>th</sup> Int. Paleolimnol. Symp., Espoo*: 105.
- Verbeek, W. C. E. P., van Duinen, G.-J., Moller Pillot, H. K. M. and Esselink, H. 2003a. *Lasiodiamesa gracilis* (Chironomidae: Podonominae) new for the Dutch fauna. - *Ent. Ber., Amst.* 63: 40-42.
- Vīksne, J. 2003a. *Ramsāres konvencijas vietas - Engures ezera - monitorings 2003. gadā. (Ramsar convention site Lake Engure monitoring in 2003.)* - Latv. Univ., Biol. Inst., Salaspils. 91 pp.
- Villalobos, L., Woelfl, S., Parra, O. and Campos, H. 2003a. Lake Chapo: a baseline study of a deep, oligotrophic North Patagonian lake prior to its use for hydroelectricity generation: II. Biological properties. - *Hydrobiologia* 510: 225-237.
- Vinogradova, E. B. and Petrova, N. A. 2003a. Synanthropization in chironomids: *Chironomus riparius* (Diptera, Chironomidae) as an example. - *Proc. Zool. Inst. Russ. Acad. Sci.* 299: 187-196.
- Walker, I. 2003a. The benthic insect fauna of the blackwater forest stream Rio Tarumã-Mirím (Manaus, Amazonas): patterns of population dynamics and their implications for ecosystem stability. - *Amazoniana* 17: 471-480.
- Walker, I. R., Levesque, A. J., Pienitz, R. and Smol, J. P. 2003a. Freshwater midges of the Yukon and adjacent Northwest Territories: a new tool for reconstructing Beringian paleoenvironments? - *J. N. Am. benthol. Soc.* 22: 323-337.
- Wang, B. and Yang, L. 2003a. (Bioassessment of Qinhuai River using a river biological index.) - *Shengtai Xuebao [Acta ecol. sin.]* 23: 2082-2091.
- Wantzen, K. M. 2003a. Cerrado streams - characteristics of a threatened freshwater ecosystem type on the Tertiary Shields of Central South America. - *Amazoniana* 17: 481-502.
- Wappler, T. 2003a. Die Insekten aus dem Mittel-Eozän des Eckfelder Maeres, Vulkaneifel. - *Mainz. naturw. Arch. Bh.* 27: 234 + 36 pp.
- Ward, J. V. and Uehlinger, U. (eds.) 2003a. Appendix V. List of benthic and hyporheic taxa. - *Ecology of a glacial flood plain*, pp. 291-297. Kluwer Acad. Publs, Dordrecht, Boston, Lond.
- Watanabe, M., Kikawada, T. and Okuda, T. 2003a. Increase of internal ion concentration triggers trehalose synthesis associated with cryptobiosis in larvae of *Polypedilum vanderplanki*. - *J. exp. Biol.* 206: 2281-2286.
- Watts, M. M., Pascoe, D and Carroll, K. 2003a. Exposure to 17 $\alpha$ -ethinylestradiol and bisphenol A - effects on larval moulting and mouthpart structure of *Chironomus riparius*. - *Ecotoxic. envir. Saf.* 54: 207-215.
- Weigel, B. M. 2003a. Development of stream macroinvertebrate models that predict watershed and local stressors in Wisconsin. - *J. N. Am. benthol. Soc.* 22: 123-142.
- Weigel, B. M., Wang, L., Rasmussen, P. W., Butcher, J. T., Stewart, P. M., Simon, T. P. and Wiley, M. J. 2003a. Relative influence of variables at multiple spatial scales on stream macroinvertebrates in the Northern Lakes and Forest ecoregion, U.S.A. - *Freshwat. Biol.* 48: 1440-1461.
- Weigelhofer, G. and Waringer, J. 2003a. Vertical distribution of benthic macroinvertebrates in riffles versus deep runs with differing contents of fine sediments (Weidlingbach, Austria). - *Int. Rev. Hydrobiol.* 88: 304-313.

- Weigelhofer, G. and Waringer, J. 2003b. Response of macroinvertebrates to fine sediment accumulations within the hyporheic zone of a calcareous sandstone stream (Weidlingbach, Austria). - *Arch. Hydrobiol. Suppl. 147, Large Rivers 14*: 327-346.
- Werner, D. and Pont, A. C. 2003a. Dipteran predators of simuliid blackflies: a worldwide review. - *Med. vet. Ent. 17*: 115-132.
- White, J. and Irvine, K. 2003a. The use of littoral mesohabitats and their macroinvertebrate assemblages in the ecological assessment of lakes. - *Aquat. Conserv. mar. Freshwat. Ecosyst. 13*: 331-351.
- Williams, D. D. 2003a. The brackishwater hyporheic zone: invertebrate community structure across a novel ecotone. - *Hydrobiologia 510*: 153-173.
- Williams, D. D. and Fulthorpe, R. R. 2003a. Using invertebrate and microbial communities to assess the condition of the hyporheic zone of a river subject to 80 years of contamination by chlorobenzenes. - *Can. J. Zool. 81*: 789-802.
- Williams, L. R., Taylor, C. M. and Warren, M. 2003a. Influence of fish predation on assemblage structure of macroinvertebrates in an intermittent stream. - *Trans. Am. Fish. Soc. 132*: 120-130.
- Williams, L. R., Taylor, C. M., Warren, M. L. Jr. and Clingenpeel, J. A. 2003a. Environmental variability, historical contingency, and the structure of regional fish and macroinvertebrate faunas in Ouachita Mountain stream systems. - *Envir. Biol. Fishes 67*: 203-216.
- Wolnicki, J., Myszkowski, L. and Kamiński, R. 2003a. Effect of supplementation of a dry feed with natural food on growth, condition and size distribution of juvenile tench *Tinca tinca* (L.). - *J. appl. Ichthyol. 19*: 157-160.
- Wong, D. C. L., Whittle, D., Maltby, L. and Warren, P. 2003a. Multivariate analyses of invertebrate community responses to a C<sub>12</sub>-<sub>15</sub>AE-3S anionic surfactant in stream mesocosms. - *Aquat. Toxic. 62*: 105-117.
- Wright, J. F., Clarke, R. T., Gunn, R. J. M., Winder, J. M., Kneebone, N. T. and Davy-Bowker, J. 2003a. Response of the flora and macroinvertebrate fauna of a chalk stream site to changes in management. - *Freshwat. Biol. 48*: 894-911.
- Wymer, D. A. and Cook, S. B. 2003a. Effects of Chironomidae (Diptera) taxonomic resolution on multivariate analyses of aquatic insect communities. - *J. Freshwat. Ecol. 18*: 179-186.
- Yamamoto, Y., Koshikawa, K., Terui, N., Mita, H., Matsuoka, A. and Shikama, K. 2003a. <sup>1</sup>H-NMR study of dynamics and thermodynamics of Cl<sup>-</sup> binding to ferric hemoglobin of a midge larva (*Tokunagayusurika akamusi*). - *Biochim. biophys. Acta 1652*:136-143.
- Yan, Y. and Liang, Y. 2003a. (Energy flow of macrozoobenthos community in an algal lake, Houhu Lake (Wuhan, China).) - *Chin. J. Oceanol. Limnol. 21*: 229-244.
- Yan, Y. and Liang, Y. 2003b. (Energy flow of macrozoobenthic community in a macrophytic lake, Biandantang Lake.) - *Shengtai Xuebao [Acta ecol. sin.] 23*: 527-538.
- Yin, D., Hu, S., Jin, H. and Yu, L. 2003a. Deriving freshwater quality criteria for 2,4,6-trichlorophenol for protection of aquatic life in China. - *Chemosphere 52*: 67-73.