

# ***Cryptochironomus*. An identification key to the larvae and pupal exuviae in Europe**

Henk J. Vallenduuk and E. Morozova

With 35 figures and 5 tables

**Keywords:** *Cryptochironomus*, Diptera, Insecta, The Netherlands, Volga, Russia, Europe morphology, identification, distribution, larva, pupa, exuvia

**Schlagwörter:** *Cryptochironomus*, Diptera, Insecta, Niederlande, Wolga, Russland, Europa, Morphologie, Bestimmung, Verbreitung, Larve, Puppe, Exuvie

The genus *Cryptochironomus* Kieffer is represented in Europe by about ten species. An illustrated key to the pupal exuviae and larvae is given to the known and suggested species. The autecology unfortunately is little known.

## **1 Introduction**

For the assessment of species specific characters it is necessary to have larvae for which the specific identification is certain. Only by rearing larvae to adults can this be achieved. Some years ago we began collecting larvae and rearing them to adult. When successful, this provides the adult and the pupal and larval exuviae. The pupal exuviae enable accurate identifications using the keys of Langton (1991) and Langton & Visser (2003). (The pupal exuviae of *Cryptochironomus defectus* (Kieffer) was not included in the earlier key, but as a result of our rearing program, has been included in the 2003 one.) The adults of *Cryptochironomus* are very difficult to separate. Morozova has also investigated the larvae cytologically. When all instars of all species have been obtained, it will be possible to make a full revision of the genus; Morozova has embarked on this. In the meantime, the interpretation of the species is that of Langton & Visser (2003).

## **2 Method used for rearing larvae**

For rearing larvae, Vallenduuk used small polystyrene culture boxes with a ventilation cap. If placed the right way up as shown in the photograph, the measurements are: bottom 47x27mm, height 65mm. In the cap there is a hole made for putting in a Pasteur pipette, which is cut off. If the adult flies to the light, it will be captured in the top of the pipette. Now it is easy to collect the adult by pinching the "neck" and pouring some alcohol into the pipette.

If the adult does not fly into the pipette, some tissue soaked in ethyl acetate is put in the pipette. In this case the cap has to be closed tightly.

In a single culture box only one larva is placed. On the bottom was put some material from the collection site, and the box filled with water also from the site. Prepupae are best, for they do not eat and the adult emerges in a few days. The larvae were not fed.

Every day the cultures were checked. *Cryptochironomus* adults usually fly to the tops of the pipettes. The larvae build cases from small grains of sand. Pupation occurs in the bottom of the box. The pupal exuviae always float on the water surface. Usually it was possible to find the larval exuviae in the water or still in the pupal case.



### 3 Taxonomy

The family Chironomidae is usually divided into eight subfamilies. The genus *Cryptochironomus* belongs to the subfamily Chironominae and to the tribe Chironomini.

The first descriptions of species that are now in the genus *Cryptochironomus* are from Kieffer, Walker, Harnisch and Pagast. The descriptions are very incomplete and are virtually unusable: the few characters used differ from description to description. Despite some later authors (e.g. Shilova 1966, Reiss 1968) providing additional characters, it is important to produce new descriptions. The adults obtained from the rearing program will be used for comparison with type species.

Based on the interpretation of Langton in Langton (1991) and Langton & Visser (2003) we think the following species exist (Tab. 1). We primarily used in the list the taxonomy of Ashe & Cranston (1990).

The larvae as well as the pupal exuviae of the *Cryptochironomus ussouriensis* differ greatly from all other species of the genus. It appears justifiable to place this species in another genus or a separate subgenus. For the time being it is retained in the genus *Cryptochironomus*.

**Tab. 1: The species of the genus *Cryptochironomus* Kieffer, 1918. \* = exuviae, obtained by rearing larva(e)**

Species	Ashe & Cranston (1990)	Langton (1991) Pe	Vallenduuk rearing larva	Literature
albofasciatus (Staeger, 1839)	+	+ *		
crassiforceps Goetghebuer, 1931	+			Shilova (1966) Pankratova (1983)
defectus (Kieffer, 1913)	+		+	
denticulatus (Goetghebuer, 1921)	+	+ *		
obreptans (Walker, 1856)	+	+ *	+	
psittacinus (Meigen, 1830)	+	+ *	+	
redekei (Kruseman, 1933)	+	+	+	
rostratus Kieffer, 1921	+	+ *	+	
supplicans (Meigen, 1830)	+	+ *	+	
ussouriensis (Goetghebuer, 1933)	+			Shilova (1959), Pankratova (1983), Kiknadze et al. (1991)

#### 4 European distribution of *Cryptochironomus* (Tab. 2)

There is little known about the distribution of the species in The Netherlands and other European countries, because it has not been possible to identify the larvae and information is only slowly accruing using pupal exuviae. For the purpose of this paper we can use only the data from pupal exuviae. Because the species *defectus* and *supplicans* both run to *supplicans* in Langton's 1991 key; prior to the publication of Langton & Visser's 2003 key these two species were recorded under the one name, *supplicans*.

**Tab. 2: Distribution of *Cryptochironomus***

Species	Ashe & Cranston (1990)	Klink & Moller-Pillot Netherlands	Number of localities Netherlands
albofasciatus	Europe	+	?1
crassiforceps	Caspian Sea		
defectus	Europe	+	20
denticulatus	Europe		
obreptans	Europe	+	71
psittacinus	Europe, Nearctic	+	23
redekei	Europe, Asia	+	14
rostratus	Europe, Asia	+	13
supplicans	Europe, Asia		16
supplicans/defectus*		+	77
ussouriensis	Eastern-Europe		

\*Identifications of *supplicans* pupal exuviae and pupae can belong to both species.

## 5 Ecology of European *Cryptochironomus*

The genus *Cryptochironomus* is widespread over a great part of the world. The larvae occur in various water types. All species dwell in the upper part of the water body, seldom being found on hard substrata near the bottom. They prefer sandy substrata with some detritus and fine organic material. They do not occur between dense vegetation. When the depth of detritus increases, the larvae move out. When the depth becomes too great, eggs are no longer laid there. The presence of their food, Oligochaeta and Chironomidae, seems to be the most important factor. Lenz thinks they are partly predators (based on Luferov 1958), because the presence of detritus in the gut. It appears that all species require a good oxygen level and clear water. Regularly more than one species is found at the same site. These are vague generalisations: clearly, further investigation is needed to determine the ecology of each species.

The larvae occur in many different water types. They seldom occur in waters with the pH lower than 5,0. They probably need an oxygenpercentage of at least 40 %. For The Netherlands, in the rivers Rhine, Lek, IJssel and Waal, the common species is *C. rostratus*, though *C. defectus* does occur. In the faster flowing river Meuse *Cryptochironomus* species have not yet been found. In the delta region, close to the North Sea, and in sites with sand bottoms, lakes, watercourses and canals, mostly *C. defectus*, *C. obreptans* and *C. supplicans* occur. In pools, smaller lakes, ponds and "wiels" the species *C. psittacinus* and *C. redekei* may occur. We have found that they do not occur in waters with dense vegetation and seldom in waters with a peat bottom. They seldom occur in narrow ditches (less than 3 meters) nor in temporary waters. *Cryptochironomus* species are rare in brackish or humic waters; they are found in oligo-/mesotroph as well as in eutrophic or oligo- and mesosaprobic waters.

## 6 Key to the larvae of the genus *Cryptochironomus*

In contrast with many chironomids the larvae of *Cryptochironomus* are easy to recognise when collecting: the body blood red in colour, with a comparatively small, posteriorly swollen, orange-coloured head. The larvae often move by strong wiggling. They lose their red colour in alcohol; the head is usually held backwards; the body stretched and rather stiff.

### Methods and terminology

The length and the width of the head can best be measured by placing the larvae dorsoventrally on a slide. In this position the width can be measured (ensure that the head is fully submerged in the alcohol). For making the head transparent lactic acid is recommended: a solution of one part lactic acid with three parts of 70 % alcohol works well. For making detailed measurements, the larva is best placed on a slide with coverslips placed alongside to the height of the head before a coverslip is placed on top. More alcohol may be added with a pipette to keep the animal submerged.

The distance between the two S3 setae is best taken with the head dorsal side uppermost.

### Terms, which are used in the key

The terminology is based on Sæther (1980) and Cranston & Reiss (1983). Some new terms are here proposed.

A1 = first antennal segment

Antenna, first segment length = from base to end of sclerotized ring

IC index = head width divided by its length

LMP = total length from tip of midmental tooth to the ventral edge of the postoccipital margin (Fig. 1)

Mandible length must be made with the mandible dissected from the head.

S3-S3 = distance between the two S3 setae (Fig. 2)

S3/A1 index = S3-S3 divided by the length of the first antennal segment (Fig. 2)

VmP length = length of the ventromental plate from the anterior to the posterior border (Fig. 4)







Fig. 1: *Cryptochironomus obreptans*. Head ventral. a = mentum, b = gula, c = postocipital margin, d = antenna (from Kiknadze et al. (1991))

Fig. 2: *Cryptochironomus spec.* Head dorsal, frontal apotome and labral sclerites. a = frontal apotome, b = labral sclerite 1, c = antenna, first segment (A1), d = measuring A1 length (Morozova)

Fig. 3: *Cryptochironomus psittacinus*. Mandible with SSd (a) and mandible length (Vallenduuk)

Fig. 4: *Cryptochironomus obreptans*. a = ventromental plate (VmP) length (after Kiknadze et al. 1991)

### Key to the instar III or IV (Tab. 3)

The key is based on characters observed in instar IV larvae. Only larvae in this instar are identifiable with certainty; prepupal larvae, with obviously swollen thorax, are reliably instar IV. If instar III is not recognised, the identification will be wrong in many cases.

- 1 Gula with obvious pigmentation 2
- Gula with at most indistinct pigmentation 3
  
- 2 Body length 6,5 mm or more. Head width (380)410-504  $\mu\text{m}$   
*C. obreptans* and *C. supplicans*, instar IV
- Body length less than 6,5 mm. Head width 280-340  $\mu\text{m}$   
*C. obreptans* and *C. supplicans*, instar III
  
- 3 Body length 7,5 mm or more instar IV
- Body length less than 7,5 mm 4
  
- 4 Head length 520  $\mu\text{m}$  or more instar IV
- Head length less than 520  $\mu\text{m}$  *C. rostratus*, *C. denticulatus* instar IV  
sometimes *C. defectus*, instar IV  
all other species instar III (see key to the instar IV)

**Tab. 3: *Cryptochironomus*. Pigmentation of gula and measurements of larvae with a body length less than 7,5 mm and a head length less than 520  $\mu\text{m}$**

species	instar	pigmentation gula	Total length (mm)	Head length ( $\mu\text{m}$ )	Head width ( $\mu\text{m}$ )
psittacinus	III	-	6,0	500	400
defectus	IV	-	7,0	496	355
redekei	III	-	6,5	480	320
redekei	III	-	6,5	480	320
rostratus	IV	-	?	460	320
rostratus	IV	-	?	450	320
redekei	III	-	7,1	450	310
redekei	III	-	7,1	450	310
psittacinus	III	-	5,5	440	360
obreptans	?IV	+	6,2	418	308
obreptans	?III	+	6,5	352	286
obreptans	III	+	5,8	340	260
obreptans	III	+	6,1	330	?
defectus	III	-	5,0	320	240
obreptans	III	+	6,0	310	?
obreptans	III	+	4,7	280	240
defectus	III	-	4,1	240	320

## Key to the instar IV-larvae

With this key it is not possible to identify *C. albofasciatus* or *C. crassiforceps*: no larvae were available and no data were found in the literature. The measurements of larvae from The Netherlands often differ from those from Russia. In the key these are amalgamated, in table 5 they are kept separate.

Ensure that the larva is instar IV – see the preceding key to instar III and IV.

- 1 Midmental tooth not pigmented (Fig. 5). Last lateral tooth of mentum ending much higher than first lateral tooth. Midmental tooth without incisions 2
  - Midmental tooth completely pigmented (Fig. 7). Lateral teeth of mentum horizontally placed. Midmental tooth with incisions (known only from Russia) *ussouriensis*
- 2 Gula with pigmentation (Fig. 9). The pigmentation often reaches to about half the length of the gula, but can also be a small spot in front of the postoccipital margin (Fig. 1c). Postoccipital margin ventrally as pigmented as the teeth of the mandible. Head length 530-680  $\mu\text{m}$  3
  - Gula without any sign of pigmentation (Fig. 10). Postoccipital margin in most of the cases completely pale or dark-yellow; if with dark ventral pigmentation than the head length nearly always 800  $\mu\text{m}$  or more. (Once larvae of *C. psittacinus* were collected with a head length of 600 and 620  $\mu\text{m}$ . In such cases use the table with measurements -see Tab. 5, p. 17) 4
- 3 *obreptans* and *supplicans*

Both species are very difficult to separate. We found the following differences: however the values overlap greatly

  - a S3-S3 (Fig. 2) 45-75  $\mu\text{m}$ . S3/A1-index (Fig. 2); mean: 0,66 (see table below). Moderately long claws of anterior parapods with relatively blunt teeth (Fig. 8) *obreptans*
  - b S3-S3 (Fig. 2) 60-86  $\mu\text{m}$ . S3/A1-index (Fig. 2); mean: 0,92 (see table below). Moderately long claws of anterior parapods with relatively sharp teeth (Fig. 6) *supplicans*

**Tab. 4: Measurements of *C. obreptans* and *C. supplicans***

		Antenna 1 ( $\mu\text{m}$ )	S3-S3 ( $\mu\text{m}$ )	S3/A1 Index	n	LMP ( $\mu\text{m}$ )	n
obreptans	range	78-105	45-75	0,53-0,79	25	210-260	15
	mean	94	62	0,66		235	
	stdev	9	7	0,10		9	
supplicans	range	70-90	60-80	0,76-1,04	28	200-238	13
	mean	80	74	0,92		223	
	stdev	8	8	0,08		9	




Fig. 5: *Cryptochironomus obreptans*. Mentum (from Kiknadze et al. 1991)  
 Fig. 6: *C. supplicans*. Moderately long claws of anterior parapods (Morozova)  
 Fig. 7: *C. ussouriensis*. Mentum (from Kiknadze et al. 1991)  
 Fig. 8: *C. obreptans*. Moderately long claws of anterior parapods (Morozova)  
 Fig. 9: *C. obreptans*. Head ventral with pigmentation (from Kiknadze et al. 1991)  
 Fig. 10: *C. defectus*. Head ventral without pigmentation (Morozova)

4	Head length 395-635 $\mu\text{m}$ . Head width 265-435 $\mu\text{m}$	5
-	Head length (600) 800-925 $\mu\text{m}$ . Head width 530-720 $\mu\text{m}$	7
5	LMP (Fig. 1) 180-240 $\mu\text{m}$ . VmP length (Fig. 4) 45-63 $\mu\text{m}$ (larvae from The Netherlands: 53-63 ). Head length 475-635 $\mu\text{m}$ . VmP striae reaching the anterior border of the ventromental plate (Fig. 11) (species occurs in stagnant, sometimes slow flowing, waters)	<i>defectus</i>
-	LMP (Fig. 1) 135-170 $\mu\text{m}$ . VmP length (Fig. 4) 35-50 $\mu\text{m}$ (larvae from The Netherlands: 48-50). Head length 395-460(?) $\mu\text{m}$ . VmP striae at most reaching $\frac{3}{4}$ the length of the ventromental plate (Fig. 12); species occur in running waters	6
6	Fourth lateral tooth of mentum lower than the fifth tooth (Fig. 12). LMP 135-170 $\mu\text{m}$ . VmP striae rather close along each other (Fig. 12). Head length 395-460 $\mu\text{m}$ . Head width 265-325 $\mu\text{m}$ . Antennal segment 2 and 3 about equally long. SSd (Fig. 3) reaches till the top of the first lateral tooth (species occur only in rivers)	<i>rostratus</i>
-	Fourth lateral tooth of mentum equally high as the fifth tooth (Fig. 13). LMP 150 $\mu\text{m}$ . VmP striae relatively coarse and widely separated; some lines converge to one line (Fig. 13). Head length unknown. Head width ? 380 $\mu\text{m}$ . Antennal segment 2 shorter than segment 3. SSd (Fig. 3) reaches to $\frac{3}{4}$ of the second lateral tooth (species occurs in brooks, very rare)	<i>denticulatus</i>
7	Postoccipital margin ventrally pale or darkish yellow. Head length (680)800-900 $\mu\text{m}$ . LMP (Fig. 1) 330-360 $\mu\text{m}$ . VmP length 70-75 $\mu\text{m}$ . Mandible length 225-270 $\mu\text{m}$ (Fig. 15)	<i>redekei</i>
-	Postoccipital margin ventrally pigmented, as dark as the mandible teeth. Head length 600-820 $\mu\text{m}$ . LMP (Fig. 1) 238-250 $\mu\text{m}$ . VmP length 55-65 $\mu\text{m}$ . Mandible length $\pm$ 215 $\mu\text{m}$ (Fig. 14)	<i>psittacinus</i>

**Tab. 5: *Cryptochironomus* larvae. Important characters and measurements: look p. 17**




Fig. 11: *Cryptochironomus defectus*. Mentum with ventromental plate (Vallenduuk)

Fig. 12: *C. rostratus*. Mentum with ventromental plate (Vallenduuk)

Fig. 13: *C. denticulatus*. Mentum with ventromental plate (Vallenduuk)

Fig. 14: *C. psittacinus*. Mandible (Vallenduuk)

Fig. 15: *C. redekei*. Mandible (Vallenduuk)

## 7 Key for pupal exuviae of *Cryptochironomus*

### Introduction

The pupal exuviae of the genus are easily distinguished from other genera. The key is based mainly on Langton (1991). In most cases we were able to use our own material through collection from the water surface or by rearing larvae to adults. The terminology is based on Sæther (1980) and Langton (1991).

### Methods

The exuviae are put into a small Petri dish with 70 % alcohol. Identification is possible with magnifications of 40-100x. If higher magnification is required, one can mount the exuviae in alcohol on a slide.

A distinct character for *Cryptochironomus*, except for *C. ussouriensis*, is the forked anal processus between the anal lobes. In most species the cephalic tubercles are hooked, making accurate measurements of them difficult. The segments have a dorsal plate, the tergite, and a ventral plate, the sternite.

### Terminology

Anal lobe (Fig. 16)

Anal process (Fig. 16a)

Cephalic tubercle (Fig. 20)

Dorsal anterior thoracic mound (Fig. 19a)

Dorsal posterior thoracic mound (Fig. 19c)

Genital sac (Fig. 16b)

Hook row on tergite II (Fig. 17b)

Pedes spurii B (Fig. 17a)

Thoracic horn (Fig. 19d)

Thorax (Fig. 19)

Ventral anterior thoracic mound (Fig. 19b)

Morozova found larvae in Russia (Saratov) which look very like *Cryptochironomus redekei* in all instars, but might belong to a new species. More about this in the revision of the genus *Cryptochironomus* by Morozova (in preparation).




Fig. 16: *Cryptochironomus defectus*. Anal lobe (female). a = forked anal process, b = genital sac (length) (Morozova)

Fig. 17: *C. defectus*. Tergite II. a = pedes spurii B (PSB), B = hook row (Morozova)

Fig. 18: *C. obreptans*. Sternite I. a = anterior hump, b = antero-lateral hump, c = posterior point row (Morozova)

Fig. 19: *C. obreptans*. Thorax. a = dorsal anterior mound, b = ventral anterior mound, c = posterior mound, d = thoracic horn without setae (Morozova)

Fig. 20: *C. obreptans*. Cephalic tubercles. a = dorsal view, b = lateral view (Morozova)

### Key to the exuviae

- 1 Hook row on tergite II with a gap (Fig. 23). The anal process is forked (Fig. 16). Thoracic horn has a main stem and 3-4 additional bundles of branched filaments 2
  - Hook row on tergite II is complete (Fig. 21). The anal process is triangular (Fig. 22). Thoracic horn comprises a main stem and a single bundle of branched filaments *ussouriensis*
- 2 Tergites generally without reticulation (Fig. 23) 3
  - Tergites with a fine (Fig. 24) or thick (Fig. 25) reticulation 6
- 3 Thorax with only a ventral anterior mound (Fig. 26) 4
  - Thorax with a dorsal and ventral anterior mound (Fig. 27) *obreptans*
- 4 Pedes spurii B present (Fig. 23). Exuviae brownish-yellow. Granulation on the anterior cephalothorax relatively small, at most 8  $\mu\text{m}$  *defectus*

Note: This species is very similar to *albofasciatus* and *supplicans*

  - Pedes spurii B absent. Exuviae yellow or golden-yellow. Granulation on the anterior cephalothorax relatively large, 10  $\mu\text{m}$  or more 5
- 5 *albofasciatus* and *supplicans*

Both species are difficult to identify because they vary so much. Further characters are needed

  - a Cephalic tubercle at the apex mostly straight, its length 180-255  $\mu\text{m}$  (Fig. 28). Gap of the hook row 55-135  $\mu\text{m}$  (Fig. 30). Gap in hook row half the length of hook row at one side or more. Granulation on the anterior cephalothorax crowded and forming ridges on the antepronotum *albofasciatus*
  - b Cephalic tubercle mostly hooked (sometimes straight) at the apex, its length 275-515  $\mu\text{m}$  (Fig. 29). Gap of the hook row 20-90  $\mu\text{m}$  (Fig. 31). Gap in hook row much less than half the length of hook row at one side. Granulation on the anterior cephalothorax widely separated *supplicans*




Fig. 21: *Cryptochironomus ussouriensis*. Tergite II (Shilova 1959, Pankratova 1983)

Fig. 22: *C. ussouriensis*. Tergite VIII and anal lobe (Shilova 1959, Pankratova 1983)

Fig. 23: *C. defectus*. Tergites II, III (Morozova)

Fig. 24: *C. Pe1*. Reticulation on tergite IV (Langton 1991)

Fig. 25: *C. psittacinus*. Reticulation on tergite IV (Langton 1991)


Fig. 26: *C. supplicans*. Thorax (Morozova)

Fig. 27: *C. obreptans*. Thorax (Morozova)

Fig. 28: *C. albofasciatus*. Cephalic tubercles (Morozova)

**Fig. 29: *C. supplicans*. Cephalic tubercles (Langton 1991)**

- 6 Reticulation thick (Fig. 25). Pedes spurii B without groups of small points. (Fig. 33) 7
- Reticulation very fine (Fig. 24). Pedes spurii B with groups of small points (Fig. 32). (Gap in the hook row about the same length as the length of a single half row. Reticulation on tergites I-VI, less clear on VII and VIII, with small points) *denticulatus*
- 7 Pedes spurii B present. Gap in the hook row is equal in length or even longer than the length of a single half row (Fig. 33) 8
- Pedes spurii B absent. Gap in the hook row does not exceed the length of a single half row (Fig. 34) 9
- 8 Points of posterior transverse row on tergites I-VII are yellowish-brown. Walls of the reticulation without points (as in Fig. 25) *rostratus*
- Points of posterior transverse row on tergites I-VII are dark brown to black. Walls of the reticulation with small points (Fig. 35) *crassiforceps*
- 9 Reticulation complete on tergites I-VI, partial on VII and VIII. Posterior thoracic mound mostly with two swellings *redekei*
- Reticulation complete on tergites I-V, partial on VI. Posterior thoracic mound always with one swelling *psittacinus*

**Fig. 30: *Cryptochironomus albofasciatus*. Hook row of tergite II (Langton 1991)****Fig. 31: *C. supplicans*. Hook row of tergite II (Langton 1991)****Fig. 32: *C. denticulatus*. Tergite II, PSB with points (Langton 1991)****Fig. 33: *C. rostratus*. Hook row on tergite II (Langton 1991)****Fig. 34: *C. redekei*. Hook row on tergite II (Morozova)**



continued		VmPL	n	LMP	n	A1	n	S3-S3	n	S3/A1	n	Mand.L	n
ussouriensis	RUS	75-80	4	350-370	4	90-100	4	100-105	2	1,0-1,11	2	250-260	4

## 8 Supplementary notes to the European species of *Cryptochironomus* with particular reference to The Netherlands and the Volga region of Russia

### *Cryptochironomus albofasciatus* (Staeger, 1839)

Langton reports that this species corresponds to *albofasciatus* of Edwards (1929), Reiss (1968) and Pinder (1978) and *defectus* of Shilova (1966)

Shilova (1966) gives characters for the adult, but not for the pupal exuviae; Pankratova (1983) gives a description of the exuviae. There are some doubts as to whether their identification is correct. Here, the characters given by Langton & Visser (2003) have been adopted. So far the authors have not reared larvae which gave such pupal exuviae.

### *Cryptochironomus crassiforceps* Goetghebuer, 1931

Distribution: Caspian Sea region. The species is unlikely to occur in Western Europe. The larvae are unknown.

### *Cryptochironomus defectus* (Kieffer, 1913)

The pupal exuviae are nearly identical to those of *C. supplicans*. However, the larvae more resemble those of *C. rostratus*.

In The Netherlands the larvae occur in many water types, such as waters in cities, watercourses and pools. The substratum is always sandy mixed with some detritus and fine organic material. *C. defectus* is often found with *C. obreptans* and *C. supplicans*. Together with *C. rostratus*, *C. defectus* also occurs in slow flowing parts of rivers. Within the genus a common species.

### *Cryptochironomus denticulatus* (Goetghebuer, 1921)

This species has until now been known only from England (information Dr. P. Langton). However, Ashe & Cranston (1990) mention Belgium (Virton) and England. Klink (pers. comm.) reports this species from the Meuse at Hastière (Belgium).

### *Cryptochironomus obreptans* (Walker, 1856)

The pupal exuviae are easily distinguished from those of the other species. Recognizing the larvae, on the other hand, is rather difficult. They are very like those of *C. supplicans*. Regularly both species are found at the same sites, often with *C. defectus*.

In The Netherlands they occur in many water types, such as waters in cities, watercourses, borders of canals and pools. The habitats always have sandy substrata mixed with some detritus and fine organic material. Pupal exuviae have also occurred in large rivers. Within the genus a common species.

### ***Cryptochironomus psittacinus* Meigen (1830)**

Within the genus a large species. The pupal exuviae are easy to distinguish from *C. redekei*, but that is not so with the larvae. Shilova (1966) and Langton (1991) give characters for this species, but the descriptions differ. Langton's interpretation is followed here.

This species is known from deep infiltration waters as well as from shallow pools. Within the genus an uncommon species.

### ***Cryptochironomus redekei* (Kruseman, 1933)**

Within the genus a large species. The pupal exuviae are easily distinguished from those of *C. psittacinus*, but that is not so with the larvae. Shilova (1966) and Langton (1991) give characters for this species, but the descriptions differ. Langton's interpretation is followed here.

Although this species has been found once by the authors together with *C. psittacinus*, that would appear to be a rare exception.

Adults and pupal exuviae are known from larger shallow pools. Vallenduuk collected also larvae in shallow pools. However there are larvae collected in a deep infiltration water. Within the genus a rare species.

### ***Cryptochironomus rostratus* Kieffer, 1921**

For identification of this species we used the characters of Langton (1991). Shilova mentioned pupae from Uzbekistan (Russia) and gave a drawing with differing reticulation. That material is unfortunately lost.

In The Netherlands this species is found at several locations on the river Waal. Pupal exuviae are regularly found in the rivers Rhine and IJssel; once from the river Meuse. The larvae are always found in slow flowing parts, the substratum containing detritus and some fine organic material. Within the genus a rather common species.

### ***Cryptochironomus supplicans* (Meigen, 1830)**

The pupal exuviae of this species are easily confused with those of *C. defectus*. The larvae of both species, however, are easily distinguished. Larvae of *C. supplicans* and *C. obreptans* differ very little. Shilova (1966) and Langton (1991) give characters for this species, but the descriptions differ. Langton's determination

is here followed. Nevertheless the cephalic tubercles depicted (Fig.13 and 14) by Langton (1991) differ from our material.

In comparison with *C. defectus* this species is found much less frequently, though both species regularly occur at the same sites. Pupal exuviae are known also from the rivers IJssel and Meuse. In The Netherlands the species is found in many water types. Within the genus a common species, perhaps less common than *C. obreptans*.

### *Cryptochironomus ussouriensis* (Goetghebuer, 1933)

Only known from Russia. It is not to be expected in western Europe.

#### Acknowledgements

We wish to thank Dr. H. M .P. Moller Pillot, Tilburg/NL for helpful comments. With pleasure we want to thank Dr. P. Langton for his co-operation.

#### References

- Ashe, P. & P. S. Cranston (1990): Psychodidae/Chironomidae.- In: Soos A. & L. Papp (eds): Catalogue of Palaearctic Diptera 2: 1-355, Budapest
- Beck, E. C. & W. M. Beck (1969): Chironomidae (Diptera) of Florida.III. The Harnischia complex (Chironominae).- Bulletin of the Florida State Museum biological sciences 13: 277-313, Gainesville, Fla.
- Chernovski, A.A., 1949. Identification of larvae of the midge family Tendipedidae.- Opredeliteli po Faune SSSR izdavaemue Zoologiceskim institutom Akad. Nauk SSSR 31: 1-185, Moskva (in Russian)
- Cranston, P.S. & F. Reiss (1983): The larvae of Chironomidae (Diptera: Chironomidae) of the Holarctic region. Keys to subfamilies.- In: Wiederholm (ed.). Chironomidae of the Holarctic region - Keys and diagnoses. Part 1. Larvae- Entomologica Scandinavica Supplementum 19: 11-15, Copenhagen
- Curry, L. (1958): Larvae and pupae of the species of *Cryptochironomus* (Diptera) in Michigan.- Limnology and Oceanography. 3: 427- 442, Lawrence, Kansas
- Edwards, F. (1929): British non biting midges (Chironomidae).- Transactions of the Entomological Society of London 77: 279-430, London
- Fittkau, E. & F. Reiss (1978): Chironomidae.- In: Illies (ed.): Limnofauna Europaea. 2.Aufl.: 404-440, G. Fischer Stuttgart
- Goetghebuer, M. (1937): 13c. Tendipedidae (Chironomidae). b) Subfamilie Tendipedinae (Chironominae).A. Die Imagines.. In: Lindner, E: (ed.): Die Fliegen der paläarktischen Region. 3,2, Lief. 107:1-138, Stuttgart
- Harnisch, O. (1923): Metamorphose und System der Gattung *Cryptochironomus* K.- Zoologische Jahrbücher Abt. Systematik, Geographie und Biologie der Tiere 47: 271-308, Jena
- Izvekova, E.I. (1975): Feeding and feeding relations of larvae of mass species of chironomids of the Uchinsk reservoir.- Moscow State University, 1-18 (in Russian)
- Izvekova, E.I. (1980.): Ecology of chironomids (Diptera, Chironomidae). Feeding.- In: Benthos Uchinskogo vodochranilishcha. Moscow, ed. Nauka: 72-100. (in Russian)
- Kieffer, J. J. (1913): Nouvelle contribution a la connaissance des Tendipedides d'Allemagne.- Bulletin de la Societé d'Histoire Naturelle de la Moselle 28: 16, Metz
- Kieffer, J. J. (1918): Beschreibung neuer auf Lazarettsschiffen des östlichen Kriegsschauplatzes und bei Ignalino in Litauen von Dr. W. Horn gesammelten Chironomiden mit Übersichtstabellen

- einiger Gruppen von palaarktischen Arten.- Entomologische Mitteilungen 7: 35-53, Berlin-Dahlem
- Kieffer, J. J. (1921): Neue Chironomiden aus Mitteleuropa. - Archiv für Hydrobiologie Supplement 2: 785-808, Stuttgart
- Kiknadze, I.I., A.I. Shilova, I.E. Kerkis et al. (1991): Kariotipy i morfologiya lichinok triby Chironomini. Atlas.- Nauka, Sibirskie otdelenie. Novosibirsk: 1-115. (in Russian)
- Klink, A. & H. K. M. Moller-Pillot (1996): Lijst van de Nederlandse Chironomidae, bijgewerkt tot 1 januari 1996. Themanummer 08: 1-17. Werkgroep Ecologisch Waterbeheer.
- Kruseman, G. (1933): Tendipedidae Neerlandicae. Pars I. Genus *Tendipes* cum generibus finitimis.- Tijdschrift voor Entomologie, 76: 119-216, s'Gravenhage
- Langton, P. H. (1991): A key to the pupal exuviae of West Palaearctic Chironomidae. Coleraine.-386 pp., Cambridgeshire.
- Langton, P. H. & H. Visser (2003): Chironomidae exuviae. A key to pupal exuviae of the West Palaearctic region.- CD-Rom, ETI/STOWA/RIZA, Amsterdam
- Lenz, F. (1921): Chironomidenpuppen und -larven. Bestimmungstabellen.- Deutsche Entomologische Zeitschrift 3: 148-162, Berlin
- Lenz, F. (1923): Die Vertikalverteilung der Chironomiden im eutrophen See.- Internationale Vereinigung für theoretische und angewandte Limnologie. Verhandlungen 1: 144-167, Stuttgart
- Lenz, F. (1926): Chironomiden aus dem Balatonsee.- Archivum Balaticum. 1: 129-144, Tihany
- Lenz, F. (1935): Die Metamorphose von *Cryptochironomus lateralis* Goetgh.- Zoologischer Anzeiger 109(3/4): 81-84, Leipzig
- Lenz, F. (1941): Die Metamorphose der Chironomiden-Gattung *Cryptochironomus*.- Zoologischer Anzeiger 133: 29-41, Leipzig
- Lenz, F. (1954-62): Tendipedidae (Chironomidae).b. Subfamilie Tendipedinae. B. Die Metamorphose der Tendipedinae.- In Linder, E. (ed.). Die Fliegen der palaarktischen Region. Stuttgart. 13c: 139-250, Stuttgart
- Lindeman, R. L. (1941): Seasonal food-cycle dynamics in a senescent lake.- American Midland Naturalist 26(3): 636-673, Notre Dame, Ind.
- Lipina, N. N. (1928): Larvae and pupae of chironomids. Ecology and systematics.- Izdatelstvo Instituta rybnogo khozyaistva 3: 1-179, Moscow (in Russian)
- Luferov, V. P. (1956): Some data on the behaviour of predatory Tendipedidae larvae.- Doklady Akademii Nauk SSSR 3: 466-469, Moskva (in Russian)
- Luferov, V. P. (1958): On the feeding relations of predatory Tendipedidae in the Rybinsk reservoir.- Byulleten Instituta biologii vodohranilish AN SSSR, 2: 16-19, Moscow (in Russian)
- Moller-Pillot, H. K. M. & R. F. M. Buskens (1990): De larven der Nederlandse Chironomidae (Diptera).- Nederlandse Faunistische Mededelingen 1C. Autekologie en verspreiding: 1-87, Stichting European Invertebrate Survey Nederland, Leiden.
- Muttkowski, R. A. (1918): The fauna of Lake Mendota: A qualitative survey with special reference to insects.- Trans. wis. Acad. Sci. Arts Lett. 19: 374-482, Riga
- Pagast, F. (1931): Chironomiden aus der Bodenfauna des Usma-See in Kurland.- Folia Zoologica et Hydrobiologica 3:199-248, Riga
- Pagast, F. (1933): Über die Metamorphosestadien von *Chironomus vulneratus* Zett. (Gruppe *Cryptochironomus* s.str.).- Konowia 11: 155-161, Wien
- Pankratova, V. Y. (1983): Larven en poppen van de muggen van de subfamilie Chironominae van de USSR fauna.- Opredeliteli po Faune SSSR izdavaemue Zoologiceskim institutom Akad. Nauk SSSR 134: 153-166, Moskva
- Pinder, L. C. V. & F. Reiss (1983): The Larvae of Chironomidae (Diptera, Chironomidae) of the Holarctic region. Keys and diagnoses.- In: Wiederholm (ed.). Chironomidae of the Holarctic region - Keys and diagnoses. Part 1. Larvae.- Entomologica Scandinavica Supplementum 19: 293-435, Copenhagen

- Reiss, F. (1968): Okologische und systematische Untersuchungen an Chironomiden (Diptera) des Bodensees. Ein Beitrage zur lakustrischen Chironomiden fauna des nordlichen Alpenvorlandes.- Archiv für Hydrobiologie 64: 176-323, Stuttgart
- Saether, O. A. (1980): Glossary of chironomid morphology terminology.- Entomologica Scandinavica Supplementum 14: 1-51, Copenhagen
- Shilova, A. I. (1959): *Cryptochironomus ussouriensis* Goetgh. (= *nigridentis* Tshern.) and some dates about biology.- Trudy Instituta biologii vodohranilitsh ANSSSR 2: 109-116, Moscow
- Shilova, A. I. (1965): Metamorphose and biology of some species of the genus *Cryptochironomus*. - Voprosy Gidrobiologii. Tezisy dokladov I sezda VGBO 456-457, Moscow (in Russian)
- Shilova, A. I. (1966): Taxonomie van *Cryptochironomus* ex. Gr. defectus Kieff. (Diptera, Chironomidae).- Trudy Instituta Biologii Vnutrennich Vod 12: 214-238, Moskva (in Russian)
- Shilova, A.I. (1976): Chironomids of the Rybinsk reservoir.- 249 pp., (ed. Nauka), Leningrad (in Russian)
- Townes, H. (1945): The Nearctic species of Tendipedini (Diptera, Tendipedidae (Chironomidae).- American Midland Naturalist 34:1-206, Notre Dame, Ind.
- Werkgroep Exuviae (DZH, PWN, RIZA, WBB) (1996) Exuviae als biologische parameter.- N.V. Duinwaterbedrijf Zuid-Holland, Vorburg & N.V. PWN Waterleidingbedrijf rd-Holland, Bloemendaal
- Wesenberg-Lund, C. (1943): Biologie der Süßwasserinsekten.- 682 pp., (Springer) Berlin

*Addresses of the authors:* Dr. E. E. Morozova, Saratov State University, Chapaeva Street 6A-70, 410045 Saratov, Russia, email: moroz@san.ru

Henk J. Vallenduuk, Spoorven 431, 5464 PA Veghel, The Netherlands, email: hj.vallenduuk@tiscali.nl

*Received:* 2004-08-25