



CHIRONOMUS NEWSLETTER ON CHIRONOMIDAE RESEARCH

Co-Editors: Ruth CONTRERAS-LICHTENBERG

Naturhistorisches Museum Wien, Burgring 7, A-1014 WIEN, Austria

Peter H. LANGTON

5 Kylebeg Avenue, Mountsandel, Coleraine, Co.

Londonderry, Northern Ireland, BT52 1JN - Northern Ireland

Bibliography: Odwin HOFFRICHTER

Institut f. Biologie I, Albert-Ludwigs-Universität Freiburg, Hauptstrasse 1

D-79104, Germany

Treasurer: Trond ANDERSEN: Museum of Zoology,

University of Bergen, Museplass 3, N-5007 Bergen - Norway

ISSN 0172-1941

No. 13

September 2000

CONTENTS

Chironomid Work in Munich to Continue	1
New curator at the Zoologische Staatssammlung Munich	2
Contributions in SPIXIANA in Memory of Dr. Reiss.....	4
To Iya Kiknadze at 70.....	5
Current Research	7
Short – Communications	19
Notice Board	21
List of regional representatives 2000	24
Current Bibliography	26
Registration Form (International Conference: „Small Rivers“, Togliatti, RU, April 25-28, 2001)	43

CHIRONOMID WORK IN MUNICH TO CONTINUE

A year has now passed since the shocking, very premature death of Dr. Reiss whom so many of us had relied upon as a guiding light in chironomid science, not only around the world but of course especially in Germany and Munich. Naturally, having lost somebody who meant so much you can only hope that the void left behind may be somehow refilled, not on the personal level but at least partially with respect to knowledge and work. It was a second hard blow, therefore, when the management of ZSM decided, for various "objective" reasons, to not make chironomids a high priority in hiring a new leader of the Diptera section.

However, we are very glad to be able to report to you that the new curator chosen, Dr. Marion

Kotrba, is doing her best to maintain the relevant resources at ZSM as available and useful to all of us in the chironomid community as they have been, although the chironomids are (so far) not part of her personal research interests. Please look for the separate article by Dr. Kotrba in this CHIRONOMUS, in which she introduces herself to you.

Meanwhile, research on chironomids has been continuing at and around ZSM. Here we aim to give a brief overview of current activities. Also in this CHIRONOMUS issue you can find an announcement of papers in memory of Dr. Reiss which are being published this year in the journal SPIXIANA. For more information on the ZSM Diptera section and the people associated with it, please visit the website at

http://www.zsm.mwn.de/wiss_start.htm and select Diptera.

For the past year, the chironomid group at ZSM has consisted of four doctoral candidates, all supervised by Prof. E. J. Fittkau.

Angela M. Sanseverino from Brazil is in Munich on a fellowship from the German Academic Exchange Service (DAAD). She is in her second year of work on the taxonomy, systematics and phylogeny of Neotropical *Tanytarsus* VAN DER WULP, including a full evaluation of existing species and groups, and the description of new species.

Elisabeth Stur and Sofia Wiedenbrug have recently completed their respective theses and are now, among other things, working together on the chironomids of groundwater springs in Berchtesgaden National Park in the German Alps. This three-year project involves rearing larvae in situ and determining species from one year of continuous emergence trapping at two different localities.

Also, Martin Spies hopes to wrap up his academic program this year. The last publications from his study of nuisance midges in southern California have been published or are in press (e.g., a paper on the general results in Odwin Hoffrichter's Proceedings from the 1997 International Symposium in Freiburg, and "A contribution to the knowledge of Holarctic *Parachironomus* LENZ (Diptera: Chironomidae), with two new species and a provisional key to Nearctic adult males" just published in Tijdschrift voor Entomologie 143).

For the entire year 2000 we have been happy to have another Munich graduate, Prof. Dr. Rodulfo Ospina Torres from Bogotá (Colombia), back here at ZSM on a one-year fellowship from the Humboldt Foundation. Rodulfo has been mostly sorting out the taxonomy of many adult specimens collected in Andean streams. He and Sofia are also

collaborating on publishing excerpts from their extensive doctoral dissertations on Neotropical pupal exuviae.

A new project taken up by Martin Spies has developed out of a recent, EU-funded visit to Paris to study the chironomids in the J. W. Meigen collection kept at the French National Museum of Natural History since 1840. Meigen established a significant share of the oldest chironomid names still in use today, especially for central Europe. In spite of this, a number of current interpretations of Meigen's species appear not to be properly based on the specimens in his collection. For example, there are several supposed type series from which no specimen has apparently ever been slide-mounted to verify the species. On the other hand, erroneous type designations have been subsequently published in some cases, e.g. selecting a male as lectotype although Meigen's original description treats the female only.

Many specimens have been taken to Munich to reevaluate their specific identities and type status. This will involve searching out and comparing further material determined by Meigen reportedly existing at other institutions. Concurrently, a comprehensive review of chironomid names published until 1840 is under way. The long-term goal is to strengthen the taxonomic basis of all chironomid work by tying species names to appropriate type specimens wherever necessary and possible, and by integrating the presently scattered or outdated information on chironomid taxa and their types in a global name list and catalog.

Angela Sanseverino, Elisabeth Stur,
Sofia Wiedenbrug, Rodulfo Ospina
and Martin Spies
c/o Zoologische Staatssammlung
München Münchhausenstr. 21, D-
81247 München, Germany

NEW CURATOR OF DIPTERA AT THE ZOOLOGISCHE STAATSSAMMLUNG MUNICH

Dear Chironomid Workers,

The ZSM houses one of the most extensive and important chironomid collections of the world. It comprises about 1/3 of all described chironomid species and about 2/3 of those known from the Palaearctic, including type material of about 500 species. The extensive library of chironomid literature is almost complete for publications on taxonomy and

systematics of the family. These collections are predominantly based on the material generated by A. Thienemann and collaborators, E. J. Fittkau, and F. Reiss. During their tenures in Munich, Drs Fittkau and Reiss have turned the ZSM's Diptera section into a „hot spot“ of chironomid research. In addition to their own extremely productive research activities they attracted and supervised countless graduate

and postgraduate students and cooperated with colleagues all over the world.

In April 2000 I was appointed the new curator of Diptera at the ZSM. As many of you know this position had been vacant since the tragic and premature death of Dr. Friedrich Reiss in August 1999 (obituary in Chironomid Newsletter 12). Although I am myself not a chironomid worker (see CV below), I will do my best to manage and further extend the chironomid collection at the ZSM, and I extend a sincere invitation to all of you who wish to

work with our material. I would especially like to point out that there is still extensive undescribed material from all regions of the world, much of it already dissected and mounted. Many new species are sorted out and await their study and description!

Please do contact me at marion.kotrba@zsm.mwn.de if you need any further information. You can also visit our homepage at <http://www.zsm.mwn.de>

CV Marion Kotrba

1957 born in Nuremberg, Germany

1985 University of Regensburg, Diploma in Zoology. Thesis: „Studies on female selection in *Cyrtodiopsis whitei* CURRAN (Diopsidae, Acalyptrata, Diptera).“

1992 University of Regensburg, Dr. rer. nat. magna cum laude. Thesis: „The reproductive system of *Cyrtodiopsis whitei* (Diopsidae, Diptera), especially regarding the internal female reproductive tract.“

1993-94 Postdoctoral Fellow, Department of Zoology, University of Maryland. Research topic: „Phylogeny of the higher flies (Diptera, Brachycera), and especially the acalyptrate Schizophora, by comparative morphology of the internal female reproductive tract“ (Research stipend DFG).

1994-96 Postdoctoral Fellow, National Museum of Natural History, Smithsonian Institution, Washington DC, Department of Entomology (Research stipend DFG / Postdoctoral Fellowship SI).

1997-2000 Curator of Diptera, Museum of Natural History, Berlin.

2000 Curator of Diptera, Zoologische Staatssammlung, Munich.

Research Collaborator of the National Museum of Natural History, Smithsonian Institution, Washington DC, Department of Entomology.

Fields of interest: Phylogeny and systematics of Diptera, especially of the acalyptrate Schizophora; reproductive biology; comparative morphology and functional morphology of the internal female reproductive tract; sperm transfer, sperm storage, and fertilization; sperm competition; postcopulatory female choice; viviparity; phylogeny and biology of Diopsidae; convergent evolution of head projections such as antlers and stalk eyes in Diptera.

Research trips: Jamaica (1982), Malaysia (1986), South Africa (1992, 1994), Costa Rica (1995), Ecuador (1999).

Dr. Marion Kotrba, Sektion Diptera, Zoologische Staatssammlung
München, Münchhausenstr. 21, D-81247 München, Germany
Tel.: +49 89 8107 147 Fax.: +49 89 8107 300

PUBLICATION ANNOUNCEMENT

Contributions in SPIXIANA in Memory of Dr. Reiss

In July of this year, the special publication announced in CHIRONOMUS No. 12 has appeared as SPIXIANA volume 23, issue 2. Although authors had to add this to all their other work under tight time constraints, the editors received more contributions than could be fit within a regular SPIXIANA, even though the issue was enlarged from the normal 96 to 128 pages. Some papers thus had to be delayed until the next issue to be published in November 2000.

To all colleagues involved in various parts of this production: thank you very much again!

You have made this a great tribute to the man we have had to say farewell to. Due to the contributions received this publication contains almost exclusively original scientific papers, but we believe Dr. Reiss himself would have preferred this to eulogies of his person. He would have enjoyed applying them in practical work, and so, we hope, will the readers.

Here is an excerpt from the table of contents, giving authors' names and abbreviated titles of the scientific papers:

- BAEHR, M. & M. SPIES (eds) 2000: Contributions to chironomid research in memory of Dr. Friedrich Reiss. – SPIXIANA 23(2).
- PETER S. CRANSTON: *Austrobrillia* Freeman: immature stages, and new species from the Neotropics
- PATRICK ASHE: *Reissmesa*, nom. nov., a replacement name for *Reissia* Brundin
- EUGENYI A. MAKARCHENKO: *Cricotopus (Cricotopus) reissi*, spec. nov. from Chukchi Peninsula, northeastern Russia
- SEBASTIÃO J. DE OLIVEIRA : A new, non-marine species of the genus *Thalassomya* Schiner, 1856
- CLAUS ORENDT: Chironomids of small Alpine water bodies (springs, spring brooks, pools, small lakes) of the northern Calcareous Alps
- RICHARD E. JACOBSEN & SUE A. PERRY: A review of *Beardius* Reiss & Sublette, with description of a new species from Everglades National Park, Florida
- JAMES E. SUBLETTE & MIR S. MULLA: *Chironomus strenzkei* Fittkau, a new Pan-American distribution, with a review of recent similar additions to the Nearctic midges
- WOLFGANG F. WÜLKER & JON MARTIN: Northernmost *Chironomus* karyotypes
- KALMAN BIRÓ: Chironomidae from Hungary 2. New records of *Lipiniella moderata* Kalugina, 1970
- MARIA C. MESSIAS: *Oukuriella reissi*, a new species of the genus *Oukuriella* Epler, 1986
- BROUGHTON A. CALDWELL: First Nearctic record of *Neostempellina* Reiss, with description of a new species
- SUSANA TRIVINHO-STRIXINO & GIOVANNI STRIXINO: A new species of *Caladomyia* Säwedal, 1981, with description of the female and immature stages
- NICOLA REIFF: Review of the mainly Neotropical genus *Caladomyia* Säwedal, 1981, with descriptions of seven new species
- PETER H. LANGTON & XAVIER-FRANÇOIS GARCIA: A review of *Cladotanytarsus conversus* (Johannsen) with first records from Europe
- ANGELA M. SANSEVERINO & SOFIA WIEDENBRUG: Description of the pupa of *Tanytarsus cuieirensis* Fittkau & Reiss
- MALCOLM G. BUTLER: *Tanytarsus aquavolans*, spec. nov. and *Tanytarsus nearcticus*, spec. nov., two surface-swarming midges from arctic tundra ponds
- ELISABETH STUR & TORBJØRN EKREM: *Tanytarsus usambarae*, spec. nov. from West Usambara Mts., Tanzania, East Africa

Papers to follow in SPIXIANA 23(3), November 2000:

ASHE, P., J. P. O'CONNOR & D. A. MURRAY: Larvae of *Eurycnemus crassipes* (Panzer) (Diptera, Chironomidae) ectoparasitic on prepupae/pupae of *Hydropsyche siltalai* Döhler (Trichoptera, Hydropsychidae), with a summary of known chironomid/trichopteran associations

KYEREMATEN, R. A. K., T. ANDERSEN & O. A. SÆTHER: A review of Oriental *Rheotanytarsus* Thienemann & Bause, with descriptions of some new species

PAGGI, A. C. & D. AÑON SUAREZ: A new species of *Ablabesmyia* from Rio Negro province, Argentina, with descriptions of the adult female and preimaginal stages

PAPUSHEVA, E., V. PROVIZ, A. BLINOV & B. GODDEERIS: Phylogeny of the endemic Baikalian *Sergentia*

If you are interested in ordering, please note that approximately 50% of issue 3 will consist of papers not involving chironomids.

The two issues are available only from the publishers of SPIXIANA (they are doing all of us a favor by handling distribution!): Verlag Dr. Friedrich Pfeil, Wolfratshauer Str. 27, D-81379 München, Germany E-mail: 100417.1722@compuserve.com; Fax: +49 89 - 72 42 772 Tel.: +49 89 - 74 28 270

The price for issues 2+3 together is Euro/US\$ 40 plus postage, issue 2 alone is Euro/US\$ 30

plus postage, issue 3 alone is Euro/US\$ 20 plus postage. These prices are based on the approximate equivalence of Euro and US\$ as of June 2000. Should the exchange rate significantly differ by the time of your payment, the publisher reserves the right to modify prices accordingly. Postage overseas is not cheap, collective rather than individual orders could significantly reduce cost as well as the work load at SPIXIANA. Credit card payment is possible. Please, contact Dr. Pfeil for all further details.

Martin Spies
Munich, Germany

TO IYA KIKNADZE AT 70

On February 9th, 2000, we celebrated the 70th



anniversary of Doctor of Biological Sciences, Professor of the Novosibirsk State University, Honoured Scientist of the Russian Federation and founder of both the Siberian school of cytogenetics and the

laboratory of Cell Biology, Institute of Cytology and Genetics, Novosibirsk, Russia, Iya Iwanowa Kiknadze.

Iya Kiknadze was one of the first scientists to come to Novosibirsk in January 1958. She was personally invited by Aleksandra Prokofeva-Belgovskaya, the instigator of Soviet cytogenetics, who participated actively in the organisation of the Institute of Cytology and Genetics. By that time, Iya Kiknadze was already a mature researcher. Having graduated

from the biological Department of Leningrad University and getting her Ph. Degree in 1955, she was thoroughly trained in cytology, working in the Laboratory of Cell Physiology, organised by Dimitry Nikolaevich Nasonov, with the Zoological Institute of the Soviet Academy of Sciences, under the guidance of the oldest Russian cytologist Ivan Ivanovich Sokolow.

When she settled in Novosibirsk, Iya Kiknadze became actively involved in the direction of cytological research within the Institute of Cytology and Genetics. She proposed a long-term program on investigating interphase chromosomes, chromomeres and nucleolus, whose actual existence were peremptorily rejected during those years when Lysenko's pseudoscientific theory was still governing. This direction is still efficiently developed at the Institute. When the Laboratory of General Cytology was combined with the Institute of Cytology and Genetics, Iya Kiknadze became its head and held that position for over 30

years. These years brought to life and put into orbit a constellation of her numerous apprentices and followers. The works of the Laboratory of General Cytology have attracted the attention of domestic and foreign researchers since the beginning of the 60s. Young scientists not only from the entire soviet Union, but also from Poland, Bulgaria, Czechoslovakia, Germany and India came to learn from Iya Kiknadze. Students of hers became candidates and doctors of science, leading researchers and heads of laboratories, and can be met in many countries of the world.

The prime scientific love of Kiknadze, which she never deserted, is the polytene chromosome of Chironomidae. She has studied the polytene chromosomes from different standpoints. First, she was interested to find out how the polytene chromosome was organised and how it functioned, and pioneered together with the laboratory staff in solving this problem. The work performed resulted in a concept of discrete location of genes in the chromosome. It was demonstrated that the gene activities changed at different stages of cell life and that puffs – loose, swollen regions of polytene chromosomes – were actively working genes. This allowed Kiknadze to formulate the concept of the polytene chromosome as a model of the interphase chromosome. The next stage of her work was focused on investigation of the products of gene function. Study of the tissue-specific genes of Balbiani rings - the most actively working regions of the polytene chromosomes – required development of new research methods, some of them still remaining unique. Most important issue of this work was the insight into molecular cytological organisation of the tissue-specific Balbiani ring Bra; isolation of the tissue-specific protein encoded by it, and isolation, cloning and molecular analysis of the corresponding DNA region. The results obtained were presented in detail at the International Symposium on Organisation and Expression of tissue-specific Genes (Novosibirsk, Akademgorodok, 1982), organised under her leadership. This symposium underlay biennial International

Workshops on Balbiani rings held in different parts of the world.

Studies on the system of Balbiani rings in polytene chromosomes of different chironomid species required comparison of their chromosome sets. This investigation gave Iya Kiknadze the understanding of the course of chromosome evolution within the Chironomidae and the role of chromosome rearrangements during specification. Iya Kiknadze and her colleagues have so far thoroughly studied the karyotype of over 150 chironomid species. Two monographs on the extensive data obtained became table books for both Russian and foreign researchers-chironomidologists.

Iya Kiknadze worked at the Chair of Cytology and Genetics with the Novosibirsk University from its very beginning. Her first students still remember her exciting and inspirational lectures on general cytology. She was their author as at that time no other university in this country gave such a course.

Typical of Kiknadze is a phenomenal capacity for work, which is still envied by young people. Everything she ever wrote is well set, fundamental, accurate and reliable. Many of her works have become classics. Iya Kiknadze is known and loved all over the world and many well-known scientists from Australia, Austria, Belgium, Bulgaria, Germany, the Netherlands and USA count working in her team as an honour.

No age can make her young spirit older or diminish her plans and designs. Iya Kiknadze is still queen not only in scientific activities; she can do everything - organise a celebration, go on a walking tour, to a theatre, museum or concert hall – she is always the leader, and everybody follows her. We all wish her good health, inexhaustible energy and constant interest in Life and Science. Dear Iya Ivanovna, we wish you many more interesting years full with new designs and successes.

Staff of the Laboratory of Cell Biology, Institute of Cytology and Genetics, Siberian Branch of the Russian Academy of Sciences.

CURRENT RESEARCH

A POINT OF VIEW ON CHIRONOMID DEFORMITIES INVESTIGATION

By Larisa Nazarova

Kasan State University, Kazan, Russia

Since 1995, investigations connected with different aspects of chironomid ecology were started in Kazan State University, focusing in particular on influence of different pollutants on chironomid morphology in natural waterbodies as well as in laboratory conditions. Although chironomid deformities have been investigated for a long time already practically everywhere else in the world, Russia has not been much involved in this specific and very interesting study. There are only a very few Russian references briefly reporting the occurrence of chironomid deformities and I am sure they are absolutely unknown to foreign specialists (e.g. SKALSKAYA 1994; ZINTCHENKO et al. 1997).

But the time is really past for the mere description of a new (even very lovely) deformity. This is the time to seek out and try to understand the internal mechanisms involved; relationships between biochemistry and physiology of organisms are very important now. How do the pesticides, trace metals (acknowledged teratogens) and other chemicals, which are supposed to be teratogens „work“. What are their targets in metabolic processes, which of them exert an influence at the genetic level and is this influence reversible or not?... and so on. These are the kind of question available for specialists fascinated by this area of chironomid study. It is not possible to forget the valuable contribution of Dr. Warwick, who formulated basic approaches and methods of investigation in this area (WARWICK 1985, 1989, 1990, 1991 etc). Also well known are brilliant investigations of Canadian, Belgian and other specialists (VAN DE GUCHTE & VAN URK 1989; JANSSENS DE BISTHOVEN et al. 1992; MADDEN et al. 1992; VERMEULEN, JANSSENS DE BISTHOVEN et al. 1994; VERMEULEN 1995 etc.) - and we hope to be admitted to this society.

Our first steps were made traditionally: a study of deformities in natural water bodies

(NAZAROVA 1997, ZINTCHENKO, NAZAROVA 1997). As a next step the influence of Cu and polluted sediments on chironomid larvae in laboratory conditions were investigated (NAZAROVA, LATYPOVA, TUHVATULLINA 1999). Cu was chosen as it is one of the most important pollutants in all waterbodies in our region. Currently we are concentrating on analyses of teratogenic influences of some biochemical compounds (the newly created class of cholinesterase inhibitors) in combination with their harmful influence on karyotype of laboratory chironomids in chronic experiments over a few generations. Our experiments have revealed that in the second generation there appeared even more morphological abnormalities than in the first one, in spite of the fact that the second generation was growing up in clean water. Changes in normal chromosome structure were found in both generations. So at present we believe that cholinesterase inhibitors (compounds which are expected to be used in pharmacology and perhaps as a base for some pesticides) with their influence on a chironomid's metabolic processes can cause morphological and chromosomal abnormalities which persist into the next generation. Of course this study is just in its early stages, but we hope that its development will help us to understand better the diversity of relationships between living and nonliving nature and the role of anthropogenic factors in it.

Bibliography

- JANSSENS DE BISTHOVEN, L., TIMMERMANS, K.R., AND OLLEVIER F. (1992). The concentration of cadmium, lead, copper and zinc in *Chironomus* gr. *thummi* larvae (Diptera: Chironomidae) with deformed versus normal menta. *Hydrobiologia*..-**239**: 141-149.
- MADDEN, C.P., SUTER, P.J., NICHOLSON, B.C. and AUSTIN A.D (1992). Deformities in chironomid larvae as indicators of pollution (pesticide) stress.-*Neth. J. Aquat. Ecol.* **26**: 551-557.
- NAZAROVA L. (1997). Effect of main pollutants of oil-extracting region on incidence of

- mentum deformities in Chironomidae (Diptera) larvae. -In: HOFFRICHTER, O. (ed.) 13th International Symposium on Chironomidae, Freiburg im Breisgau, 5-9 September 1997, Abstracts.: p. 86.
- NAZAROVA, L.B., LATYOVA, V.Z., TUCHVATULLINA, L.G. (1999). Cu teratogenic influence on chironomid larvae. Toxicological review.- No. 3.(in russian).
- SKALSKAYA N.A. (1994). The reaction of Darwin reservoir lakes zooperiphiton on acidification// Structure and functioning of acid lakes. - ecosystem. Proceedings of IBVB, _.:Nauka.- v.70 (73).- pp.170-185.
- GUCHTE van DE, C. & VAN URK, G. (1989). Discrepancies in the effects of field and artificially heavy metal contaminated aquatic sediments upon midge larvae.- In: Heavy Metals in the Environment, 7th International Conference, Geneva.
- VERMEULEN, A.C. (1995). Elaborating chironomid deformities as bioindicators of toxic sediment stress: the potential application of mixture toxicity concepts.- Annls zool. fenn. 32: 265-285.
- VERMEULEN, A.C., JANSSENS DE BISTHOVEN, L., POSTMA, J. & OLLEVIER, F. (1994). Mechanical wear of *Chironomus riparius* (Diptera, Chironomidae) larvae reared in two artificial substrates. - First Benelux-congress of Zoology, 4-5 November, University of Leuven, Leuven, Belgium.
- WARWICK, W.F. (1985). Morphological abnormalities in Chironomidae (Diptera) larvae as measures of toxic stress in freshwater ecosystems: indexing antennal deformities in *Chironomus* Meigen.- Can. J. Fish. Aquat. Sci. 42: 1881-1914.
- WARWICK W.F. (1989). Morphological deformities in larvae of *Procladius* SKUSE (Diptera: Chironomidae) and their biomonitoring potential. - Can. J.Fish.Aquat.Sci. 46: 1255-1270.
- WARWICK, W.F. (1990). Morphological deformities in Chironomidae (Diptera) larvae from the Lac St. Louis and Laprairie Basins of the St. Lawrence River. - J.Great Lakes Res. 16: 185-208.
- WARWICK W.F. (1991). Indexing deformities in ligulae and antennae of *Procladius* larvae (Diptera: Chironomidae): application to contaminant-stressed environments.- Can.J.Fish.Aquat.Sci. 1991. 48: 1151-1166.
- ZINCHENKO, T.D., GOLOVATUK, L.V., MARCHENKO, N.A.. (1997). Ecological state of Chapaevka-river basin in condition of anthropogenic press (bioindication). Ecological safe and stable development of Samara region.- Togliatti. 3: 124-145. (in russian)
- ZINCHENKO, T.D., NAZAROVA, L.B. (1997). Use of the morphological deformities as biological screening tool for assessment of the ecological state of water reservoirs.- III. Regional conference „Actual ecological problems of RT“- Kazan: 95-96 (in russian).

POLYTENE CHROMOSOMES OF DIFFERENT SUBFAMILIES OF FAMILY CHIRONOMIDAE, DIPTERA.

By P. Michailova

Institute of Zoology, Bulgarian Academy of Sciences,
1, Tzar Osvoboditel boul., 1000 Sofia

The Chironomidae has still not been fully studied from the systematic aspect. The conventional morphometric method does not always provide an opportunity for determining separate forms which should, on the basis of a number of biological characteristics be considered "well differentiated species". The morphometric method can "work" for some species of chironomids only at one stage of ontogenesis, while for others it may not "work" at all. There are no clearly distinct differences in the larval stage of many related species and many of them are thus known merely as groups of species or larval forms. The polytene chromosomes appeared to be particularly promising for taxonomic diagnosis of chironomid species combined with the morphological characteristics of the separate developmental stages.

The diagnostic value of karyotype characteristics like the number and morphology of the polytene chromosomes, marker zones, number and localization of "weak points", Balbiani rings and nucleoli is great. The giant size and constant specific band structure of polytene chromosomes allows for its wide application as a stable diagnostic character in the systematics of the Chironomidae. Every band represents a complex of identical chromomeres adjacent to each other of solid spiralized DNP regions (BEERMANN 1952). The reasons of the pairing of homologues in polytene chromosomes is not known yet. It may be that the physical joining between sister chromosomes in homologous sections plays an important role. The protein interaction between sister chromomeres can not be excluded either (ANANIEV, BARSKY

1985). In most species of Chironomidae the homologues are paired, with some exception in genera: *Chironomus* (KEYL 1962) and *Glyptotendipes* (MICHAILOVA 1989a). An asynapsis in the polytene chromosomes reveals differences in the chromomere and chromonema organization of the homologues and their cyclic state (PROKOFIEVA-BELGOVSKAJA 1986). The asynapsis in polytene chromosomes can also be favoured by heterozygous inversions or gene differences in homologues. These peculiarities are easily seen in inter-species hybrids. Each chromosome in the karyotype of every species has its unique band pattern. Sometimes under the influence of various factors (e.g. infection, salinity, different environments) the banding pattern changes: becoming diffused or super contracted or shortened. The shortening of the chromosomes is caused by the fusion of groups of neighbouring bands into chromatin blocks. In these cases the inactivation of transcription processes leads to DNA condensation in earlier active regions accompanied by association of separate bands, resulting in shortened chromosomes. However, there is no alteration to the basic banding markers of the chromosome.

Sequences of band pattern can be changed through different chromosome rearrangements: inversions, translocations (homozygous reciprocal translocations and fusions). In most cases heterozygous inversions ensure plasticity of the species. When the aberration has a high selective value, it is further stabilized by selective forces and spread in the population, so becoming a constant characteristic of the species. Such chromosome differences are directly linked to isolating mechanisms (disturbances occurring in the meiosis of hybrids, unbalanced gametes formed). Among species of the successful genus *Chironomus* a great similarity of chromosome band patterns has been found. The genetic material has been redistributed through different chromosome aberrations. On the basis of homozygous reciprocal translocations species of this genus may have the chromosome arm combination: AB, CD, EF, G or AE, BF, CD, G etc.. Species having the same chromosome arm combination may be united into complexes (KEYL 1962). So, the karyotype based on polytene chromosomes plays an integrating role in the systematics of the Chironomidae. In these complexes species are differentiated on the basis of homozygous inversions (KEYL 1962). Homozygous inversions are responsible for divergence of sibling species and in the microevolution differentiation of the species

(MICHAILOVA 1989, a,b), (KIKNADZE et al. 1991). In the most primitive subfamilies (Diamesinae, Orthocladiinae) different types of rearrangements have been incorporated during the karyotype evolution, so that only a few sections of the polytene chromosomes of related species have common band patterns or distinguishable by homozygous inversions (MICHAILOVA 1989 a).

The special appearance of some polytene chromosome sections, the so called "weak points" can be used as a species characteristic. There are constructions of chromosomes which are underreplicated during polytenization. Also, there are differences in the band structure organisation of chromosomes of related species. These can be revealed by "C" and "Q" banding methods. Common banding patterns have been found in arms of different species belonging to other genera. These common banding patterns are considered as "basic" or "plesiomorphic", existing in a hypothetical stem species before separation (WÜLKER 1980). Such basic patterns have been established in genera *Glyptotendipes*, *Chironomus*, *Endochironomus*, *Micropsectra*, *Brillia*, *Orthocladius*, *Cricotopus*, *Prodiamesa* (MICHAILOVA 1989 a).

The number and position of Balbiani rings and nucleoli are a significant taxonomic character for the family. Within the larval salivary glands of chironomids a functional differentiation of distinct glandular regions has been established in some species (*Acricotopus lucidus*, *Axarus* sp.). This is reflected in the level of the polyteny of the chromosomes and in the expression of cell specific Balbiani rings. More primitive subfamilies (Orthocladiinae, Diamesinae) have great functional activity, realized by Balbiani rings. They can be localized in one and the same chromosome (Diamesinae) or distributed in different chromosomes (Orthocladiinae). In these subfamilies the position of nucleolus is not fixed. In the subfamily Chironominae very often (with few exceptions only) the nucleolus and a Balbiani ring are situated in the short chromosome.

An important taxonomic character is the manifestation of the centromere region in polytene chromosomes. In most species of the genera *Chironomus*, *Endochironomus*, *Micropsectra*, *Dicrotendipes*, *Acricotopus* the centromere region is represented by a dark band. In species of genera *Cricotopus*, *Orthocladius*, *Glyptotendipes* the centromere region is a large dark block. The chromosomes of some species of the genera *Cricotopus*,

Parachironomus, *Orthocladius* (MICHAILOVA 1989 a) and *Lipinella* (KERKIS et al. 1985) have been combined in a chromocentre. It has been formed as a result of a number of ectopic contacts arising between the separate replicates in the precentromere regions of the chromosomes. The inference to be drawn is that the evolution of chromosomes of these groups has proceeded much more rapidly than the evolution of external morphological characters. This phenomenon of a high karyotypic diversity among closely related species manifests the so called "chromosome tachytely" evolution (MARKS 1983).

References

- ANANIEV, E. V. & BARSKY, V. E. (1985). Electron microscope map of *Drosophila* polytene chromosomes: 1-95.
- BEERMANN, V. (1952). Chromosomenkonstanz und spezifische Modifikation der Chromosomenstruktur in der Entwicklung und Organdifferenzierung von *Chironomus tentans*. - *Chromosoma*, **52**: 139-198.
- KERKIS, I. E., KIKNADZE, I. I. & SHILOVA, A. I. (1985). Karyological characteristics of *Lipinella arenicola* SHILOVA (Diptera, Chironomidae). - *Tsitologia*, **27**: 1410-1413.
- KEYL, H. (1962). Chromosomenevolution bei *Chironomus*. II. Chromosomenumbauten und phyogenetische Beziehungen der Arten. - *Chromosoma*, **13**: 496-541.
- KIKNADZE, I., SHILOVA, A., KERKIS, I., SHOBANOV, N., ZELENZOV, N., GREBENJUK, L., ISTOMINA, A., PRESLOV, B., (1991). Karyotype and morphology of larvae of Chironomini. N. Novosibirsk, 115 p.
- MARKS, I. (1983). Rates of Karyotype evolution. - *Syst. Zool.* **32**, 2: 207-209.
- MICHAILOVA, P. (1989 a). The polytene chromosomes and their significance to the systematics of the family Chironomidae, Diptera. - *Acta Zool. Fennica*, **186**: 1-107.
- MICHAILOVA, P. (1989 b). Cytotaxonomic studies on the experimental hybrid of *Glyptotendipes barbipes* (STAEGER) and *Glyptotendipes salinus* MICHAILOVA (Diptera, Chironomidae). - *Biol. Debr. Oecol. Hung.*, **32**, 2: 141-150.
- PROKOFIEVA-BELGOVSKAJA, A. A. (1986). Heterochromatin regions of chromosomes. 431 p.
- WÜLKER, W. (1980). Basic patterns in chromosome evolution in the genus *Chironomus* (Diptera). - *Z-zool. Syst. Evolutionforsch.*, **18**, 2: 112-123.

CHIRONOMID MIDGE SWARMS ASSOCIATED WITH SLOW SAND FILTRATION WORKS

By P. D. Armitage, J. H. B. Blackburn and I. T. Harris

CEH Dorset (Formerly the Institutes of Terrestrial Ecology and Freshwater Ecology). Winfrith Technology Centre, Corchester, Dorset DT2 8ZD, UK

The mass emergence of chironomid midges frequently causes problems to people living next to freshwater habitats. A local water company, Bournemouth and West Hampshire Water (BWHW), operate slow sand filters for drinking water treatment beside the River Abon at Mill Road, Christchurch and at Francis Avenue, Bournemouth. In recent years, there have been increased incidences of swarms of midges occurring at such densities as to cause problems to local residents, particularly those occupying new housing developments near the slow sand filters.

The emergence of large numbers of midges results in swarms which are carried or are attracted to, nearby housing. It has not been possible to predict the occurrence of these swarms due to a lack of basic ecological data linked to operational procedures and timing. The project is funded by BWHW and is attempting to determine the chironomid faunal

composition and the identity of nuisance species at the two locations, initially over a period of one year. We are also investigating the life-history of the dominant species and trapping adults to determine emergence time and the distribution of adults around the sites. These data will be linked to information on filter bed operation and local weather conditions.

In general terms, the filter beds represent temporary water bodies. The beds are vertical-sided rectangular ponds which have a base of porous concrete overlain with gravel and then by a thick layer of sand which is smoothed and levelled to produce a flat homogeneous bottom. The depth of water is also uniform at about 1 m, depending on operational practice. The continuous supply of water carrying particulate matter, which is trapped at the sand surface, provides a rich source of food for organisms resulting in the build up of large populations of oligochaetes (worms),

nematodes, protozoans and chironomid midge larvae. Periodically the beds are emptied (about every 12 weeks at Christchurch and over 20 weeks at Bournemouth) and the filtering medium cleaned, thus only organisms which can complete their development within the period the beds are run can maintain populations. Chironomid midges are well adapted to survive in such environments.

Of the twenty-three taxa recorded from the filter beds only seven were common *Ablabesmyia monilis*, *Macropelopia nebulosa*, *Cricotopus (Isocladius) sp.? tricinatus*, *Psectrocladius barbimanus*, *Orthocladius glabripennis*, *Micropsectra lindrothi*, and *Tanytarsus gregarius/inaequalis*. Of these only the *Tanytarsini* and *Orthocladius glabripennis* occurred in large enough numbers to cause a nuisance.

The filter beds at both Christchurch and Bournemouth are characterised by their long periods of operation. At Bournemouth the mean bed run for the filters which were sampled, was approximately 150 days and at Christchurch the period was shorter (105 days - based on previous run times). These periods are up to three times longer than at the Thames Water Works at Ashford Common where the longest run observed during our 1990 study was 74 days and the mean 29 days (WOTTON et al 1992). The implications of this are that complex communities can develop in the beds with many overlapping generations. In the summer periods chironomid species can complete their development in as little as 17 days. In beds which are run for periods of less than 30 days only one generation of these species would be able to complete their life cycle. In addition, longer bed runs allow colonisation by a wide range of species and the Bournemouth and Christchurch beds supported a pond-like fauna. These characteristics have made it difficult to estimate the development time of individual taxa as one generation overlaps the other and there is continual recruitment from egg-laying females. At the

same time the relative complexity of the community has prevented dominance by one generation which can lead to mass emergence. However post-winter conditions (increase in light and temperature) result in maturation of the community as a whole and it is at these times that mass emergence, resulting in large swarms, has occurred in recent years. The perception of these swarms as nuisance is determined largely by the wind speed and direction which control movement off site towards residential areas.

This study will continue through the summer and end in October. In the meantime I would be interested to hear of any other similar investigations in other parts of the world. Slow sand filters provide an environment to examine both basic ecological aspects (competition, resource use, adaptation, distribution, behaviour of adults and larvae) and applied problems (filtration rates, water quality, nuisance problems etc.) and can provide a rich source of information on chironomid ecology and behaviour (WOTTON & ARMITAGE 1994).

References:

- WOTTON, R.S., ARMITAGE, P.D., ASTON, K., BLACKBURN, J.H., HAMBURGER, M. & WOODWAR, C.A. (1992). Colonization and emergence of midges (Chironomidae: Diptera) in slow sand filter beds. *Netherlands Journal of Aquatic Ecology*, **26**(2-4): 331-339.
- WOTTON R. S. & ARMITAGE P. D. (1994). Size of midges (Diptera: Chironomidae) emerging from newly-filled ponds. In: *Chironomids from genes to ecosystems* (ed. P. S. CRANSTON): 355-361, CSIRO Publications.

Patrick Armitage

CEH Dorset (Formerly the Institutes of Terrestrial Ecology and Freshwater Ecology) Winfrith Technology Centre
Dorchester Dorset DT2 8ZD UK e-mail:pdar@ceh.ac.uk www.ceh.ac.uk

NEWS FROM THE RUSSIAN FAR EAST


By **Eugenyi A. Makarchenko**

Institute of Biology and Soil Sciences, Far East Branch, Russian Academy of Sciences,
(FEB RAS) 690022, Vladivostok, Russia

Last year I introduced my working group, which consists of **Marina A. Makarchenko** (Orthocladiinae), **Oksana V. Zorina** (Chironominae) and **Eugenyi A. Makarchenko** (Podonominae, Diamesinae, Prodiamesinae and Orthocladiinae). The main purpose of our investigation is to prepare a Key for Chironomidae (males, pupae and larvae) of the Russian Far East.

During the second part of 1999 and in 2000 we have studied the taxonomy of the Orthocladiinae from Vrangal Island, Chukchi Peninsula and Primorye (territory from Vladivostok to Khabarovsk), and Chironominae (tribe Chironomini) from the south part of the Russian Far East (Primorye Territory, Khabarovsk Region, Sakhalin and Kurile Islands) (Fig. 1).

Fig.1. The map of the Russian Far East



From Vrangal Island we identified for the first time 27 species of Orthocladiinae: *Bryophaenocladus nitidicollis* (GOETGH.), *Chaetocladus festivus* (HOLMGR.), *Chaetocladus* sp. n. 1, *Chaetocladus* sp. n. 2, *Chaetocladus* sp. n. 3, *Corynoneura arctica* KIEFF., *Cricotopus* (*C.*) *tibialis* (MG.), *Hydrobaenus fusistylus* (GOETGH.), *Limnophyes brachytomus* (KIEFF.), *L. pumilio* (HOLMGR.), *Limnophyes* sp. n. 1, *Limnophyes* sp. n. 2, *Metriocnemus eurynotus* (HOLMGR.), *M. intergerivus* SÆTHER, *M. ursinus* (HOLMGR.), *Orthocladus* (*Eudactylocladius*)

gelidus KIEFF., *O.* (s.str.) *hazenensis* SOPONIS, *O.* (*Pogonocladus*) *consobrinus* (HOLMGR.), *Pseudosmittia recta* (ED W.), *P. nanseni* (KIEFF.), *Rheocricotopus* (s.str.) *reduncus* SÆTHER et SCHNELL, *Smittia extrema* (HOLMGR.), *S. joganbrevicosta* SASA et OKAZAWA, *Tokunagaia kibunensis* (TOK.), *T. rectangularis* (GOETGH.).

Sixty five species of Orthocladiinae genera *Antillocladius*, *Bryophaenocladus*, *Camptocladus*, *Corynoneura*, *Cricotopus* (s.str.), *C.* (*Pseudocricotopus*), *C.* (*Nostococladus*), *C.* (*Isocladus*), *Eukiefferiella*, *Gymnometriocnemus*, *Heterotrissocladus*, *Hydrobaenus*, *Krenosmittia*, *Limnophyes*, *Metriocnemus*, *Mesosmittia*, *Nanocladus*, *Orthocladus* (s.str.), *O.* (*Euorthocladus*), *Pseudorthocladus*, *Parametriocnemus*, *Paratrachocladus*, *Pseudosmittia*, *Smittia*, *Stilocladus*, *Thienemanniella*, *Tsudoyusurika*, *Tvetenia* from Primorye Territory were found, from which seven species of *Bryophaenocladus*, *C.* (*Pseudocricotopus*), *Limnophyes*, *Pseudorthocladus*, *Smittia*, *Stilocladus* and *Tsudoyusurika*, appear to be new for science. One species, *Hydrobaenus calvescens* SÆTHER, is recorded for the Palaearctic region for the first time.

One hundred ninety two species of Chironomini from the south part of the Russian Far East were identified, of which about 20 species of the genera *Cryptochironomus*, *Cryptotendipes*, *Dicrotendipes*, *Microtendipes*, *Paratendipes*, *Polypedilum*, *Stenochironomus* and *Stictochironomus* appear to be new to science; 9 species, *Harnischia incidata* TOWNES, *Kloosia dorsenna* (SÆTHER), *Phaenopsectra* ? *profusa* (TOWNES), *P.* ? *dyari* (TOWNES), *Polypedilum exilicaudatus* SÆTHER et SUNDAL, *P. albinodus* TOWNES, *Stictochironomus* ? *lutosus* (TOWNES), *S.* ? *naevus* (MITCHELL), are recorded for Palaearctic region for the first time. Twenty species were known previously only from Japan.

More detailed information can be found on the Russian Chironomid Home Page, which was opened in Vladivostok in March 14, 2000. Address of this site is:

<http://www.tendipes.febras.ru>

CHIRONOMID FAUNA OF THE RIVER ANGARA

By Erbaeva E. A.

Scientific-Research Institute of Biology at Irkutsk State University, Irkutsk, Russia

Introduction

The Angara is the main river of Eastern Siberia. It flows out of the oligotrophic Lake Baikal and after 1779 km joins the river Yenisey. The water of the Angara remains baikalian for a significant distance. Even at the mouth of the river 45% of the total water is baikalian. According to R. A. GOLYSHKINA (1970) the substratum is composed mainly of stones, due to relief and geological structure. The Angara previously was a mountain river with a high current velocity of 7 km h⁻¹ in depositing areas and 12 - 15 km h⁻¹ in eroding areas.

Different authors have studied the chironomid fauna of the Angara. 144 species and forms of Chironomids (Tanypodinae - 10, Diamesinae - 10, Prodiamesinae - 2, Orthocladiinae - 63, Chironominae - 59) have been found in it. 5 species (*Diamesa baicalensis* CHERN., "*Orthocladius compactus* LINEVICH", *O. gregarius* LINEVICH, *O. setosus* LINEVICH, *Neozavrelia minuta* LINEVICH) are endemics of Baikal; *Cricotopus angarensis* LINEVICH is an endemic of the Angara.

Chironomid fauna before the dams

A. A. LINEVICH (1953, 1957, 1981) studied the chironomid fauna over the region from its outflow from Lake Baikal to the city of Bratsk (670 km) from material collected during the 1940-s. Later, in the middle of 1960-s N. V. VERSHININ (1967) studied the chironomids from the city of Svirsk to Bratsk (470 km). The chironomid fauna of the lower part of the Angara from the settlement Motygin to its mouth (123 km) was studied by I. I. GREZE (1953) in the 1950- s.

The Angara from the Lake Baikal outflow to Bratsk is divisible into three parts: 1) from outflow to Irkutsk, 2) from Irkutsk to the inflow of the river Kitoy, 3) from the mouth of Kitoy to Bratsk.

According to A. A. LINEVICH (1953, 1957, 1981) over the first part representatives of the subfamilies Diamesinae and Orthocladiinae dominated (listed in order of abundance): *Diamesa baicalensis* CHERN., "*Parorthocladius tridentifer* LINEVICH", "*Orthocladius compactus* LINEVICH", *O. frigidus* (ZETT.), *O. trigonolabis* ED W., *Eukiefferiella coerulea* KIEFF., *E. clypeata* KIEFF., "*E. longicalcar* KIEFF.", *E. similis*

GOETGH., *Diplocladius cultriger* KIEFF., *Pagastia lanceolata* (TOK.), *Potthastia longimana* (KIEFF.), *Pseudodiamesa nivos* (GOETGH.), *Lauterbornia* sp., *Polypedilum* sp. (*Chironomina genuina* N3 LIPINA), "*Stictochironomus psammophilus* CHERN.", i. e. mainly lithorheophilic species.

The second part was dominated by the following species: *D. baicalensis*, *O. trigonolabis*, *O. frigidus*, *O. compactus*, *Polypedilum convictum* (WALK.), *Pagastia lanceolata*, *Potthastia gaedii* (MEIG.), *Pseudodiamesa nivos*, *E. coerulea*, *O. consobrinus* (HOLMG.), *C. angarensis*, *Eukiefferiella* sp., *Pagastia orientalis* (CHERN.). So, over the second part most species of the genus *Eukiefferiella* are absent, but lithorheophilic species (subfamilies Diamesinae and Orthocladiinae) continue to dominate.

Over the third part the following species are mentioned: *Pagastia lanceolata*, *Potthastia gaedii*, *Eukiefferiella coerulea*, *O. trigonolabis*, *O. compactus*, *O. frigidus*, *Polypedilum convictum*, *Pseudodiamesa nivos*, *Monodiamesa bathyphila* (KIEFF.), *Stictochironomus psammophilus*, *Polypedilum bicrenatum* KIEFF., *O. consobrinus*, *Cryptochironomus* gr. *defectus* (KIEFF.) *Harnischia curtilamellata* (MALLOCH), *Paracladopelma camptolabis* (KIEFF.), *Paratendipes "connectens"* N3 LIPINA, *Tanytarsus exiguus* JON., *Parorthocladius nudipennis* (KIEFF.), *D. baicalensis* and *Brillia bifida* (KIEFF.) In terms of numbers *P. lanceolata* and *P. gaedii* are first; in terms of the number of species, the subfamily Chironominae dominates.

Unfortunately, I.I. GREZE (1953) does not give a full list of the species of the lower part of the river, but only the dominant ones. The dominant species inhabiting vegetation are the larvae of *Tanytarsus exiguus*, *Polypedilum nubeculosum* MEIG., and *O. thienemanni* KIEFF. Vegetation on pebbles is inhabited by *T. exiguus*, *Micropendipes pedellus*, *Polypedilum scalaenum* (SCHRANK) and *Glyptotendipes gripenkoveni* KIEFF.. Sand is dominated by *Chernovskia orbiculus* TOWNES (*Chernovskia ra* ULOMSKY), *Robackia demeijerei* (KRUS.) and *Beckidia zabolotzkyi* (GOETGH.), species not found in the upper parts of the river. There are pelorheophilic and psammorheophilic forms dwelling in silted

sand: the most common among them are *Polypedilum bicrenatum* KIEFF., *P. scalaenum* (SCHRANK), *Cladotanytarsus* gr. *mancus* (WALK.), *Chironomus* gr. *thummi* (KIEFF.), and *Ch. pr. plumosus-reductus* LIPINA. Silt is inhabited by *Chironomus* gr. *thummi*. General features of the chironomid fauna of the lower part seem to be quite different from those of the upper parts. A rheophilic complex, including baikalian species and mainly composed of species belonging to the Diamesinae and Orthocladiinae, inhabits the typical stony biotopes of the upper parts of the river; this is replaced in the lower parts by a common complex of river forms belonging mainly to the Chironominae. In sandy biotopes the psammorheophils *C. ra*, *R. demejerei*, and *Bekidia zabolotzkyi* are found.

Present state of the chironomid fauna after erecting the complex of dams

The river Angara is unaffected initially (about 7 km), from Irkutsk to settlement Telma (about 80 km) and below the dam of the Ust-Ilim hydropower station to its mouth (962 km). Close to its outflow from Lake Baikal the same species that dwelled here before the regulation remain; the 5 baikalian endemic species mentioned above are still to be found. Lithorheophilic forms found in the 40-s dominate. After the regulation in the middle 70-s we have investigated the river near Angarsk (at 10 km). Comparison of the chironomids before and after the regulation has shown some changes in composition (TOMILOV & al. 1977). *O. frigidus*, *P. lanceolata*, *P. gaedi*, *O. consobrinus* and *S. psammophilus* which were here before regulation are no longer to be found. *O. gr. olivaceus* and *P. tridentifer* dominate and *D. baicalensis* occurs in stony substrata. In the bottom vegetation *O. saxicola* and *C. angarensis* prevail. *Polypedilum* sp. (Chironominae sp. N3 LIPINA) and *Monodiamesa bathyphila* (KIEFF.) dominate in silted sand. During 1973-1975 the chironomid fauna near the future Ust-Ilim reservoir over the 302 km from Bratsk to Ust-Ilimsk was investigated. 91 species were found, 42 species belonging to the subfamily Chironominae: *D. baicalensis* and *O. olivaceus* dominate in stony sediments; *Prodiamesa olivacea* and *Chironomus obtusidens* on small pebbles and silted sand; *Cricotopus sylvestris*, *C. biformis*, *O. frigidus* and *Diplocladius cultriger* on stones covered with algae; *Chironomus cingulatus* MEIG., *Paratendipes albimanus* (MEIG.), "*Paratrichocladius inaequalis*

KIEFF." and *Tanytarsus* gr. *gregarius* KIEFF. in silted sediments.

At the end of the 80-s to the beginning of the 90-s we studied the chironomids of the Angara below the dam of the Ust-Ilim power station (KOZHOVA & al. 1993) from the city of Ust-Ilimsk to the mouth of the river Kata (about 90 km). Here there are lot of eroding currents with high velocity: 51 species of Chironomidae were found, about 50 % representatives of the subfamily Orthocladiinae. *D. baicalensis*, *P. lanceolata* and *Pagastia orientalis* dominate on stony sediments; *Cricotopus sylvestris*, *C. biformis* and *O. saxicola* dominate on stones covered with *Ulothrix*; *P. nivosa*, *P. olivacea*, *O. olivaceus*, *Diamesa insignipes* KIEFF., *Eukiefferiella coeruleascens*, *P. inaequalis* and *Micropsectra junci* (MEIG.) in silted sand and pebbles; *Paratendipes albimanus* in silt. *O. frigidus*, *O. consobrinus*, *O. compactus*, *O. gregarius*, *C. angarensis*, *P. inaequalis*, *Cr. gr. defectus*, *Cladopelma viridula* (L.), *Parachironomus pararostratus*, *Polypedilum bicrenatum*, *E. albipennis*, *Microtendipes pedellus* (DE GEER) and *Cladotanytarsus* gr. *mancus* (WALK.) found in upper parts are not found here.

Conclusion

In the river Angara from its outflow from Lake Baikal to its junction with the River Yenisey hydrologic conditions change: decrease of current velocity, increase of water temperature, decrease of transparency, and increase in deposition of sand and silt, i.e. the transformation of a mountain river into the usual Siberian river, and consequent changes in the chironomid fauna take place. In the upper parts of the river lithorheophils of the subfamilies Diamesinae and Orthocladiinae prevail, whereas in the lower part, pelorheophils of the subfamily Chironominae dominate. The main characteristic feature of the river Angara - the influence of Lake Baikal on the fauna (the presence of baikalian endemics) - occurs mainly in the upper part of the river.

References:

- GOLYSHKINA, R.A. 1970. Zoobenthos of the river Angara. Ph. D. Thesis. Irkutsk. 35 pp. (In Russian).
- GREZE I. I. 1953. Hydrobiology of the lower parts of the river Angara. Proc. of the Aquatic Ecology Society of the USSR., 5: 203-211. (In Russian)

- KOZHOVA, O. M., ERBAEVA, E. A., IZMEST'EVA, L.R. & al. 1993. The distribution of hydrobionts in the river Angara in the region of waste waters of Ust-Ilmsk timber processing plant inflow. *VINITI*, N 1972-B93. 57 pp. (In Russian)
- LINEVICH, A.A. 1953. Tendipedids of the upper part of the river Angara. *Proc. of Irkutsk State University*, 7 (1-2): 153-175. (In Russian)
- LINEVICH, A.A. 1957. Concerning the larvae of tendipedids of the upper part of the river Angara. *Proc. of Biologic-Geographic Scientific-Research Institute*, 17 (1-4): 144-154. (In Russian)
- LINEVICH, A.A. 1981. Chironomidae of Baikal and Pribaikalye. *Novosibirsk, Nauka*. 152 pp. (In Russian)
- TOMILOV, A.A., ERBAEVA, E.A., MEKHANIKOVA, I.V. & al. 1977. Macrozoobenthos of the river Angara near the city of Angarsk. In: *Biological investigations of water bodies of Eastern Siberia*. Irkutsk, pp. 44-70. (In Russian)
- VERSHININ, N.V. 1967. The feeding resources of fishes of the middle Angara river before the regulation (bottom fauna). In: *Fishes and feeding resources of basins of rivers and reservoirs of Eastern Siberia*. Krasnoyarsk, pp. 261-290. (In Russian)

CHIRONOMID TYPES AT MUSÉUM NATIONAL D'HISTOIRE NATURELLE, PARIS (Meigen collection not included)

By Torbjørn Ekrem

Museum of Zoology, University of Bergen, Muséplass 3, N-5007 Bergen, Norway
(email: Torbjorn.Ekremoo.uib.no)

The chironomid type collection in Paris is, except for the Meigen collection, not very well known. The list presented here was generated during my visit to the Muséum National d'Histoire Naturelle as part of the European Union's PARASYST program in June/July 2000. Notes on the current taxonomic status of the species are included. These notes are, when not available on the labels, collected from the catalogues of Neotropical, Palaearctic, Oriental and Afrotropical Chironomidae (SUBLETTE & SUBLETTE 1973; FREEMAN & CRANSTON 1980; ASHE & CRANSTON 1990; SPIES & REISS 1996). A list of the chironomid specimens in the Meigen collection, which is currently under revision by Martin Spies (pers. comm.), has previously been published by FITTKAU and REISS (1976).

The specimens of Goetghebuer and Kieffer all bear identical "cotype" labels. These are probably not original, and have been placed on their respective pins by another scientist. Freeman and Macquart types are located in separate boxes. The other type specimens form part of the general chironomid collection.

Acknowledgements

I would like to thank Dr. Cristophe Daugeron at Muséum National d'Histoire Naturelle for his hospitality during my stay and Prof. Daniel F. Goujet and Dr. François Bouvier for answering my many questions and organizing my visit. Thanks also to Martin Spies for all

helpful information and to Trond Andersen for comments on the manuscript.

References:

- ASHE, P. & CRANSTON, P.S. (1990). Family Chironomidae. pp 113-355 in SOÓS, A. & PAPP, L. (eds): *Catalogue of Palaearctic Diptera*. Volume 2. Psychodidae - Chironomidae. Budapest, Akadémiai Kiadó.
- BLANCHARD, E. (1852). *Orden IX. Dípteros*. pp 1-471 in Gay, C. (ed): *Historia física y política de Chile*. Zoología. Paris & Santiago de Chile.
- FITTKAU, E.J. & REISS, F. (1976). *Die Chironomidentypen und ihr Erhaltungszustand in der Sammlung des Muséum d'Histoire Naturelle, Paris*. *Chironomus*, 1: 146-150.
- FREEMAN, P. (1956). Some Chironomidae (Diptera) from French West- Africa. *Bull. Inst. fr. Afr. noire*, 18: 93-96.
- FREEMAN, P. (1961). A collection of Chironomidae and Culicidae subfamily Dixinae (Diptera, Nematocera) from Madagascar. *Mém. Inst. scient. Madagascar, Sér. E*, 12: 238-255.
- FREEMAN, P. & CRANSTON, P.S. (1980). II. Family Chironomidae. Pp 175-202 in CROSSKEY, R. W. (ed): *Catalogue of the Diptera of the Afrotropical Region*. London, British Museum (Natural History).
- GOETGHEBUER, M. (1919). Observations sur les larves et les nymphes de quelques Chironomides de Belgique. *Annls Biol. lacustre*, 9: 51-78.
- KIEFFER, J.J. (1912). H. Sauter's Formosa - Ausbeute. *Suppl. Ent.*, 1: 32-43.

- KIEFFER, J.J. (1922). Chironomides nouveaux ou peu connus de la région paléarctique. *Annls Soc. scient. Brux.*, **42**: 71-180.
- MACQUART, J. (1826). Insectes Diptères du Nord de la France. I. Tipulaires. *Mém. Soc. Sci. Agric. Lille*, **1823-1824**: 59-224.
- MACQUART, J. (1838). Diptères exotiques nouveaux ou peu connus. *Mém. Soc.Sci. Agric. Lille*, **2**: 9-225.
- ROSSARO, B. & DELETTRE, Y.R. (1992). Description of *Smittia celtica*, n. sp. (Diptera, Chironomidae). *Annls Soc. ent. Fr.*, **28**: 365-370.
- SÉGUY, E. (1965). Deux nouveaux Tendipédides des Iles Crozet (Insectes Diptères Nématocères. *Bull. Mus. natn. Hist. nat., Paris*, **37**: 285-289.
- SERRA-TOSIO, B. (1982). Description du male de *Belgica albipes* (SÉGUY, 1965), n. comb., rare chironomidé microptère des îles Crozet (Diptera). *Revue fr. Ent.*, **4**: 97-100.
- SERRA-TOSIO, B. (1986). Un nouveau chironomide antarctique des îles Crozet, *Parochlus crozetensis* n. sp. (Diptera, Nematocera). *Nouv. Revue Ent. (N.S.)*, **3**: 149-159.
- SPIES, M. & REISS, F. (1996). Catalog and bibliography of Neotropical and Mexican Chironomidae (Insecta, Diptera). *Spixiana Suppl.*, **22**: 61-119.
- SUBLETTE, J.E. & SUBLETTE, M.F. (1973). Family Chironomidae. Pp 389-422 in Delfinado, M. D. & Hardy, D. E. (eds): A catalog of the Diptera of the Oriental region. Honolulu, The University Press of Hawaii.

Table 1. Types in the general chironomid collection in Muséum National d'Histoire Naturelle, Paris

Number	Species	Author	Type status	Synonyms/present genus	Locality
21	"nitidus"	"MACQ" *	"Type", good*	" <i>Paratendipes albimanus</i> , GTGH."	Not given
3683	<i>Belgica albipes</i>	(SÉGUY, 1965)	Allotype male det. Serra-Tosio (1982) on slide		Crozet: Baie du Navirre Rochers, 25.III.1972, Ph. Dreux
	<i>Chaetocladius angustus</i>	FREEMAN, 1961	Holotype, male, good	<i>Bryophaenocladius angustus</i> FREEMAN, 1961	Madagascar Centre, Pic Boby 2400m, Andringitra, Ambalavao, 11-14.I.1958, B. Stuckenberg
	<i>Chaetocladius angustus</i>	FREEMAN, 1961	Paratype, male, good		As holotype
	<i>Chironomus bequaerti</i>	GOETGHEBUER, 1921	Cotype, male, good	<i>Einfeldia longipes</i> STAEGER, 1839	Destelbergen, 11.V.1915, M. Goetghebuer
	<i>Chironomus brayi</i>	GOETGHEBUER, 1921	Cotype, female, good	<i>Nilothauma brayi</i>	Virton, 2.IX.1920, M. Goetghebuer
	<i>Chironomus (Cryptochironomus) nigrofascia</i>	FREEMAN, 1961	Holotype, male, good	<i>Parachironomus nigrofasciatus</i> FREEMAN, 1961	Madagascar Centre, Pic Boby 2400m, Andringitra, Ambalavao, 11-14.I.1958, B. Stuckenberg
	<i>Chironomus (Endochironomus) pruinosa</i>	FREEMAN, 1961	Holotype, male, good	<i>Endochironomus pruinosa</i> FREEMAN, 1961	Madagascar Nord, Montagne Ambre 1000m, det. Diego Suarez, 23.XI-4.XII.1958, B. Stuckenberg
	<i>Chironomus latidens</i>	GOETGHEBUER, 1921	Cotype, male, good	<i>Einfeldia pagana</i> MEIGEN, 1838	Destelbergen, 1.VI.1920, M. Goetghebuer
	<i>Chironomus latidens</i>	GOETGHEBUER, 1921	5 specimens as cotype but without cotype label		

	<i>Chironomus mucronatus</i>	GOETGHEBUER, 1919	Cotype, male, good	<i>Parachironomus parilis</i> WALKER, 1856	Destelbergen, 11.V.1915, M. Goetghebuer
	<i>Chironomus obtusidens</i>	GOETGHEBUER, 1921	Cotype, male, good		Destelbergen, 23.IX.1916, M. Goetghebuer
	<i>Chironomus varus</i>	GOETGHEBUER, 1921	Cotype, male, good	<i>Parachironomus varus</i>	Destelbergen, 18.V.1920, M. Goetghebuer
	<i>Cricotopus carnosus</i>	KIEFFER, 1912	Cotype, male, fair	<i>Cricotopus (Isocladius) ornatus</i> (MEIG., 1818)	Tainan, Formosa, XI.1909, H. Sauter
	<i>Cricotopus carnosus</i>	KIEFFER, 1912	1 male and 2 females in same series but without cotype label		Tainan, Formosa, XI.1909, H.Sauter
	<i>Glyptotendipes sigillatus</i>	KIEFFER, 1922	Lectotype, female, design. Contreras-Lichtenberg, good		Aisne, Saint Gobert, July 1912, J. Surcouf
	<i>Glyptotendipes sigillatus</i>	KIEFFER, 1922	1 male Cotes du Nord, Lamballe, VIII.1910, J. Surcouf		
	<i>Metriocnemus cunepennis</i>	FREEMAN, 1961	Holotype, male, good	<i>Paraphaenocladius cuneipennis</i> Freeman 1961	Madagascar Centre, Moramanga 1000m, 18.-24.XI.1957, B. Stuckenberg
	<i>Metriocnemus cunepennis</i>	FREEMAN, 1961	Paratype, male, poor		As holotype
	<i>Microtendipes bicoloripennis</i>	FREEMAN, 1961	Holotype, female, good		Madagascar, Sambirano, Lokobe Nossibé 6m, 9.-23.XI.1957, B. Stuckenberg
	<i>Orthocladius stuckenbergi</i>	FREEMAN, 1961	Holotype, male, good		Madagascar Centre, Moramanga, Route d'Anosibe, 18.-24.XII.1957, B. Stuckenberg
	<i>Orthocladus stuckenbergi</i>	FREEMAN, 1961	Paratype, male, good		As holotype
3681	<i>Parochlus crozetensis</i>	SERRA-TOSIO, 1986	Holotype, male on slide		Iles Crozet: Pointe Basse, 12.XI.1979, D, Vernon
3682	<i>Parochlus crozetensis</i>	SERRA-TOSIO, 1986	Paratype male on slide		As holotype
	<i>Polypedilum airense</i>	FREEMAN, 1956	Holotype, male, fair	<i>Polypedilum aegyptium</i> KIEFFER, 1925	Irabellaben, Mts Baguezans, 1200-1300m, 26.-31.VIII.1947, L. Chopard, A. Villiers
	<i>Polypedilum bicoloratum</i>	FREEMAN, 1961	Holotype, male, good		Madagascar Nord, Montagne Ambre 1000m, deeeet. Diego Suarez, 23.XI-4.XII.1958, B. Stuckenberg
	<i>Polypedilum pallidiventris</i>	FREEMAN, 1961	Holotype, male, good	<i>Polypedilum pallidiventre</i> FREEMAN, 1961	Madagascar Centre, Vakoana 1520m, Andringitra

	<i>Polypedilum pelostolum</i>	KIEFFER, 1912	Cotype, female, fair	<i>Polypedilum nubifer</i> SKUSE, 1889	Ambalavao, 21.-24.I.1958, B. Stuckenberg Tainan, Formosa, X.1908, H. Sauter
	<i>Polypedilum pelostolum</i>	KIEFFER, 1912	2 males and 2 females in same series but without cotype label		
	<i>Polypedilum stuckenbergi</i>	FREEMAN, 1961	Holotype, male, good		Madagascar Centre, Pic Boby 2400m, Andringitra, Ambalavao, 11-14.I.1958, B. Stuckenberg As holotype
	<i>Polypedilum stuckenbergi</i>	FREEMAN, 1961	Paratype, male, good		
	<i>Procladius formosanus</i>	KIEFFER, 1912	Cotype, male, fair	<i>Tanypus formosanus</i>	Tainan, Formosa, II.1909, H. Sauter
3741	<i>Smittia celtica</i>	ROSSARO & DELETTRE, 1992	Paratype male on slide		Paimpont, Brittany, NL 13.XII.1988
3740	<i>Smittia celtica</i>	ROSSARO & DELETTRE, 1992	Paratype male on slide		Paimpont, Brittany, PA4 13.XII.1988
3742	<i>Smittia celtica</i>	ROSSARO & DELETTRE, 1992	Paratype larva on slide		Paimpont, Brittany, NPA 13.XII.1988
3739	<i>Smittia celtica</i>	ROSSARO & DELETTRE, 1992	Holotype male on slide		Paimpont, Brittany, NL 13.XII.1988
	<i>Stenochironomus unicalar</i>	FREEMAN, 1961	Holotype, female, good		Madagascar Nord, Montagne Ambre 1000m, det. Diego Suarez, 23.XI-4.XII.1958, B. Stuckenberg
	<i>Tanytarsus formosanus</i>	KIEFFER, 1912	Cotype, female, fair		Tainan, Formosa, II.1909, H. Sauter
	<i>Tanytarsus formosanus</i>	KIEFFER, 1912	2 males and 2 females in same series but without cotype label		Tainan, Formosa, II.1909 and X.1908, H. Sauter
	<i>Tendipes tainanus</i>	KIEFFER, 1912	Cotype, male, good	<i>Nilodorum tainanus</i>	Tainan, Formosa, II.1909, H. Sauter
	<i>Tendipes tainanus</i>	KIEFFER, 1912	1 male and 2 females in same series but without cotype label		Tainan, Formosa, XI.1909 and II.1909, H. Sauter

Table 2. Chironomid types in the Macquart collection

Number	Species	Author	Type status	Synonyms/present genus	Locality
1229	<i>Chironomus maculatus</i>	MACQUART, 1826	1 specimen missing hypopygium	<i>Chironomus maculosus</i> Macq. 1834 nom. dub.	North France
1230	<i>Chironomus trimaculatus</i>	MACQUART, 1838	1 specimen missing hypopygium	nom. dub.	Brazil

1231	<i>Chironomus ferrugineus</i>	MACQUART, 1838	1 female	nom. dub.	Brazil
1232	<i>Chironomus maculipennis</i>	BLANCHARD, 1852	3 males	<i>Chironomus maculosipennis</i> Kieffer 1906 replacement name. Unplaced in Chironiminae	Chile
1233	<i>Chironomus pallidulus</i>	BLANCHARD, 1852	3 males	Jun. hom. of <i>C. pallidulus</i> Meigen. Unplaced in Chironominae	Chile
1234	<i>Chironomus obscurellus</i>	BLANCHARD, 1852	1 specimen missing hypopygium	Unplaced in Chironominae	Chile
1235	<i>Chironomus tessellatus</i>	BLANCHARD, 1852	1 male	Unplaced in Chironominae	Chile
1236	<i>Chironomus articuliferus</i>	BLANCHARD, 1852	3 males	Unplaced in Chironomidae	Chile

SHORT COMMUNICATIONS

WHAT'S THE POINT

By Peter H. Langton

.....or, for that matter, the spine, spinule, tooth, tubercle or granule? These are terms I coined (LANGTON 1984, 1991) for different kinds of projections on the pupal cuticle of Chironomidae; to me a spinule is a very different structure to a point. Basically a point is the sharp end of a cuticular ridge, which when flattened vertically is v-shaped, resembling the point of a nail, whereas a spine is an elongate, more or less cylindrical structure, resembling the spine of a hedgehog (*Erinaceus*); a spinule is a small spine. Towards the posterior margin of the tergites in *Kloosia pusilla* there is a sudden gradation from the general covering of points through spinous points (where the cuticular ridge has a projecting spinous tip) to spines: usually there is no difficulty in ascribing a structure to one or the other category. The importance of distinguishing these in descriptions can be appreciated when comparing the exuviae of *Micropsectra atrofasciata*, *M. bidentata* and *M. lindrothi*; on tergite IV the lateral longitudinal bands on *atrofasciata* are of spines, on *bidentata* of spinules and on *lindrothi* of points. Not to make the distinction between spinule and point is to lose a useful descriptor. These terms are necessarily relative: a spine on a *Virgatanytarsus* would be no more than a spinule when superimposed on a *Dicrotendipes*. However, this is not a problem as the different sized exuviae are observed under different magnifications.

The tooth is a large conical or flattened more or less v-shaped projection, as in the posterior transverse tergite bands on *Diamesa* and *Cryptochironomus*. Tubercles and granules are blunt protrusions, the tubercle taller than broad at base, the granule broader than high. Shagreen is composed of minute spinules or points, usually arranged in short transverse rows which frequently form a reticulate pattern.

References:

- LANGTON, P.H. 1984. A key to pupal exuviae of British Chironomidae. 334 pp. Privately published.
 LANGTON, P.H. 1991. A key to pupal exuviae of West Palaearctic Chironomidae. 386 pp. Privately published.

KARYOSYSTEMATICS OF SPECIES OF PLUMOSUS GROUP (CHIRONOMIDAE, DIPTERA) FROM SOME MACEDONIAN LAKES

By P. Michailova¹ and Smilkov, S.²

¹Institute of Zoology, Bulgarian Academy of Sciences, Sofia 1000.

²Prirodno-Matemat.Fac.Biocoski Institut, Skopje, Macedonia.

The polytene karyotypes of two chironomid species from Ochrid and Doiran lakes are being studied. Both species (*Chironomus* sp.1 from Ochrid and *Chironomus* sp. 2 from Doiran) belong to the *thummi* complex, with chromosome set 8 and arm combination AB,CD,EF and G. The chromosomes AB,CD are metacentric, EF submetacentric and G telocentric. They have well expressed centromeric heterochromatin bands. A high frequency of heterozygous inversion in arm A of both species is established. Both species have a similar banding pattern to some species of the *plumosus* group. *Chironomus* sp. 1: arms A,B,E,F,G are identical with those of *C. plumosus*; arm C is identical with that of *C. entis*; arm D is similar to that of *C. entis* but a small group of bands are in an inverted position. *Chironomus* sp. 2: arm A is identical with that of *C. agilis*; arms B,C,D,E,F,G are as those of *C. plumosus*. However, arm B of this species has well formed BR, arm G has one BR only. Both species are closely related to those of the *plumosus* group. After detail analysis of all stages of metamorphosis we will consider the taxonomic position of these species.

PROFESSOR SÆTHER'S VISIT TO CHINA

Invited by Nankai University, Tianjin, China, and supported by the National Science Foundation of China (NSFC) and the Norwegian Research Council (NFR),

By Xinhua Wang

Professor Ole A. Sæther (Bergen, Norway) visited China from March 2 to 28 this year.

Due to his long-term support and great contributions to Chinese chironomid studies, on March 15, Prof. Sæther was awarded the title of honoured Professor of Nankai University. The president of Nankai Univ. bestowed the certificate in the ceremony.

During his month-long visit, Prof. Sæther delivered 6 guest lectures for the staff and students of Nankai University.

1. The canalized evolutionary potential.
2. The myth of objectivity and the electronic Ouija board, Different views on similarity.
3. Phylogeny of the Culicomorpha (Diptera).
4. Phylogeny of the subfamilies of Chironomidae (Diptera).
5. Zoogeographical patterns in Chironomidae (Diptera).
6. Morphological adaptations of tropical chironomids (Diptera).

At the same time, Prof. Sæther worked on a joint project initiated by Dr. Wang titled "Biosystematic studies on Chinese Chironomidae, with emphasis on immature stages" completing 3 joint papers:

1. The larvae of *Propiloscerus sinicus* WANG et SÆTHER and *P. paradoxus* (LUNDSTRÖM) (Diptera: Chironomidae).
2. First record of the genus *Paratrissocladius* ZAVREL from the Oriental region (Diptera: Chironomidae).
3. Two new species from China of the orientalis group of *Rheocricotopus* (*Psilocricotopus*) (Diptera: Chironomidae).

From March 18-25 Prof. Sæther together with Dr. Wang visited Shanghai and Hainan Island (Southern China) and had a field trip in Bawangling National Natural Conservation. A field trip to The New Territories in Hong Kong was arranged by Dr. David Dudgeon. Prof. Sæther left from Hong Kong on February 28.

THE FIRST MEETING AND SYMPOSIUM OF THE AQUATIC ENTOMOLOGIST'S SOCIETIES IN EAST ASIA (AESEA MEETING)

By Xinhua WANG

Dept. of Biology, Nankai University, China

The first meeting and symposium of the Aquatic Entomologist's Societies in East Asia (AESEA Meeting) was held in Chiaksan, Korea, from May 17-20, 2000. Sixty-three aquatic entomologists from Korea, China, Japan, Mongolia, Thailand Vietnam, Russia (Far East) and USA attended the meeting. The theme of the meeting was "The 21st Century and Aquatic Entomology in East Asia". More than 40 papers on systematics, ecology, behavior, zoogeography and environmental biology were given in oral presentations and posters. During the meeting a field trip was taken to Chiaksan National Park, about 100 Km southeast of Seoul.

According to the resolution of the meeting, AESEA meeting will convene every second year. The next meeting will take place in Japan in 2002 and the third in China in 2004.

The papers and abstracts on chironomid research presented at the meeting include:

CHON, T. S. & al.: Water quality monitoring by community dynamics of benthic macroinvertebrates in Youngjae stream, Han River, Korea.

KUO, C. H.: The succession of a chironomid community in an urban stream.

MAKARCHENKO, E. V. & al.: Review of the Chironomidae (Diptera) of the Russian Far East.

SERGEEVA, I. V. and Makarchenko, E. V.: Morphokaryological investigations of the subfamily Tanypodinae (Diptera, Chironomidae) of the Russian Far East.

WANG, X. H.: Zoogeography of Chironomidae from China.

YOUN, B. J.: Faunistic study of the larval chironomids (Diptera) in the stream of Kyong-sang Providence, South Korea.

ZORINA, O. V.: Fauna and systematics of the Tribe Chironomini (Diptera, Chironomidae) of the Russian Far East (Primo rye, Khabarovsk and Sahalin Territories).

NOTICE BOARD

NEW CHIRONOMID KEYS

The larval volume of Chironomidae of the Holarctic region (WIEDERHOLM, T. ed. 1983) is sold out and now longer available from the publisher. However, a new key to genera published in Hungary by Science Herald is just out:

SÆTHER, O. A., ASHE, P. & MURRAY, D. E. 2000. Family Chironomidae. Pp. 113-334 in: PAPP, L. and DARVAS, B. (eds): Contributions to a Manual of Palaearctic Diptera (with special reference to the flies of economic importance). Vol. 4. A.6.- Science Herald, Budapest. Volume 4 also includes keys to 15 other families of Diptera including Culicidae and Blepharoceridae.

Volume 4 is obtainable from E. W. Classey Ltd., Oxford House, Marlborough Street, Farringdon, Oxfordshire SW7 /JP, UK at a price of GBP 150, Order code 16272. The e-mail is: Peter@Classey.demon.co.uk.

The key mostly follows WIEDERHOLM (ed.) 1983, 1986, 1989, but especially the pupal keys to the orthoclads is improved. The keys also lists the names of species and their geographical distributions in revised genera with less than about 30 species. New or revised genera and subgenera appearing since the last volume of the Holarctic key are incorporated. There are several new synonyms and combinations. Japanese species are included as far as possible without revision of all species. The reference list is updated as far as possible, but some recent publications appearing after the first submission of the manuscript may be missing. The keys should be useful also for chironomid workers from North America as there are very few genera which are exclusively Nearctic.

Ole A. Sæther

NEW PUBLICATION

MASCHWITZ, DAVID E. and Edwin F. COOK. 2000. Revision of the Nearctic species of the genus *Polypedilum* KIEFFER (Diptera: Chironomidae) in the subgenus *P. (Polypedilum)* KIEFFER and *P. (Uresipedilum)* OYEWU and SÆTHER. Ohio Biological Survey Bulletin New Series Volume 12 Number 3. vii + 135 p.

This publication is available through the Ohio Biological Survey, 1315 Kinnear Road, Columbus, Ohio 43212-1192, U.S.A

KEY TO PUPAL EXUVIAE OF WEST PALAEARCTIC CHIRONOMIDAE UPDATE ON THE GENUS CHIRONOMUS 1995 (PAN-PALAEARCTIC VERSION)

Peter H. Langton

This update is no longer available and is under extensive revision. So far 77 taxa from the Palaearctic Region are included. I am extremely grateful to all those colleagues who have sent me associated pupal exuviae, particularly of karyotypically identified stock. Material of new species recently published or soon to appear would be much appreciated. Also, I have material of only a few eastern Palaearctic species and would be grateful for any positively named exuviae from that region.

LEN FERRINGTON NEW ADDRESS:

Beginning on 5 July 2000 Len Ferrington will have a new mailing address, telephone number and e-mail address.

The mailing address is: Leonard C. Ferrington Jr., Department of Entomology
Hodson Hall, 1980 Folwell Avenue, University of Minnesota, St. Paul, MN 55108

The new telephone number is: 612-624-3265

The new e-mail address is: ferri016@umn.edu" (Please note that the last three digits before @ are zero {not o}, one and six) (See also List of Regional Representatives)

PETER CRANSTON NEW ADDRESS

CRANSTON, Peter S., Department of Entomology, University of California, One Shields Avenue,
Davis, CA 95616, US Email: pscranston@ucdavis.edu

A REQUEST FOR ASSISTANCE

Art Borkent has undertaken a project to interpret the immatures (eggs, larvae, pupae) of Ceratopogonidae at a generic level (remember, they used to be a subfamily of the Chironomidae!). At the present time, there are generic keys available only for some European larvae and none for pupae. Such an undertaking requires associated material and if you have any reared specimens (or pupae with pharate adults) he would be very grateful for any loans. If you have such specimens, please contact Art at the following address:

Dr. A. Borkent 1171 Mallory Road, R1-S20-C43, Enderby, British Columbia,, V0E 1V0,
Canada E-mail: aborkent@jetstream.net, Ph (250) 833-0931, FAX (250) 832-2146

ANOTHER REQUEST FOR ASSISTANCE

Any specimens (larvae, pupae and especially adults-males and females of westpalaearctic *Glyptotendipes* s.str. - species: *Glyptotendipes mancurianus*, *Glyptotendipes foliicola*, *Glyptotendipes scirpi*, *Glyptotendipes imbecillis*, *Glyptotendipes varipes*, *Glyptotendipes viridis*, *Glyptotendipes caulicola*, *Glyptotendipes aequalis* and *Trichotendipes signatus* will be wellcome for the second part of my Revision of the westpalaearctic *Glyptotendipes*.

Please contact: **R. Contreras-Lichtenberg**, Naturhistorisches Museum Wien 2. Zoologische Abteilung
Burgring 7 P.O.Box 417 A-1014 VIENNA- AUSTRIA

**THE INTERNATIONAL CONFERENCE „SMALL RIVERS:MODERN
ECOLOGICAL STATE: ACTUAL PROBLEMS.**

APRIL 25 – 28, 2001, TOGLIATTI, RUSSIA

Dear colleagues,

Russian Academy of Science
Samara Scientific Centre of RAS
Institute of Ecology of Volga Basin of RAS
Hydrobiological Society of RAS
Moscow State University

invites you to attend the International Conference "Small rivers: Modern ecological State, actual problems". It is a pleasure to inform you that the Conference will be a part of the activities of the Institute of Ecology of the Volga Basin of the Russian Academy of Science.

At the conference the following questions are to be discussed:

1. Biodiversity, as an index of river system state. Bioindication and monitoring.
2. Influence of anthropogenic factors on structural – functional organisation of river ecosystem. Objective laws of organization and function.
3. Criteria and methods of anthropogenic load level and water quality estimation.
4. Balanced approach to river ecosystem investigation.
5. Methodological aspects of landscape zonation of rivers.
6. Hydroecological safety of small rivers.

There will be a special session for Chironomidae where all kind of investigations connected with this group of invertebrates are welcome. There will be sessions at the Conference for oral presentation and posters.

Instruction for authors:

TITLE (14, bold, capitals)

Names of authors (12, bold)

Organisation, City (12, italic)

Abstracts must not exceed 1 printing page, typed in single-line spacing in Times New Roman, 12 pt, with following margin width: upper and lower - 1,5 cm, left - 2,5 cm, right - 2 cm and should be sent to:

Organising Committee in two copies.

Electronic form: Files should be sent in Word 6 for Windows 95
in 3 1/2-inch diskette or via e-mail.

Registration form and abstracts should be sent to:

420008 Russia

Kazan, Kremliovskaya str., 18

Kazan State University, Ecological Faculty

Dr. Nazarova Larisa

e-mail: larisa.nazarova@ksu.ru

The invitation and the Conference program will be sent in the 2-nd Information letter in February of 2001.

List of Regional Representatives 2000

Javier ALOCER, UNAM Campus Iztacala, Aquatic Ecology Lab, Environmental Conservation & Improvement Project Calle 15 #51 San Pedro de los Pinos, Mexico City, D.F. **03800 – MEXICO** Email: jalocer@unam vm1.dgsca.unam.mx

Josef S. AMAKYE, Institute of Aquatic Biology (C.S.I.R.), P.O.Box 38, Achimota, Accra, GHANA - **WEST AFRICA** Phone: +233 ? 775511

Peter BITUSIK, Faculty of Ecology, Technical University, Masarykova 24, 960 53 Zvolen - **SLOVAKIA**

Ian K.G. BOOTHROYD, National Institute of Water and Atmospheric Research, P. O. Box 11-115, Hamilton, - **NEW ZEALAND** Phone: 07 8561737; Fax: 07 8560151; Email: (work) I.boothroyd@niwa.cri.nz (home) Ian.b@xtra.co.nz"

Arnoldine BURGOS, Bureau voor Openbare Gezondheidszorg, Centraal Laboratorium, Rode Kruislaan 13 Postbus 1911, Paramaribo - **SURINAME**

Youn BYEONG-JIN, Pusan National University, Department of Biology, #30 Changjeon-dong Kumjeong-Ku, Pusan 609-735 - **SOUTH KOREA**

Zenaida Batac CATALAN, Institute of Environmental Science and Management, The University of the Philippines at Los Banos, College Laguna - **PHILIPPINES**

Marcos CALLISTO, Lab. Ecologia de Bentos/ liminologia - Universidade Federal de Minas Gerais, ICB, Depto. Biologia Geral, Lab. Limnologia/Ecologia de Bentos, CP. 486, CEP. 30.161-970, Belo Horizonte, MG, **BRAZIL**. Brazil Tel: +55 31 499 25 95. Fax: +55 31 499 25 67. Email : callisto@mono.icb.ufmg.br"

P.K. CHAUDHURI, Department of Zoology - Entomology, University of Burdwan, Burdwan - 713104 - W.B. **INDIA** Phone: +91 BDN 2371-75 [37]

Pavel CHVOJKA, National Museum, Department of Entomology, Kunratice 1, 14800 Praha - **CZECH REPUBLIC**

Ruth CONTRERAS, Naturhistorisches Museum, 2.Zoologische Abteilung, Burgring 7, P.O.Box 417 - **AUSTRIA** Phone: ++43 01 521 77/317; Fax: ++43 01 523 52 54; E-mail: Ruth.Contreras@nhm-wien.ac.at"

Peter S. CRANSTON, Department of Entomology, University of California, One Shields Avenue, Davis, CA 95616, US

AUSTRALIA Email: pscranston@ucdavis.edu

Carlos DE LA ROSA, Brooker Creek Preserve, Pinellas Co. Environmental Mgmt. Dept., 1001 Lora Lane, Tarpon Springs, Florida 34689 - U.S. **CENTRAL AMERICA** Phone: (727) 943-4004; Cell Phone: (727) 580-2573; Fax: (727) 943-4002. Email: cdelaros@co.pinellas.fl.us

György DEVAI, L. Kossuth University, Ecological Institute, H4010 Debrecen - **HUNGARY**

Engelsina ERBAEVA, Institute of Biology, University of Irkutsk, P.O.Box 24, 664003 Irkutsk - Russia, **LAKE BAIKAL AND RIVER ANGARA, SOUTH SIBERIA** Phone +7? 243079 or 242186 e-mail: root@bio.isu.runnet.ru

Leonard C. FERRINGTON Jr Department of Entomology Hodson Hall, 1980 Folwell Avenue University of Minnesota St. Paul, MN 55108 **U.S.A.** Phone: + 612-624-3265 E-mail: ferri016@umn.edu

Boudewijn GODDEERIS, Koninklijk Belgisch Instituut voor Natuurwetenschappen, Vautierstraat 29, B-1040 Brussel - **BELGIUM** Phone: +32 2 6274314; Fax: 32 2 6464433 E-mail: godderi@kbinirsnb.be

Arthur HARRISON, 111 A Berg Road, Fish Hoek 7975 - **SOUTH AFRICA**

A.R. ISMAIL, Jabatan Biologie, University Pertanian Malaysia, 43400 UPM Serdang, Selangor - **MALAYSIA**

Toshio IWAKUMA, National Institute for Environmental Studies, 16-2 Onogawa, Tsukuba, Ibaraki 305 - **JAPAN** Phone: 0298 51 6111; Fax: 0298 51 4732 E-mail: iwakuma@ees.hokudai.ac.jp

Richard K. JOHNSON, University of Agricultural Sciences, P.O. Box 7050, S-75007 Uppsala - **SWEDEN** Fax: +46 18673156 E-mail: Richard.Johnson@ma.slu.se

Kulli KANGUR, Vvrstjdrv Limnological Station, EE2454 Rannu - **ESTONIA** E-mail: andu@lim.tartu.ee

Esa KOSKENNIEMI, National Board of Waters and the Environment Water and Environment District of Vaasa, Vaasan Vesi, Ja Ymparistopiiri, P.O. Bo. 262, SF-65101

Vaasa - **FINLAND** Fax: +358 (9)61 256601 E-mail: Esa.Koskenniemi@vyh.fi

Adrzej KOWNACKI, Institute of Freshwater Biology - Polish Academy of Sciences, ul.Slawkowska 17, PL-31-016 Krakow - **POLAND** Phone: +48 12 222115

Jehoshua KUGLER, Department of Zoology, Tel Aviv University, Tel Aviv 69978 - **ISRAEL**

Peter H. LANGTON, 5 Kylebeg Avenue, Mountsandel, Coleraine, Co. Londonderry, Northern Ireland, BT52 1JN - Northern Ireland

GREAT BRITAIN E-mail: 106375.1453@compuserve.com

Henri LAVILLE, Laboratoire d'Hydrobiologie, Université Paul Sabatier, 118. Route de Narbonne, F-31062 Toulouse Cedex - **FRANCE** Fax: +61 55 6470 or +61 55 6000 E-mail: limnol@cict.fr

Heather A. LESLIE, Department of Aquatic Ecology and Ecotoxicology, ARISE, University of Amsterdam, Kruislaan 320, 1098 SM Amsterdam - **THE NETHERLANDS** Tel. +31-(0)20-525 77 18. Fax: +31-(0) 20-525 77 09. Email: leslie@bio.uva.nl "

Claus LINDEGAARD, Freshwater Biological Laboratory, University of Copenhagen, 51 Helsingorsgade, DK-3400 Hillerød - **DENMARK** E-mail: linddegaard@ibm.net

Brigitte LODS-CROZET, Laboratoire d'Ecologie et de Biologie Aquatique, 18 Chemin des Clochettes, University of Geneva, CH-1206 GENEVE - **SWITZERLAND** Tel: (+41 22) 705 71 00; fax (+41 22) 789 49 89; E-mail: Brigitte.Lods@leba.unige.ch

Eugeniy A. MAKARCHENKO, Institute of Biology and Pedology, Far Eastern Branch of the Russian Academy of Sciences, 690022 Vladivostok, Russia - **FAR EAST OF RUSSIA** Fax: 0423 225 8484 E-mail: emakar@mail.ru (home), zoology@eastnet.febras.ru (work)

Julieta MASAFERRO, Department of Entomology, Natural History Museum, Cromwell Road, London SW7 5BD, U.K. **ARGENTINA** Tel: (0) 207 942 5198; Fax: (0) 207 942 5229; Email: J.Massaferrero@nhm.ac.uk

Paraskeva MICHAILOVA, Institute of Zoology, Bulgarian Academy of Sciences, Institute of Zoology, Boul. Osvoboditel 1, Sofia 1000 - **BULGARIA** E-mail: parmich@mail.bol.bg

Seyed Karim MOUSAYI, Department of Fisheries and Aquatics, College of Natural

Resources, University of Tehran, Karaj 31584 - **IRAN** E-mail: karim@nfh.uit.no

Declan MURRAY, Department of Zoology, University College Dublin, Belfield, Dublin 4 - **IRELAND** Phone: +353 1 706 2336; Fax: +353 1 706 1152; E-mail: DMURRAY@ollamh.ucd.ie

JON SIGURDUR OLAFSSON, University of Iceland, Institute of Biology, Grensasvegi 12, 108 Reykjavik - **ICELAND** Email: jsol@rhi.hi.is

Narcis PRAT, Department d'Ecologia, University of Barcelona, Avgda. Diagonal 645, ES-08028 Barcelona - **SPAIN** Fax: +34 3 4111438 E-mail: narcis@porthos.bio.ub.es

Bruno ROSSARO, Univ. of Milano, Dept. Biology, Sect. Ecology, via Celoria 26, I 20133, Milano - **ITALY** Fax: +39-2-26604261 E-mail: rossaro@mailserver.unimi.it

Martin SPIES, Schraemelstr. 151, D-81247 München - **GERMANY** Email: spies@zi.biologie.uni-muenchen.de

Maria-Monica TUDORANCEA "Babes-Bolyai" University Faculty of Biology and Geology Department of Ecology-Genetics 1 M. Kogalniceanu Str. Cluj-Napoca, 3400 **ROMANIA** Fax: (40) (64) 191906 or 40 64 431858 Email: mtudor@biolog.ubbcluj.ro"

Ian R. WALKER, Department of Biology, Okanagan University College, 3333 College Way, Kelowna, British Columbia V1V 1V7 - **CANADA** Phone: (250) 762-5445 Ext. 7559; Fax: (250) 470-6004; E-mail: iwalker@okanagan.bc.ca

Xinhua WANG, Dept. of Biology, Nankai University, Tianjin, 300071. - **P. R. CHINA** Fax: +86 22 344853 E-mail: xhwang@sun.nankai.edu.cn

Endre WILLASSEN, Museum of Zoology, University of Bergen, Museplass 3, N-5007 Bergen - **NORWAY** Fax: +47 5 321153 . Email: Endre.Willassen@zmb.uib.no

Tatiana ZINCHENCO, Institute of Ecology of the Volga River Basin, Russian Academy of Sciences, 445003 Togliatti, Russia - **TOGLIATTI**

**DEADLINE FOR CHIRONOMUS 14
1st of July 2001**

Current Bibliography: 1 Jan. 1998 - 31 Dec. 1999

by Odwin Hoffrichter

[© 2000 by CHIRONOMUS, and O. Hoffrichter - Associated Editor]

Supplement to 1998 Current Bibliography:

additions & corrections (earlier parts only to be found on the Chironomid home page of the Internet).

- Andreeva, E. N., Kiknadze, I. I. i Aimanova, K. G. 1998a. Sravnitel'nyi analiz risunka diskov politennykh khromosom blizkikh vidov *Glyptotendipes salinus* Michailova i *G. barbipes* Staeger (Diptera, Chironomidae). (Comparative analysis of polytene chromosome banding pattern of closely related species *Glyptotendipes salinus* Michailova and *G. barbipes* Staeger (Diptera, Chironomidae).- *Tsitologiya* 40: 972-979.
- Ashe, P., O'Connor, J. P. and Murray, D. A. 1999a. A checklist of Irish aquatic insects.- *Occ. Publ. Ir. biogeogr. Soc.* 3: 80 pp.
- Banning, M. 1998a. Auswirkungen des Aufstaus größerer Flüsse auf das Makrozoobenthos - dargestellt am Beispiel der Donau.- *Essen. ökol. Schr.* 9: 285 + 88 pp. Westarp Wiss. Verl., Hohenwarsleben.
- Benbow, M. E., Burky, A. J. and Way, C. M. 1998a. Influence of microhabitat stream flow on larval biomass, sexual dimorphism, and sex ratio of an endemic Hawaiian midge.- *Bull. ecol. Soc. Am.* 79, Suppl.: 147.
- Bouckaert, F. W. and Davis, J. 1998a. Microflow regimes and the distribution of macroinvertebrates around stream boulders.- *Freshwat. Biol.* 40: 77-86.
- Boulton, A. J. and Fosdter, J. G. 1998a. Effects of buried leaf litter and vertical hydrologic exchange on hyporheic water chemistry and fauna in a gravel-bed river in northern New South Wales, Australia.- *Freshwat. Biol.* 40: 229-243.
- Burel, F., Baudry, J., Butet, A., Clergeau, P., Delettre, Y., Le Coeur, D., Dubs, F., Morvan, N., Paillat, G., Petit, S., Thenail, C., Brunel, E. and Lefeuvre, J.-C. 1998a. Comparative biodiversity along a gradient of agricultural landscapes.- *Acta oecol.* 19: 47-60.
- Burky, A. J., Benbow, M. E. and Way, C. M. 1998a. Larval sexual dimorphism and sex ratio of a large, tropical chironomid.- *Bull. N. Am. benthol. Soc.* 15: 218.
- Burky, A. J., Benbow, M. E., Driscoll, K., Dwyer, P. M., Kearns, M. F., Valaitis, J. S. and Way, C. M. 1998a. The influence of stream discharge and drainage basin size on macroinvertebrate drift from Hawaiian streams.- *Bull. N. Am. benthol. Soc.* 15: 214.
- Chattopadhyay, S. and Chaudhuri, P. K. 1998b. Life stages of *Dicrotendipes pelochloris* (Kieffer) with a note on its biology (Diptera: Chironomidae).- *Proc. zool. Soc. Calcutta* 51: 167-177.
- Clenaghan, C., Giller, P. S., O'Halloran, J. and Hernan, R. 1998a. Stream macroinvertebrate communities in a conifer-afforested catchment in Ireland: relationships to physico-chemical and biotic factors.- *Freshwat. Biol.* 40: 175-193.
- Cranston, P. S. 1998b. The Chironomidae (Diptera) of the Musselbrook region.- *R. geogr. Soc. Qd Geogr. Monogr. Ser.* 4: 59-66.
- Dole-Olivier, M.-J. 1998a. Surface water-groundwater exchanges in three dimensions on a backwater of the Rhône River.- *Freshwat. Biol.* 40: 93-109.
- Felder, D. P., D'Surney, S. J., Rodgers, J. H. Jr. and Deardorff, T. L. 1998a. A comprehensive environmental assessment of a receiving aquatic system near an unbleached kraft mill.- *Ecotoxicology* 7: 313-324.
- Guerra, J. A. de, Guerra Martinez, A. y Rebaza Vigo, M. (1997) 1998a. (The "red worm of rice" in Jequetepeque Valley, La Libertad, Peru).- *Revta per. Ent.* 40: 139-153.
- Hickey, C. W. and Clements, W. H. 1998a. Effects of heavy metals on benthic macroinvertebrate communities in New Zealand streams.- *Envir. Toxic. Chemy* 17: 2336-2346.
- Hillbricht-Ilkowska, A., Dusoge, K., Ejsmont-Karabin, J., Jasser, I., Kufel, I., Ozimek, T., Rybak, J. I., Rzepecki, M. and Wegle_ska, T. 1998a. Long term effects of liming in a humic lake: Ecosystem processes, biodiversity, food web functioning (Lake Flosek, Masurian Lakeland, Poland).- *Pol. J. Ecol.* 46: 347-415.
- Iannacone, J. A. y Alvariano, L. 1998a. Ecotoxicidad aguda del insecticida organofosforado temephos sobre *Chironomus calligraphus* Goeldi (Diptera: Chironomidae).- *Acta ent. chilena* 22: 53-55.

- Jacobsen, R. E. 1998a. *The symbiotic relationship and emergence phenology of three Chironomidae associated with the burrowing mayfly, Ephemera guttulata Pictet (Ephemeroptera: Ephemeridae).*- Ph. D. Thes., Univ. Pittsburgh. 144 pp.
- Jankovi, M. 1998a. The chironomid fauna of the Velika Morava river.- *Zb. Radova Fauni Srbije* 5: 105-139.
- Kakareko, T. 1998a. On the ecology of bream *Abramis brama* of the Woc_awek Dam Reservoir.- *Int. Rev. Hydrobiol.* 83, spec. Iss.: 409-412.
- Kiknadze, I. I., Andreeva, E. N., Istomina, A. G. i Butler, M. G. 1998a. Kariofond golarktičeskoj khironomidy *Glyptotendipes barbipes* (Staeger). (Karyofund of Holarctic midge *Glyptotendipes barbipes* (Staeger).)- *Tsitologiya* 40: 900-912.
- Lang, C. 1998a. Qualité biologique de 29 rivières vaudoises en 1997 indiquée par la diversité du zoobenthos.- *Bull. Soc. vaud. Sci. nat.* 86: 61-71.
- Liber, K., Schmude, K. L. and Rau, D. M. 1998a. Toxicity of *Bacillus thuringiensis* var. *israelensis* to chironomids in pond mesocosms.- *Ecotoxicology* 7: 343-354.
- Paggi, A. C. 1998a. Chironomidae.- In: Morrone, J. J. y Coscaron, S. (eds.): *Biodiversidad de artropodos argentinos: una perspectiva biotaxonomica*, pp. 327-337. Ed. SUR, La Plata.
- Pelli, A. e Barbosa, F. A. R. 1998b. Insetos coletados em *Salvinia molesta* Mitchell (Salviniaceae), com especial referência às espécies que causam dano à planta, na lagoa Olhos d'Água, Minas Gerais, Brasil.- *Revta bras. Ent.* 42: 9-12.
- Prat, N. y Rieradevall, M. 1998a. Criterios de evaluación de la calidad del agua en lagos y embalses basados en los macroinvertebrados bentónicos.- *Actual. Biol.* 20: 137-147.
- Rey, D., Pautou, M. and Meyran, J. 1998a. Histopathological effects of tannic acid on the midgut epithelium of some aquatic Diptera larvae.- *J. Invert. Path.* 73: 173-181.
- Saigusa, M. and Oishi, K. 1998a. Patterns of emergence in marine invertebrates: on the influence of a field light.- *Benthos Res.* 53: 95-104.
- Sasa, M., Suzuki, H. and Sakai, T. 1998b. Studies on the chironomid midges collected on the shore of Shimanto River in April, 1998. Part 2. Description of additional species belonging to Orthocladiinae, Diamesinae and Tanypodinae.- *Trop. Med.* 40: 99-147.
- Sheldon, F. and Puckridge, J. T. 1998a. Macroinvertebrate assemblages of Goyder Lagoon, Diamantina River, South Australia.- *Trans. R. Soc. S. Aust.* 122: 17-31.
- Shilova, A. I. 1998a. Novyi dlya Rossii vid khironomid *Corynocera oliveri* Lindeberg (Diptera, Chironomidae). (*Corynocera oliveri* Lindeberg (Diptera, Chironomidae), a new for the fauna of Russia species of Chironomidae, with redescription of male.)- *Ent. Obozr.* 77: 687-691.
- Spindler-Barth, M. and Spindler, K.-D. 1998b. Morphogenetic actions and mode of action of ecdysteroids in a dipteran cell line.- *Curr. Topics Steroid Res.* 1: 73-81.
- Tátrai, I., Paulovits, G., Mátyás, K., Korponai, J., and Pomogyi, P. 1998a. Long-term food web interactions in Kis-Balaton Reservoir.- *Int. Rev. Hydrobiol.* 83, spec. Iss.: 515-522.
- Tolonen, A. 1998a. Application of a bioenergetics model for analysis of growth and food consumption of subarctic whitefish *Coregonus lavaretus* (L.) in Lake Kilpisjärvi, Finnish Lapland.- *Hydrobiologia* 390: 153-169.
- Trayler, K. M. and Davis, J. A. 1998a. Forestry impacts and the vertical distribution of stream invertebrates in south-western Australia.- *Freshwat. Biol.* 40: 331-342.
- Vermeulen, A. C. 1998a. *Head capsule deformation in Chironomus riparius (Diptera): causality, ontogenesis and its application in biomonitoring.*- Ph. D. Thes., Univ. Leuven. 207 pp.
- Vrsansky, P. and Wagner, J. 1998a. Facets of fossil chironomids (Diptera: Chironomidae).- *Ent. Probl.* 29: 94.
- Vuori, K.-M., Joensuu, I., Latvala, J., Jutila, E. and Ahvonen, A. 1998a. Forest drainage: a threat to benthic biodiversity of boreal headwater streams?- *Aquat. Conserv. mar. Freshwat. Ecosyst.* 8: 745-759.
- Winterbourn, M. J. 1998a. Insect faunas of acidic coal mine drainages in Westland, New Zealand.- *N. Z. Ent.* 21: 65-72.
- Wi_niewski, J. R. 1998a. High mobility group proteins: components that modulate insect chromatin.- *Zool. Pol.* 43: 5-24.
- [to be deleted: Lobinske, R. J., Ali, A. and Stout, I. J. 1998a. Benthic macroinvertebrates and selected physico-chemical parameters in two tributaries of the Wekiva River, central Florida, USA.- *Freshwat. Biol.* 48: 219-231 - correct in Curr. Bibl. 1997!]

Current Bibliography 1998

- Adam, J. I. and Sæther, O. A. 1999a. Revision of the genus *Nilothauma* Kieffer, 1921 (Diptera: Chironomidae).- *Ent. scand. Suppl. 56*: 1-107.
- Adkins, S. C. and Winterbourn, M. J. 1999a. Vertical distribution and abundance of invertebrates in two New Zealand stream beds: a freeze coring study.- *Hydrobiologia 400*: 55-62.
- Alam, N. and Shafi, 1999a. Histopathological lesions induced by metacid 50 in the *Chironomus* larvae (Chironomidae: Diptera).- *Envir. Ecol. 17*: 637-639.
- Alzmann, N., Köhler, B. and Maier, G. 1999a. Spatial distribution, food and activity of *Gomphus pulchellus* SELYS 1840 (Insecta; Odonata; Gomphidae) from a still water habitat.- *Int. Rev. Hydrobiol. 84*: 299-313.
- Amaral, M. J. and Costa, M. H. 1999a. Macroinvertebrate communities of salt pans from the Sado estuary (Portugal).- *Acta oecol. 20*: 327-332.
- Andersen, T. and Contreras-Ramos, A. 1999a. First record of *Antillocladius* (SÆTHER) from continental South America (Chironomidae, Orthocladiinae).- *Acta zool. Acad. Scient. hung. 45*: 149-154.
- Andersen, T. and Kristoffersen, L. (1998)1999a. New species of *Xestochironomus* Sublette and Wirth (Chironomidae:Chironominae) from Chile and Costa Rica.- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae. - J. Kans. ent. Soc. 71*:
- Angradi, T. R. 1999a. Fine sediment and macroinvertebrate assemblages in Appalachian streams: a field experiment with biomonitoring applications.- *J. N. Am. benthol. Soc. 18*: 49-66.
- Anido, A. E, Cox, C. K. and Case, S. T. 1999a. Molecular fate of Balbiani ring 4 in *Chironomus tentans*.- *Molec. Biol. Cell 10 (Suppl.)*: 91a.
- Augustin, J. C. Grubaugh, J. W. and Marshall, M. R. 1999a. Validating macroinvertebrate assumptions of the shorebird management model for the lower Mississippi Valley.- *Wildl. Soc. Bull. 27*: 552-558.
- Bachmann, V. and Usseglio-Polatera, P. 1999a. Contribution of the macroinvertebrate compartment to the oxygen budget of a large regulated river: the Mosel.- *Hydrobiologia 410*: 39-46.
- Batzer, D. P., Shurtleff, A. S. and Robinette, J. R. 1999a. Managing fish and invertebrate resources in a wood stork feeding pond.- *J. Freshwat. Ecol. 14*: 159-165.
- Beaudoin, C. P., Tonn, W. M., Prepas, E. E. and Wassenaar, L. I. 1999a. Individual specialization and trophic adaptability of northern pike (*Esox lucius*): an isotope and dietary analysis.- *Oecologia 120*: 386-396.
- Beauvais, S. L., Atchison, G. J., Stenback, J. Z. and Crumpton, W. G. 1999a. Use of cholinesterase activity to monitor exposure of *Chironomus riparius* (Diptera : Chironomidae) to a pesticide mixture in hypoxic wetland mesocosms.- *Hydrobiologia 416*: 163-170.
- Belyanina, S. I., Durnova, N. A. and Shobanov, N. A. 1999a. Osobennosti kariotipa i morfologii *Glyptotendipes* sp. (Diptera, Chironomidae) s khromosomnym naborom 2n = 4. (Morphology and karyotype of *Glyptotendipes* sp. (Diptera, Chironomidae) with 2n = 4.)- *Zool. Zh. 78*: 1434-1440.
- Benbow, M. E. 1999a. *Natural history and bioenergetics of an endemic Hawaiian chironomid: Fluctuating stability in a stochastic environment.*- Diss., Univ. Dayton, Dayton, OH. 394 pp.
- Benbow, M. E., Burky, A. J. and Way, C. M. 1999a. Effects of stream flow on tropical chironomid populations: a comparison of differential flow habitat on instar standing stock biomass.- *Bull. N. Am. benthol. Soc. 16*: 209.
- Bendell-Young, L. I. 1999a. Application of a kinetic model of bioaccumulation across a pH and salinity gradient for the prediction of cadmium uptake by the sediment dwelling Chironomidae.- *Envir. Sci. Technol. 33*: 1501-1508.
- Benke, A. C., Huryn, A. D., Smock, L. A. and Wallace, J. B. 1999a. Length-mass relationships for freshwater macroinvertebrates in North America with particular reference to the southwestern United States.- *J. N. Am. benthol. Soc. 18*: 308-343.
- Berezina, N. A 1999a. Tolerance of *Omisus caledonicus* (Diptera: Chironomidae) to changes in salinity, pH, and ionic composition of water.- *Russ. J. Ecol. 30*: 58-60.
- Berezina, N. A. 1999b. Osobennosti razvitiya makrozoobentosnykh soobshchestv pod vliyaniem dreisseny (*Dreissena polymorpha* Pall.) v eksperimental'nykh mezokosmakh. (Peculiarities of development of macrozoobenthos communities under influence of *Dreissena polymorpha* Pall. in experimental mesocosms.)- *Zh. obshch. Biol. 60*: 189-198.
- Bervoets, L. and Blust, R. 1999a. Bioavailability of cadmium and zinc to midge larvae under natural and experimental conditions: Effects of some environmental factors.- *Belg. J. Zool. 129*: 269-284.

- Bhattacharya, G., Mazumdar, A. and Chaudhuri, P. K. 1999a. Incidence of deformed *Chironomus* larvae in contaminated sediment of the River Damodar, West Bengal (Diptera: Chironomidae).- *Pollut. Res.* 18: 79-82.
- Bianchi, M. M., Massaferro, J., Roman Ross, G., Amos, A. J. and Lami, A. 1999a. Late pleistocene and early Holocene ecological response of Lake El Trebol (Patagonia, Argentina) to environmental changes.- *J. Paleolimnol.* 22: 137-148.
- Bleeker, E. A. J., Leslie, H. A., Groenendijk, D., Plans, M. and Admiraal, W. 1999a. Effects of exposure to azaarenes on emergence and mouthpart development in the midge *Chironomus riparius* (Diptera: Chironomidae).- *Envir. Toxic. Chem* 18: 1829-1834.
- Bohonak, A. J. 1999a. Effect of insect-mediated dispersal on the genetic structure of postglacial water mite populations.- *Heredity* 82: 451-461.
- Boothroyd, I. K. G. (1998)1999a. Size distribution of *Maoridiamesa stouti* Brundin (Diptera: Chironomidae): a madicolous chironomid from New Zealand.- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc.* 71: 484-489.
- Boothroyd, I. K. G. 1999b. Description of *Kaniwhaniwhanus* gen. n. (Diptera: Chironomidae: Orthoclaadiinae) from New Zealand.- *N. Z. Jl mar. Freshwat. Res.* 33: 341-349.
- Boothroyd, I. K. G. 1999c. Life history of *Kaniwhaniwhanus chapmani* Boothroyd (Chironomidae: Orthoclaadiinae): population dynamics, emergence, and drift.- *N. Z. Jl mar. Freshwat. Res.* 33: 351-359.
- Boothroyd, I. and Cranston, P. 1999a. The 'ice-worm' - the immature stages, phylogeny and biology of the glacier midge *Zelandochlus* (Diptera: Chironomidae).- *Aquat. Insects* 21: 303-316.
- Borgmann, U. and Norwood, W. P. 1999a. Sediment toxicity testing using large water-sediment ratios: An alternative to water renewal.- *Envir. Pollut.* 106: 333-339.
- Bott, T. L. and Borchardt, M. A. 1999a. Grazing of protozoa, bacteria, and diatoms by meiofauna in lotic epibenthic communities.- *J. N. Am. benthol. Soc.* 18: 499-513.
- Botts, P. S. 1999a. Lake Erie coastal wetlands. A review and case study of Presque Isle invertebrates.- In: Batzer, D. P., Rader, R. B. and Wissinger, S. A. (eds.): *Invertebrates in freshwater wetlands of North America: ecology and management*, pp. 995-1012. John Wiley & Sons, N. Y., Chichester.
- Bretschko, G. 1999a. The lotic fauna in water-logged sediments beyond the waterline in the canalized Danube.- *Arch. Hydrobiol. Supp.* 115: 413-421.
- Brodersen, K. P. and Lindegaard, C. 1999a. Mass occurrence and sporadic distribution of *Corynocera ambigua* Zetterstedt (Diptera, Chironomidae) in Danish lakes. Neo- and palaeolimnological records.- *J. Paleolimnol.* 22: 41-52.
- Brodersen, K. P. and Lindegaard, C. 1999b. Classification, assessment and trophic reconstruction of Danish lakes using chironomids.- *Freshwat. Biol.* 42: 143-157.
- Brunke, M. and Gonser, T. 1999a. Hyporheic invertebrates - the clinal nature of interstitial communities structured by hydrological exchange and environmental gradients.- *J. N. Am. benthol. Soc.* 18: 344-362.
- Burmester, T. and Hankeln, T. 1999a. A globin gene of *Drosophila melanogaster*.- *Molec. Biol. Evol.* 16: 1809-1811.
- Butler, M. G., Kiknadze, I. I., Golygina, V. V., Martin, J., Istomina, A. G., Wülker, W. F., Sublette, J. E. and Sublette, M. F. 1999a. Cytogenetic differentiation between Palearctic and Nearctic populations of *Chironomus plumosus* L. (Diptera, Chironomidae).- *Genome* 42: 797-815.
- Caldwell, B. A. 1999a. Description of the adult male and larva of *Orthocladius (Orthocladius) vaillanti* (Diptera: Chironomidae).- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc.* 71: 234-240.
- Call, D. J., Liber, K., Whiteman, F. W., Dawson, T. D. and Brooke, L. T. 1999a. Observations on the 10-day *Chironomus tentans* survival and growth bioassay in evaluating Great Lakes sediments.- *J. Gt Lakes Res.* 25: 171-178.
- Call, D. J., Polkinghorne, C. N., Markee, T. E., Brooke, L. T., Geiger, D. L., Gorsuch, J. W. and Robillard, K. A. 1999a. Silver toxicity in *Chironomus tentans* in two freshwater sediments.- *Envir. Toxic. Chem* 18: 30-39.
- Callisto, M., Esteves, F. A., Gonçalves, J. F. and Leal, J. J. F. (1998)1999a. Impact of bauxite tailings on the distribution of benthic macrofauna in a small river ('igarapé') in Central Amazonia, Brazil.- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1:*

- Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc.* 71: 447-451.
- Cano, E., Jiménez, A., Cabral, J. A. and Ocete, M. E. 1999a. Acute toxicity of malathion and the new surfactant "genapol OXD 080" on species of rice basins.- *Bull. envir. Contam. Toxic.* 63:133-138.
- Carter, C. E. and McLarnon, L. A. 1999a. *Monodiamesa ekmani* (Diptera: Chironomidae) [sic!] in Lough Neagh, Northern Ireland.- *J. Freshwat. Ecol.* 14: 343-348.
- Case, S. T. and Thornton, J. R. 1999a. High molecular mass complexes of aquatic silk proteins.- *Int. J. Biol. Macromolec.* 24: 89-101.
- Chattopadhyay, S. 1999a. Mating activity and behaviour of *Chironomus samoensis* Edwards (Diptera: Chironomidae).- *Geobios* 26: 208-210.
- Choi, J., Roche, H. and Caquet, T. 1999a. Characterization of superoxide dismutase activity in *Chironomus riparius* Mg. (Diptera, Chironomidae) larvae - a potential biomarker.- *Comp. Biochem. Physiol.* 124C: 73-81.
- Chou, R. Y. M., Ferrington, L. C. Jr., Hayford, B. L. and Smith, H. M. 1999a. Composition and phenology of Chironomidae (Diptera) from an intermittent stream in Kansas.- *Arch. Hydrobiol.* 147: 35-64.
- Claret, C., Marmonier, P., Dole-Olivier, M.-J. and Castella, E. 1999a. Effects of management works on the interstitial fauna of floodplain aquatic systems (River Rhône, France).- *Biodivers. Conserv.* 8: 1179-1204.
- Cobo, F. (1998)1999a. Valor de conservación e interés faunístico del poblamiento de quironómidos (Diptera. Chironomidae) en los Monegros (Zaragoza, España).- *Boln Soc. ent. aragon.* 24: 138.
- Coffman, W. P. and de la Rosa, C. L. (1998)1999a. Taxonomic composition and temporal organization of tropical and temperate species assemblages of lotic Chironomidae.- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 2: Biology, ecology and natural history of Chironomidae.* - *J. Kans. ent. Soc.* 71: 388-406.
- Collier, K. J., Champion, P. D. and Croker, G. F. 1999a. Patch- and reach-scale dynamics of a macrophyte-invertebrate system in a New Zealand lowland stream.- *Hydrobiologia* 392: 89-97.
- Conrad, A. U., Fleming, R. J. and Crane, M. 1999a. Laboratory and field response of *Chironomus riparius* to a pyrethroid insecticide.- *Wat. Res.* 33: 1603-1610.
- Contreras-Lichtenberg, R. 1999a. Revision der westpaläarktischen Arten des Genus *Glyptotendipes* KIEFFER, 1913 (Insecta: Diptera: Chironomidae). Teil 1: Subgenus *Phytotendipes* GOETGHEBUER, 1937.- *Annln naturhist. Mus. Wien 101 B*: 359-403.
- Corley, J. C. and Massaferrero, J. I. (1998)1999a. Long term turnover of a fossil community of chironomids (Diptera) from lake Mascardi (Patagonia, Argentina).- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 2: Biology, ecology and natural history of Chironomidae.* - *J. Kans. ent. Soc.* 71: 407-413.
- Craig, A., Hare, L. and Tessier, A. 1999a. Experimental evidence for cadmium uptake via calcium channels in the aquatic insect *Chironomus staegeri*.- *Aquat. Toxic.* 44: 255-262.
- Cranston, P. S. (1998)1999a. Nearctic *Orthocladius* subgenus *Eudactylocladius* revised (Diptera: Chironomidae).- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc.* 71: 272-295.
- Cranston, P. S. 1999b. Two unusual Chironomini (Diptera: Chironomidae) from Australian rainforest streams: One new genus and a neotropical genus new for the region.- *Aust. J. Ent.* 38: 291-299.
- Cranston, P. S. and Edwards, D. H. D. 1999a. *Botryocladus* gen. n.: a new transantarctic genus of orthoclaadiine midge (Diptera: Chironomidae).- *Syst. Ent.* 24: 305-333.
- Dahl, J. and Greenberg, L. 1999a. Effects of prey dispersal on predator-prey interactions in streams.- *Freshwat. Biol.* 41: 771-780.
- Dallas, H. F., Janssens, M. P. and Day, J. A. 1999a. An aquatic macroinvertebrate and chemical database for riverine ecosystems.- *Wat. SA* 25: 1-8.
- Daneholt, B. 1999a. Pre-mRNP particles: From gene to nuclear pore.- *Curr. Biol.* 9: R412-R415.
- Davies, N. A., Edwards, P. A., Lawrence, M. A. M., Simkiss, K. and Taylor, M. G. 1999a. Biocide testing using particles with controlled surface properties (artificial sediments).- *Envir. Toxic. Chem* 18: 2337-2342.
- De Jaegher, C. and Goossens, A. 1999a. Protein contact dermatitis from midge larvae (*Chironomus thummi thummi*).- *Contact Dermatitis* 41: 173.
- Dinsmore, W. P., Scrimgeour, G. J. and Prepas, E. E. 1999a. Empirical relationships between profundal macroinvertebrate biomass and

- environmental variables in boreal lakes of Alberta, Canada.- *Freshwat. Biol.* 41: 91-100.
- Donley, S., Ferrington, L. C. Jr. and Strayer, D. (1998)1999a. The habitat of *Paraboreochlus* larvae (Diptera: Chironomidae).- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc.* 71: 501-504.
- Dorava, J. M. and Milner, A. M. 1999a. Effects of recent volcanic eruptions on aquatic habitat in the Drift River, Alaska, USA: implications at other Cook Inlet region volcanoes.- *Envir. Mgmt* 23: 217-230.
- Doucett, R. R., Giberson, D. J. and Power, G. 1999a. Parasitic association of *Nanocladius* (Diptera: Chironomidae) and *Pteronarcys biloba* (Plecoptera: Pteronarcyidae): insights from stable-isotope analysis.- *J. N. Am. benthol. Soc.* 18: 514-523.
- Driscoll, M. P. and Miranda, L. E. 1999a. Diet ecology of yellow bass, *Morone mississippiensis*, in an oxbow of the Mississippi River.- *J. Freshwat. Ecol.* 14: 477-486.
- Dudgeon, D. and Wu, K. K. Y. 1999a. Leaf litter in a tropical stream: food or substrate for macroinvertebrates?- *Arch. Hydrobiol.* 146: 65-82.
- Edlund, M. B. and Francis, D. R. 1999a. Diet and habitat characteristics of *Pagastiella ostansa* (Diptera: Chironomidae).- *J. Freshwat. Ecol.* 14: 293-300.
- Edwards, D. D. 1999a. The behavioral responses by larvae of the water mite *Unionicola foili* (Acari: Unionicolidae) to gravity, light and host chemicals: adaptations for transmission success?- *Exp. appl. Acarol.* 23: 817-826.
- Egyhazi, E., Ossoinak, A., Filhol-Cochet, O., Cochet, C. and Pigon, A. 1999a. The binding of the alpha subunit of protein kinase CK2 and RAP74 subunit of TFIIIF to protein-coding genes in living cells is DRB sensitive.- *Molec. cell. Biochem* 191: 149-159.
- Ekrem, T. 1999a. Six new *Tanytarsus* species from Ghana, West Africa (Insecta, Diptera, Chironomidae).- *Spixiana* 22: 53-68.
- Ekrem, T. 1999b. *Cladotanytarsus bukavus* (Lehmann, 1979) comb. n. and *C. congolensis* (Lehmann, 1979) comb. n. from Central Africa (Diptera: Chironomidae).- *Annls Limnol.* 35: 185-191.
- Ekrem, T. and Harrison, A. D. 1999a. *Tanytarsus minutipalpus*, spec. nov. from the saline lakes in the Rift Valley, East Africa (Insecta, Diptera, Chironomidae).- *Spixiana* 22: 199-208.
- Ekrem, T. and Reiss, F. 1999a. Two new *Tanytarsus* species (Diptera: Chironomidae) from Brazil, with reduced median volsella.- *Aquat. Insects* 21: 205-213.
- Ekrem, T., Reiss, F. and Langton, P. H. 1999a. *Tanytarsus mancospinosus* sp. n. (Diptera: Chironomidae) from eutrophic lakes in Europe.- *Norw. J. Ent.* 46: 79-87.
- Elberling, H. and Olesen, J. M. 1999a. The structure of a high latitude plant-flower visitor system: the dominance of flies.- *Ecography* 22: 314-323.
- Elke, C., Rauch, P., Spindler-Barth, M. and Spindler, K.-D. 1999a. DNA-binding properties of the ecdysteroid receptor-complex (EcR/USP) of the epithelial cell line from *Chironomus tentans*.- *Archs Insect Biochem Physiol.* 41: 124-133.
- Elser, P. 1999a. Einsatz seminaturlicher Besiedlungskörper zur modellhaften Untersuchung natürlicher Störungen in Fließgewässern.- *Verh. Ges. Ökol.* 29: 457-461.
- Elser, P. 1999b. Use of colonization baskets for the investigation of disturbance phenomena in streams under model conditions.- *Limnologia* 29: 120-127.
- Englund, G. and Evander, D. 1999a. Interactions between sculpins, net-spinning caddis larvae and midge larvae.- *Oikos* 85: 117-126.
- Epler, J. H. and Janetzky, W. J. (1998)1999a. A new species of *Monopelopia* (Diptera: Chironomidae) from phytotelmata in Jamaica, with preliminary ecological notes.- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc.* 71: 216-225.
- Epler, J. H., Harrison, A. D. and Hare, L. (1998)1999a. *Acinoretracus*, a new Afrotropical genus for some species previously placed in *Dicrotendipes* (Diptera: Chironomidae: Chironominae).- *Tijdschr. Ent.* 141: 209-220.
- Fedirko, N. V. i Klevets, M. Yu. 1999a. (Evidence of permanent Ca²⁺ entry in secretory cells of *Chironomus plumosus* salivary gland and its role in basal secretion.)- *Fiziol. Zh.* 45: 84-91.
- Ferrington, L. C. Jr. (1998)1999a. Mary and Jim Sublette. Fifty years of collaborative studies with Chironomidae.- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc.* 71: 195-198.
- Fesl, C., Humpesch, U. H. and Aschauer, A. 1999a. The relationship between habitat structure and biodiversity of the macrozoobenthos in the free-

- flowing section of the Danube in Austria - east of Vienna (preliminary results).- *Arch. Hydrobiol. Supp.* 115: 349-374.
- Fittkau, E. J. and Reiss, F. (1998)1999a. *Micropsectra pharetrophora*, a new species of Tanytarsini (Diptera, Chironomidae) constructing portable larval cases.- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc.* 71: 226-233.
- Flory, E. A. and Milner, A. M. 1999a. Influence of riparian vegetation on invertebrate assemblages in a recently formed stream in Glacier Bay National Park, Alaska.- *J. N. Am. benthol. Soc.* 18: 261-273.
- Fonseca, J. J. L. and Esteves, F. A. 1999a. Influence of bauxite tailings on the structure of the benthic macroinvertebrate community in an Amazonian lake (Lago Batata, Pará - Brazil).- *Revta bras. Biol.* 59: 397-405.
- Forrester, G. E., Dudley, T. L. and Grimm, N. B. 1999a. Trophic interactions in open systems: Effects of predators and nutrients on stream food chain.- *Limnol. Oceanogr.* 44: 1187-1197.
- Francoeur, S. N., Biggs, B. J. F. and Lowe, R. L. 1999a. Inhibition of algae and invertebrates by malathion from insecticide-diffusing substrata.- *J. Freshwat. Ecol.* 14: 179-186.
- Franquet, E. 1999a. Chironomid assemblage of a Lower-Rhone dike field: relationships between substratum and biodiversity.- *Hydrobiologia* 397: 121-131.
- Fretz, A. and Spindler, K.-D. 1999a. Hormonal regulation of actin and tubulin in an epithelial cell line from *Chironomus tentans*.- *Archs Insect Biochem. Physiol.* 41: 71-78.
- Frouz, J. 1999a. Use of soil dwelling Diptera (Insecta, Diptera) as bioindicators: a review of ecological requirements and response to disturbance.- *Agric. Ecosyst. Envir.* 74: 167-186.
- Frouz, J. and Olejník, J. 1999a. Flight patterns of Chironomidae and Dolichopodidae (Diptera) in a suburban fallow area: the effect of vegetation mosaic.- *Biología, Bratisl.* 54: 169-178.
- Galindo, P. A., Lombardero, M., Mur, P., Feo, F., Gómez, E., Borja, J., García, R. and Barber, D. 1999a. Patterns of immunoglobulin E sensitization to chironomids in exposed and unexposed subjects.- *J. invest. Allergol. clin. Immunol.* 9: 117-122.
- García-Berthou, E. 1999a. Food of introduced mosquitofish: ontogenetic diet shift and prey selection.- *J. Fish Biol.* 55: 135-147.
- Genner, M. J., Turner, G. F. and Hawkins, S. J. 1999a. Foraging of rocky habitat cichlid fishes in Lake Malawi: coexistence through niche partitioning?- *Oecologia* 121: 283-292.
- Gennito, D. and Kerans, B. L. 1999a. Effects of a diverse prey assemblage on stonefly feeding.- *J. Freshwat. Ecol.* 14: 219-231.
- Giberson, D. and Hardwick, M. L. 1999a. Pitcher plants (*Sarracenia purpurea*) in eastern Canadian peatlands. Ecology and conservation of the invertebrate inquilines.- In: Batzer, D. P., Rader, R. B. and Wissinger, S. A. (eds.): *Invertebrates in freshwater wetlands of North America: ecology and management*, pp. 401-422. John Wiley & Sons, N. Y., Chichester.
- Gordon, A. K. and Bills, I. R. 1999a. Aspects of the feeding and reproductive biology of the Lake Tanganyikan cichlid, *Lamprologus ornatipinnis* (Pisces, Cichlidae).- *Envir. Biol. Fishes* 55: 431-441.
- Green, A. J., Fox, A. D., Hughes, B. and Hilton, G. M. 1999a. Time-activity budgets and site selection of White-headed Ducks *Oxyura leucocephala* at Burdur Lake, Turkey in late winter.- *Bird Study* 46: 62-73.
- Groenendijk, D. 1999a. *Dynamics of metal adaptation in riverine chironomids*.- Acad. Proefschr., Univ. Amsterdam. 159 pp.
- Groenendijk, D., Kraak, M. H. S. and Admiraal, W. 1999a. Efficient shedding of accumulated metals during metamorphosis in metal-adapted populations of the midge *Chironomus riparius*.- *Envir. Toxic. Chem* 18: 1225-1231.
- Groenendijk, D., Opzeeland, B. van, Dionisio Pires, L. M. and Postma, J. F. 1999a. Fluctuating life-history parameters indicating temporal variability in metal adaptation in riverine chironomids.- *Archs envir. Contam. Toxic.* 37: 175-181.
- Growns, J. E., King, A. J. and Betts, F. M. 1999a. The Snag Bag: a new method for sampling macroinvertebrate communities on large woody debris.- *Hydrobiologia* 405: 67-77.
- Gunderina, L. I. and Kiknadze, I. I. 1999a. Mezvidovaya differentsiatsiya tsitogeneticheskoi struktury vidov-dvoinikov *Chironomus plumosus* L. *Chironomus balatonicus* Devai, Wulker, Scholl (Chironomidae: Diptera).- *Genetika* 35: 1-9. (Interspecific cytogenetic differentiation of the sibling species *Chironomus plumosus* L. and *Chironomus balatonicus* Dévai, Wülker, Scholl (Chironomidae: Diptera).- *Russ. J. Genet.* 35: 1023-1029.)

- Gunderina, L. I., Kiknadze, I. I. and Golygina, V. V. 1999a. Vnutrividovaya differentsiatsiya tsitogeneticheskoi struktury prirodnykh populyatsii *Chironomus plumosus* L. - tsentralnogo vida gruppy vidov-dvoinikov.- *Genetika* 35: 193-202. (Intraspecific differentiation of the cytogenetic structure in natural populations of *Chironomus plumosus* L., the central species in the group of sibling species (Chironomidae: Diptera).- *Russ. J. Genet.* 35: 142-150.)
- Gunderina, L. I., Kiknadze, I. I. and Golygina, V. V. 1999b. Vnutripopulyatsionnaya differentsiatsiya tsitogeneticheskoi struktury u vidov roda *Chironomus* (Chironomidae: Diptera).- *Genetika* 35: 322-328. (Intrapopulation differentiation of cytogenetic structure in species of the genus *Chironomus* (Chironomidae: Diptera).- *Russ. J. Genet.* 35: 256-261.)
- Gunderina, L. I., Kiknadze, I. I. and Golygina, V. V. 1999c. Differentsiatsiya tsitogeneticheskoi struktury prirodnykh populyatsii vidov-dvoinikov gruppi *plumosus Chironomus balatonicus*, *Chironomus entis*, *Chironomus muratensis* i *Chironomus nudiventris* (Chironomidae: Diptera).- *Genetika* 35: 606-614. (Differentiation of the cytogenetic structure of natural populations in the *plumosus* group of sibling species *Chironomus balatonicus*, *Chironomus entis*, *Chironomus muratensis*, and *Chironomus nudiventris* (Chironomidae: Diptera).- *Russ. J. Genet.* 35: 506-513.)
- Haden, G. A., Blinn, D. W., Shannon, J. P. and Wilson, K. P. 1999a. Driftwood: an alternative habitat for macroinvertebrates in a large desert river.- *Hydrobiologia* 397: 179-186.
- Hägele, K. 1999a. Hybrid syndrome-induced postzygotik reproductive isolation: A second reproduction barrier in *Chironomus thummi* (Diptera: Chironomidae).- *J. zool. Syst. evol. Res.* 37: 161-164.
- Hämäläinen, H. 1999a. Critical appraisal of the indexes of chironomid larval deformities and their use in bioindication.- *Annls zool. fenn.* 36: 179-186.
- Hall, R. I., Leavitt, P. R., Quinlan, R., Dixit, A. S. and Smol, J. P. 1999a. Effects of agriculture, urbanization, and climate on water quality in the northern Great Plains.- *Limnol. Oceanogr.* 44: 739-756.
- Halpern, M., Gasith, A., Teltsch, B., Porat, R. and Broza, M. 1999a. Chloramine and copper sulfate as control agents of planktonic larvae of *Chironomus luridus* in water supply systems.- *J. Am. Mosquito Control Ass.* 15: 453-457.
- Hamer, M. J., Goggin, U. M., Muller, K. and Maund, S. J. 1999a. Bioavailability of lambda-cyhalothrin to *Chironomus riparius* in sediment-water and water-only systems.- *Aquat. Ecosyst. Hlth Mgmt* 2: 403-412.
- Harthun, M. 1999a. Der Einfluß des Bibers (*Castor fiber albicus*) auf die Fauna (Odonata, Mollusca, Trichoptera, Ephemeroptera, Diptera) von Mittelgebirgsbächen in Hessen (Deutschland).- *Limnologica* 29: 449-464.
- Hatch, A. C. and Burton, G. A. Jr. 1999a. Photo-induced toxicity of PAHs to *Hyaella azteca* and *Chironomus tentans*: effects of mixtures and behavior.- *Envir. Pollut.* 106: 157-167.
- Haynes, A. 1999a. The long term effect of forest logging on the macroinvertebrates in a Fijian stream.- *Hydrobiologia* 405: 79-87.
- Haynes, J. M., Stewart, T. W. and Cook, G. E. 1999a. Benthic macroinvertebrate communities in southwestern Lake Ontario following invasion of *Dreissena*: continuing changes.- *J. Gt Lakes Res.* 25: 828-838.
- Heinrichs, M. L., Walker, I. R., Mathewes, R. W. and Hebda, R. J. 1999a. Holocene chironomid-inferred salinity and paleovegetation reconstruction from Kilpoola Lake, British Columbia.- *Geogr. phys. quat.* 53: 211-221.
- Hickey, C. W., Golding, L. A., Martin, M. L. and Croker, G. F. 1999a. Chronic toxicity of ammonia to New Zealand freshwater invertebrates: a mesocosm study.- *Archs envir. Contam. Toxic.* 37: 338-351.
- Hirabayashi, K. and Ogawa, K.-i. 1999a. The efficiency of artificial wingbeat sounds for capturing midges in black light traps.- *Ent. exp. applic.* 92: 233-238.
- Hirabayashi, K. and Okino, T. (1998)1999a. Massive flights of chironomid midge nuisance insects around a hypereutrophic lake in Japan: a questionnaire survey of tourists.- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 2: Biology, ecology and natural history of Chironomidae.* - *J. Kans. ent. Soc.* 71: 439-446.
- Hudson, P. L. and Adams, J. V. (1998)1999a. Sieve efficiency in benthic sampling as related to chironomid head capsule width.- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc.* 71: 456-468.
- Huggett, D. B., Gillespie, W. B. Jr. and Rodgers, J. H. Jr. 1999a. Copper bioavailability in Steilacoom Lake sediments.- *Archs envir. Contam. Toxic.* 36: 120-123.

- Iannacone, J. A. y Dale, W. E. 1999a. Protocolo de bioensayo ecotoxicológico para evaluar metales pesados contaminantes de agua dulce con *Chironomus calligraphus* (Diptera: Chironomidae) y *Moina macrocopa* (Crustacea: Cladocera), en el Río Rimac, Lima, Peru.- *Revta per. Ent. 41*: 111-120.
- Iannacone Oliver, J., Alvaríño Flores, L. y Gutierrez Roman, A. 1999a. Cinco ensayos ecotoxicológicos para evaluar metales pesados en el agua dulce.- *Boln Soc. quim. Peru 65*: 30-45.
- Il'inskaya, N. B., Petrova, N. A. and Matena, I. 1999a. The relationship between the level of inversion polymorphism and the type of water body, the season, and the year of observation in *Chironomus plumosus* L. (Diptera, Chironomidae).- *Russ. J. Genet. 35*: 908-917.
- Imbert, J. B. and Perry, J. A. 1999a. Invertebrate responses to stepwise and abrupt increases in non-scouring flow: The role of refugia.- *Arch. Hydrobiol. 146*: 167-187.
- Istomina, A. G., Kiknadze, I. I. i Siirin, M. T. 1999a. Karyologicheskii analiz vidov *Chironomus* gruppy *obtusidens* Altaya (Diptera: Chironomidae). (Karyological analysis of *Chironomus* gr. *obtusidens* from the Altai (Diptera, Chironomidae).)- *Tsitologiya 41*: 1022-1031.
- Itkonen, A., Marttila, V., Meriläinen, J. J. and Salonen, V.-P. 1999a. 8000-year history of palaeoproductivity in a large boreal lake.- *J. Paleolimnol. 21*: 271-294.
- Jacobsen, R. E. (1998)1999a. The symbiotic relationship of a chironomid with its ephemeropteran host in an Arizona mountain stream.- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 2: Biology, ecology and natural history of Chironomidae.* - *J. Kans. ent. Soc. 71*: 426-438.
- Johnson, A. A. and Kleve, M. G. 1999a. *Hydromermis contorta* (Kohn) and *Hydromermis pseudocontorta* n. sp. from chironomids of Lake Itasca and Long Lake, Minnesota.- *J. Parasit. 85*: 105-109.
- Johnston, T. A. and Cunjak, R. A. 1999a. Dry mass-length relationships for benthic insects: a review with new data from Catamaran Brook, New Brunswick, Canada.- *Freshwat. Biol. 41*: 653-674.
- Jorcín, A. 1999a. Temporal and spatial variability in the macrozoobenthic community along a salinity gradient in the Castillos Lagoon (Uruguay).- *Arch. Hydrobiol. 146*: 369-384.
- Jordan, S., Shiozawa, D. K. and Schmid-Arraya, J. M. 1999a. Benthic invertebrates of a large, sandy river system: The Green and Colorado Rivers of Canyonlands National Park, Utah.- *Arch. Hydrobiol. 147*: 91-127.
- Kadono, A., Takemon, Y. and Tokeshi, M. 1999a. Mobility and habitat colonization in stream invertebrates: An experimental study.- *Jap. J. Limnol. 60*: 215-222.
- Kajan, R. and Frenzel, P. 1999a. The effect of chironomid larvae on production, oxidation and fluxes of methane in a flooded rice soil.- *FEMS Microbiol. Ecol. 28*: 121-129.
- Kangur, K., Kangur, A. and Kangur, P. 1999a. A comparative study on the feeding of eel, *Anguilla anguilla* (L.), bream, *Abramis brama* (L.) and ruffe, *Gymnocephalus cernuus* (L.) in Lake Võrtsjärv, Estonia.- *Hydrobiologia 408/409*: 65-72.
- Karjalainen, J., Leppä, M., Rahkola, M. and Tolonen, K. 1999a. The role of benthivorous and planktivorous fish in a mesotrophic lake ecosystem.- *Hydrobiologia 408/409*: 73-84.
- Karouna-Renier, N. and Zehr, J. 1999a. Ecological implications of molecular biomarkers: assaying sub-lethal stress in the midge *Chironomus tentans* using heat shock protein 70 (HSP-70) expression.- *Hydrobiologia 401*: 255-264.
- Kawai, K., Inoue, E. and Imabayashi, H. 1999a. Differences in occurrence in relation to the eutrophication level among chironomid species of the genus, *Tanytarsus*, in littorals of lakes.- *Med. Ent. Zool. 50*: 65-70.
- Kay, W. R., Smith, M. J., Pinder, A. M., Mcrae, J. M., Davis, J. A. and Halse, S. A. 1999a. Patterns of distribution of macroinvertebrate families in rivers of north-western Australia.- *Freshwat. Biol. 41*: 299-316.
- Kedwards, T. J., Maund, S. J. and Chapman, P. F. 1999a. Community level analysis of ecotoxicological field studies: I. Biological monitoring.- *Envir. Toxic. Chem 18*: 149-157.
- Kemble, N. E., Dwyer, F. J., Ingersoll, C. G., Dawson, T. D. and Norberg-King, T. J. 1999a. Tolerance of freshwater test organisms to formulated sediments for use as control materials in whole-sediment toxicity test.- *Envir. Toxic. Chem 18*: 222-230.
- Khaitov, V. M., Fokin, M. F. and Nicolaeva, A. M. 1999a. Structure of communities associated with dense assemblages of the tube-dwelling polychaete *Polydora quadrilobata* Jacobi (Spionidae) in the White Sea.- *Hydrobiologia 393*: 221-226.

- King, R. S. and Brazner, J. C. 1999a. Coastal wetland insect communities along a trophic gradient in Green Bay, Lake Michigan.- *Wetlands* 19: 426-437.
- Kobayashi, T. and Kuranishi, R. 1999a. The second species in the subfamily Podonominae recorded from Japan, *Papaboreochlus* [sic!] *okinawanus*, new species (Diptera: Chironomidae).- *Raffles Bull. Zool.* 47: 601-606.
- Kobayashi, T. and Sæther, O. A. 1999a. *Sasacricotopus*, a new orthoclad genus from Japan (Diptera, Chironomidae).- *Acta zool. Acad. Scient. hung.* 45: 155-159.
- Kobayashi, T. and Suzuki, H. 1999a. *Harnischia ohmuraensis* sp. nov. and the first record of *Parachironomus monochromus* (van der Wulp, 1874) from Japan (Diptera: Chironomidae).- *Med. Ent. Zool.* 50: 79-84.
- Kobayashi, T. and Suzuki, H. 1999b. The first record of the genus *Apometriocnemus* Sæther, 1985: *A. japonicus* sp. n. from the Palaeartic Region (Diptera: Chironomidae).- *Tijdschr. Ent.* 142: 65-67.
- Kondo, S. (1998)1999a. Seasonal abundances of two halophilous chironomids (Diptera: Chironomidae) in a brackish pond of Nagoya City, Japan.- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc.* 71: 469-473.
- Kovachev, S. and Stoichev, S. 1999a. Hydrofaunistic investigations of the Karla Lake, Central Greece.- *Lauterbornia* 36: 71-73.
- Kovachev, S., Stoichev, S. and Hainadjieva, V. 1999a. The zoobenthos of several lakes along the Northern Bulgarian Black Sea Coast.- *Lauterbornia* 35: 33-38.
- Kovachev, S., Stoichev, S., Uzunov, Y., Kumanski, K. and Memeti, A. 1999a. Hydrofaunistic study of the Tetovska River, Macedonia.- *Lauterbornia* 36: 67-70.
- Kreivi, P., Muotka, T., Huusko, A., Mäki-Petäys, A., Huhta, A. and Meissner, K. 1999a. Diel feeding periodicity, daily ration and prey selectivity in juvenile brown trout in a subarctic river.- *J. Fish Biol.* 55: 553-571.
- Kugler, N., Bogner, F. X. und Kinzle, W. 1999a. Einbindung einer Dorfbachrenaturierung in den Biologieunterricht.- *Verh. Ges. Ökol.* 29: 629-621.
- Kuhns, L. A. and Berg, M. B. 1999a. Benthic invertebrate community responses to round goby (*Neogobius melanostomus*) and zebra mussel (*Dreissena polymorpha*) invasion in southern Lake Michigan.- *J. Gt Lakes Res.* 25: 910-917.
- Lacey, R., Watzin, M. C. and McIntosh, A. W. 1999a. Sediment organic matter content as a confounding factor in toxicity tests with *Chironomus tentans*.- *Envir. Toxic. Chem.* 18: 231-236.
- Lammers-Campbell, R. (1998)1999a. Ordination of chironomid (Diptera: Chironomidae) communities characterizing habitats in a Minnesota peatland.- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 2: Biology, ecology and natural history of Chironomidae.* - *J. Kans. ent. Soc.* 71: 414-425.
- Landrum, P. F., Fisher, S. W., Hwang, H. and Hickey, J. 1999a. Hazard evaluation of ten organophosphorus insecticides against the midge, *Chironomus riparius* via QSAR.- *SAR QSAR envir. Res.* 10: 423-450.
- Lang, C. 1999a. Contrasting responses of oligochaetes (Annelida) and chironomids (Diptera) to the abatement of eutrophication in Lake Neuchâtel.- *Aquat. Sci.* 61: 206-214.
- Langton, P. H. 1999(1998)a. *Micropsectra silvesterae* n. sp. and *Tanytarsus heliomesonyctios* n. sp., (Diptera: Chironomidae), two parthenogenetic species from Ellesmere Island, Arctic Canada.- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc.* 71: 208-215.
- Langton, P. H. and Casas, J. 1999a. Changes in chironomid assemblage composition in two Mediterranean mountain streams over a period of extreme hydrological condition.- *Hydrobiologia* 390: 37-49.
- Laskowski, H. P., Larsen, A. C., O'Shea, G. F. and Pittendrigh, B. R. 1999a. Effect of xanthan gum and traditional mosquito larvicides on chironomid larvae.- *Wildl. Soc. Bull.* 27: 741-745.
- Lazaridou-Dimitriadou, M., Artemiadou, V., Yfantis, G., Mourelatos, S. and Mylopoulos, Y. 1999a. Contribution to the ecological quality of Aliakmon river (Macedonia, Greece): a multivariate approach.- *Hydrobiologia* 410: 47-58.
- Leal, J. J. F. and Esteves, F. A., 1999a. Density and biomass of *Campsurus* sp. (Ephemeroptera) and other macroinvertebrates in an Amazonian lake impacted by bauxite tailings (Lago Batata, Pará, Brazil).- *Amazoniana* 15: 193-209.
- Leslie, H. A., Pavluk, T. I., bij de Vaate, A. and Kraak, M. H. S. 1999a. Triad assessment of the impact of chromium contamination on benthic macroinvertebrates in the Chusovaya River

- (Urals, Russia).- *Archs envir. Contam. Toxic.* 37: 182-189.
- Lichtwardt, R. W. and Williams, M. C. 1999a. Three Harpellales that live in one species of aquatic chironomid larva.- *Mycologia* 91: 396-399.
- Lichtwardt, R. W., Ferrington, L. C. and Lastra, C. L. 1999a. Trichomycetes in Argentinean aquatic insect larvae.- *Mycologia* 91: 1060-1082.
- Linz, G. M., Bleier, W. J., Overland, J. D. and Homan, H. J. 1999a. Response of invertebrates to glyphosate-induced habitat alterations in wetlands.- *Wetlands* 19: 220-227.
- Livingstone, D. M., Lotter, A. F., Walker, I. R. 1999a. The decrease in summer surface water temperature with altitude in Swiss Alpine lakes: A comparison with air temperature lapse rates.- *Arct. Antarct. Alp. Res.* 31: 341-352.
- López, C. C., Kannert, I., Scherbik, S. V. and Edström, J.-E. 1999a. Interspersed DNA element restricted to a specific group of telomeres in the dipteran *Chironomus pallidivittatus*.- *Gene* 233: 249-259.
- López, C. C., Rodriguez, E., Díez, J. L., Edström, J.-E. and Morcillo, G. 1999a. Histochemical localization of reverse transcriptase in polytene chromosomes of chironomids.- *Chromosoma* 108: 302-307.
- Lorenz, M., Hillisch, A., Payet, D., Buttinelli, M., Travers, A. and Diekmann, S. 1999a. DNA bending induced by high mobility group proteins studied by fluorescence resonance energy transfer.- *Biochemistry* 38: 12150-12158.
- Lydy, M. J., Belden, J. B. and Ternes, M. A. 1999a. Effects of temperature on the toxicity of m-parathion, chlorpyrifos, and pentachlorobenzene to *Chironomus tentans*.- *Archs envir. Contam. Toxic.* 37: 542-547.
- Maheshwari, G. and Maheshwari, G. 1999a. Biosystematic studies of Indian Chironomidae (Diptera).- *J. Bombay nat. Hist. Soc.* 96: 74-81.
- Makarchenko, E. A. and Zorina, O. V. 1999a. A preliminary list of Chironomidae (Diptera) of the Primorye Territory (Russian Far East).- *Far east. Ent.* 78: 1-15.
- Malloy, J. C., Meade, M. L. and Olsen, E. W. 1999a. Small-scale spatial variation of selenium concentrations in chironomid larvae.- *Bull. envir. Contam. Toxic.* 62: 122-129.
- Mangum, F. A. and Madrigal, J. L. 1999a. Rotenone effects on aquatic macroinvertebrates of the Strawberry River, Utah: a five-year survey.- *J. Freshwat. Ecol.* 14: 125-135.
- Marchant, R., Hirst, A., Norris, R. and Metzeling, L. 1999a. Classification of macroinvertebrate communities across drainage basins in Victoria, Australia: consequences of sampling on a broad spatial scale for predictive modelling.- *Freshwat. Biol.* 41: 253-268.
- Martin, J. (1998)1999a. *Chironomus forsythi* n. sp. from New Zealand, a member of the *C. zealandicus*-group with salinarius-type larvae.- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc.* 71: 247-259.
- Massaferro, M. M., Massaferro, J., Roman Ross, G., Amos, A. J. and Lami, A. 1999a. Late Pleistocene and early Holocene ecological response of Lake El Trébol (Patagonia, Argentina) to environmental changes.- *J. Paleolimnol.* 22: 137-148.
- Mastrantuono, L. and Mancinelli, T. 1999a. Long-term changes of zoobenthic fauna and submerged vegetation in the shallow Lake Monterosi (Italy).- *Limnologica* 29: 160-167.
- Mayle, F. E., Bell, M., Birks, H. H., Brooks, S. J., Coope, G. R., Lowe, J. J., Sheldrick, C., Shijie, L., Turney, C. S. M. and Walker, M. J. C. 1999a. Climate variations in Britain during the last Glacial-Holocene transition (15.0-11.5 cal ka BP): Comparison with the GRIP ice-core record.- *J. geol. Soc.* 156: 411-423.
- McAllen, R. 1999a. *Enteromorpha intestinalis* - a refuge for the supralittoral rock-pool harpacticoid copepod *Tigriopus brevicornis*.- *J. mar. biol. Ass. U. K.* 79: 1125-1126.
- McKinney, T., Ayers, A. D. and Rogers, R. S. 1999a. Macroinvertebrate drift in the tailwater of a regulated river below Glen Canyon Dam, Arizona.- *SWest Nat.* 44: 205-210.
- McLachlan, A. 1999a. Parasites promote mating success: the case of a midge and a mite.- *Anim. Behav.* 57: 1199-1205.
- McLachlan, A., Ladle, R. and Bleay, C. 1999a. Is infestation the result of adaptive choice behaviour by the parasite? A study of mites and midges.- *Anim. Behav.* 58: 615-620.
- McNamara, A. E. and Hill, W. R. 1999a. Effects of UV-B dose and irradiance: comparison among grazers.- *J. N. Am. benthol. Soc.* 18: 370-380.
- Mera, A., Cobo, F. y González, M. A. 1999a. Valor nutritivo y energético de algunas familias de macroinvertebrados dulceacuícolas pertenecientes a la base trófica del Salmón Atlántico (*Salmo salar* L., 1758) en estadios juveniles.- In: *1ª Semana del Salmón Atlántico en la Península Ibérica*, pp. 155-162. Univ. Oviedo.

- Mercier, V., Vis, C., Morin, A., and Hudon, C. 1999a. Patterns of invertebrate and periphyton size distributions from navigation buoys in the St. Lawrence River.- *Hydrobiologia* 394: 83-91.
- Messias, M. C. and Oliveira, S. J. de (1998)1999a. On a new species of the genus *Oukuriella* Epler (Diptera, Chironomidae, Chironominae).- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc.* 71: 260-262.
- Michiels, S. 1999a. Die Chironomidae (Diptera) der unteren Salzach.- *Lauterbornia* 36: 45-53.
- Minakawa, N. and Gara, R. I. 1999a. Ecological effects of a chum salmon (*Oncorhynchus keta*) spawning run in a small stream of the Pacific Northwest.- *J. Freshwat. Ecol.* 14: 327-335.
- Minakawa, N. and Kraft, G. F. 1999a. Fall and winter diets of juvenile coho salmon in a small stream and an adjacent pond in Washington State.- *J. Freshwat. Ecol.* 14: 249-254.
- Miota, F., Siegfried, B. D., Scharf, M. E. and Lydy, M. J. 1999a. Atrazine induction of cytochrome P450 in *Chironomus tentans* larvae.- *Chemosphere* 40: 285-291.
- Misra, J. K., White, M. M. and Lichtwardt, R. W. 1999a. *Furculomyces septentrionalis* reveals an unexpected distribution for this genus of Harpellales.- *Mycologia* 91: 703-706.
- Mogi, M., Sunahara, T. and Selomo, M. 1999a. Mosquito and aquatic predator communities in ground pools on lands deforested for rice field development in Central Sulawesi, Indonesia.- *J. Am. Mosquito Control Ass.* 15: 92-97.
- Mori, C., Orsini, A. and Migon, C. 1999a. Impact of arsenic and antimony contamination on benthic invertebrates in a minor Corsican river.- *Hydrobiologia* 392: 73-80.
- Moubayed, J. and Langton, P. H. 1999a. Description of *Parametricnemus valescurensis* sp. n. from temporary streams and helocrenes in southern France (Diptera, Chironomidae).- *Nouv. Revue Ent. N. S.* 16: 155-160.
- Moyo, N. and Fernando, C. H. 1999a. The food of *Sargochromis codringtonii* (Boulenger, 1908) in Lake Kariba, Zimbabwe.- *Hydrobiologia* 411: 129-137.
- Nakano, S., Fausch, K. D. and Kiatno, S. 1999a. Flexible niche partitioning via a foraging mode shift: a proposed mechanism for coexistence in stream-dwelling charrs.- *J. Anim. Ecol.* 68: 1079-1092.
- Nakano, S., Miyasaka, H. and Kuhara, N. 1999a. Terrestrial-aquatic linkages: riparian arthropod inputs alter trophic cascades in a stream food web.- *Ecology* 80: 2435-2441.
- Nelson, S. M. and Roline, R. A. 1999a. Relationships between metals and hyporheic invertebrate community structure in a river recovering from metals contamination.- *Hydrobiologia* 397: 211-226.
- Nessimian, J. L., Sanseverino, A. M. e de Oliveira, A. L. H. 1999a. (Trophic relationships of Chironomidae larvae (Diptera) and its importance on the foodwebs in a sand dune marsh on the littoral of Rio de Janeiro State.)- *Revta bras. Ent.* 43: 47-53.
- Nielsen, D. L., Hillman, T. J. and Smith, F. J. 1999a. Effects of hydrological variation and planktivorous competition on macroinvertebrate community structure in experimental billabongs.- *Freshwat. Biol.* 42: 427-444.
- Northcote, T. G. and Chapman, M. A. 1999a. Dietary alterations in resident and migratory New Zealand common smelt (*Retropinna retropinna*) in lower Waikato lakes after two decades of habitat change.- *N. Z. Jl mar. Freshwat. Res.* 33: 425-436.
- Ó Hare, M. T. and Murphy, K. J. 1999a. Invertebrate hydraulic microhabitat and community structure in *Callitriche stagnalis* Scop. patches.- *Hydrobiologia* 415: 169-176.
- Olander, H., Birks, H. J. B., Korhola, A. and Blom, T. 1999a. An expanded calibration model for inferring lakewater and air temperatures from fossil chironomid assemblages in northern Fennoscandia.- *Holocene* 9: 279-294.
- Orendt, C. 1999a. Chironomids as bioindicators in acidified streams: a contribution to the acidity tolerance of chironomid species with a classification in sensitivity classes.- *Int. Rev. Hydrobiol.* 84: 439-449.
- Otto, C.-J. and Schiegg, K. 1999a. Chironomidae (Diptera) collected in the forest reserve Sihlwald ZH, with 21 new records for Switzerland.- *Mitt. schweiz. ent. Ges.* 72: 95-103.
- Paggi, A. C. 1999a. (Chironomidae as quality indicators of freshwater environments.)- *Revta Soc. ent. argent.* 58: 202-207.
- Peglow, H.-G. 1999a. Bachstelzen fangen rennende Zuckmücken.- *Orn. Mitt.* 51: 94-95.
- Pemberton, R. W. 1999a. Natural enemies of *Trapa* spp. in Northeast Asia and Europe.- *Biol. Control.* 14: 168-180.
- Penczak, T. 1999a. Fish production and food consumption in the Warta River (Poland): continued post-impoundment study (1990-1994).- *Hydrobiologia* 416: 107-123.

- Penczak, T., Agostinho, A. A., Hahn, N. S., Fugi, R. and Gomes, L. C. 1999a. Energy budgets of fish populations in two tributaries of the Paraná River, Paraná, Brazil.- *J. trop. Ecol.* 15: 159-177.
- Pereira de Araujo, P. R. 1999a. A progress report on the performance of two artificial substrate for collecting Chironomidae larvae (Insecta: Diptera) in Guandu River, Rio de Janeiro State, Brazil.- *Ent. Vectores* 6: 306-313.
- Persson, A. and Hansson, L.-A. 1999a. Diet shift in fish following competitive release.- *Can. J. Fish. aquat. Sci.* 56: 70-78.
- Persson, L., Byström, P., Wahlström, E., Andersson, J. and Hjelm, J. 1999a. Interactions among size-structured populations in a whole-lake experiment: size- and scale-dependent processes.- *Oikos* 87: 139-156.
- Petranka, J. W. and Kennedy, C. A. 1999a. Pond tadpoles with generalized morphology: is it time to reconsider their functional roles in aquatic communities?- *Oecologia* 120: 621-631.
- Piecy_ska, E., Ko_odziejczyk, A and Rybak, J. I. 1999a. The responses of littoral invertebrates to eutrophication-linked changes in plant communities.- *Hydrobiologia* 391: 9-21.
- Pöpperl, R. 1999a. Emergence pattern of Diptera in various sections of a northern German lowland stream.- *Limnologia* 29: 128-136.
- Pont, D., Franquet, E. and Tourenq, J. N. 1999a. Impact of different *Bacillus thuringiensis* variety *israelensis* treatments on a chironomid (Diptera Chironomidae) community in a temporary marsh.- *J. econ. Ent.* 92: 266-272.
- Pope, R. J., Gordon, A. M. and Kaushik, N. K. 1999a. Leaf litter colonization by invertebrates in the littoral zone of a small oligotrophic lake.- *Hydrobiologia* 392: 99-112.
- Post, J. R., Parkinson, E. A. and Johnston, N. T. 1999a. Density-dependent processes in structured fish populations: interaction strengths in whole-lake experiments.- *Ecol. Monogr.* 69: 155-175.
- Postma, J. F. and Groenendijk, D. 1999a. Adaptation to metals in the midge *Chironomus riparius*: a case study in the River Dommel.- In: Forbes, V. E. (ed.): *Genetics and ecotoxicology*, pp. 79-101. Taylor & Francis, Lond.
- Prat, N., Toja, J., Sola, C., Burgos, M. D., Plans, M. and Rieradevall, M. 1999a. Effect of dumping and cleaning activities on the aquatic ecosystems of the Guadiamar River following a toxic flood.- *Sci. tot. Envir.* 242: 231-248.
- Prus, T., Prus, M. and Bijok, P. 1999a. Diversity of invertebrate fauna in littoral of shallow Myczkowce dam reservoir in comparison with a deep Solina dam reservoir.- *Hydrobiologia* 409: 203-210.
- Rader, R. B. and Belish, T. A. 1999a. Influence of mild to severe flow alterations on invertebrates in three mountain streams.- *Regul. Rivers Res. Mgmt* 15: 353-363.
- Rawer-Jost, C., Kappus, B., Böhmer, J., Jansen, W. and Rahmann, H. 1999a. Upstream movements of benthic macroinvertebrates in two different types of fishways in southwestern Germany.- *Hydrobiologia* 391: 47-61.
- Reinhold, J. O., Hendriks, A. J., Slager, L. K. and Ohm, M. 1999a. Transfer of microcontaminants from sediment to chironomids, and the risk for the Pond bat *Myotis dasycneme* (Chiroptera) preying on them.- *Aquat. Ecol.* 33: 363-376.
- Rempel, L. L., Richardson, J. S. and Healey, M. C. 1999a. Flow refugia for benthic macroinvertebrates during flooding of a large river.- *J. N. Am. benthol. Soc.* 18: 34-48.
- Rey, D., Pautou, M.-P. and Meyran, J.-C. 1999a. Histopathological effects of tannic acid on the midgut epithelium of some aquatic Diptera larvae.- *J. Invert. Path.* 73: 173-181.
- Ribeiro, R., Kelly, L. A., Gonçalves, F., Burton, G. A. Jr. and Soares, A. M. V. M. 1999a. New artificial sediment for *Chironomus riparius* toxicity testing.- *Bull. envir. Contam. Toxic.* 63: 691-697.
- Richardson, B. A. 1999a. The bromeliad microcosm and the assessment of faunal diversity in a Neotropical forest.- *Biotropica* 31: 321-336.
- Ristola, T., Kukkonen, J. V. K. and Pellinen, J. 1999a. Body residues and responses of the midge *Chironomus riparius* to sediment-associated 2,4,5-trichlorophenol in subchronic and chronic exposures.- *Archs envir. Contam. Toxic.* 37: 42-49.
- Ristola, T., Pellinen, J., Ruokolainen, M., Kostamo, A. and Kukkonen, J. V. K. 1999a. Effect of sediment type, feeding level, and larval density on growth and development of a midge (*Chironomus riparius*).- *Envir. Toxic. Chemy* 18: 756-764.
- Robarts, R. D., Zhulidov, A. V., Zhulidova, O. V., Pavlov, D. F., Headley, J. V., Reznikov, S. A., Matveev, A. A. and Lysenko, V. S. 1999a. Biogeography and limnology of the Lake Taimyr-wetland system, Russian Arctic: An ecological synthesis.- *Arch. Hydrobiol. Suppl.* 121: 159-200.
- Robinson, C. T. and Burgherr, P. 1999a. Seasonal disturbance of a lake outlet benthic community.- *Arch. Hydrobiol.* 145: 297-315.

- Robson, B. J., Chester, E. T. and Davis, J. A. 1999a. Manipulating the intensity of near-bed turbulence in rivers: effects on benthic invertebrates.- *Freshwat. Biol.* 42: 645-653.
- Roy, I. and Hare, L. 1999a. Relative importance of water and food as cadmium sources to the predatory insect *Sialis velata* (Megaloptera).- *Can. J. Fish. aquat. Sci.* 56: 1143-1149.
- Särkkä, J., Mäkelä, J. 1999a. Meiofauna of esker groundwaters in Finland.- *Hydrobiologia* 405: 25-37.
- Sæther, O. A. and Ekrem, T. 1999a. *Mollerella*, a new terrestrial orthoclad genus from the Netherlands (Diptera: Chironomidae).- *Acta zool. Acad. Scient. hung.* 45: 161-168.
- Sæther, O. A. and Sundal, A. (1998)1999a. *Cerobregma*, a new subgenus of *Polypedilum* Kieffer, with a tentative phylogeny of subgenera and species groups within *Polypedilum* (Diptera: Chironomidae).- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc.* 71: 315-382.
- Shahin, Y. and Arslan, N. P. 1999a. Epoche in Chironomidae (Diptera) larvae in the streams Beşik and Çamlıca, a part of the southern Sakarya river system.- *Turk. J. Zool.* 23: 265-267.
- Samietz, R. 1999a. Chironomidae = Tendipedidae.- In: Schumann, H., Bährmann, R. und Stark, A. (eds.): *Entomofauna Germanica 2: Checkliste der Dipteren Deutschlands. Studia dipterol. Suppl.* 2: 39-50. Ampyx-Verl., Halle.
- Sanders, M. D. 1999a. Effect of changes in water level on numbers of black stilts (*Himantopus novaezelandiae*) using deltas of Lake Benmore.- *N. Z. J. Zool.* 26: 155-163.
- Sasa, M. and Ogata, K. 1999a. Taxonomic studies on the chironomid midges (Diptera, Chironomidae) collected from the Kurobe Municipal Sewage Treatment Plant.- *Med. Ent. Zool.* 50: 85-104.
- Sasa, M. and Suzuki, H. 1999a. Studies on the chironomid midges of Tsushima and Iki Islands, western Japan: Part 1. Species of Chironominae collected on Tsushima.- *Trop. Med.* 41: 1-53.
- Sayer, C., Roberts, N., Sadler, J., David, C. and Wade, P. M. 1999a. Biodiversity changes in a shallow lake ecosystem: a multi-proxy palaeolimnological analysis.- *J. Biogeogr.* 26: 97-114.
- Schmid, U. 1999a. Das Makrozoobenthos des Unteren Odertals - Faunenzusammensetzung und Besiedlungsdynamik in einer Flußbaue.- In: Dohle, W., Bornkamm, R. und Weigmann, G. (eds.): *Das Untere Odertal. Auswirkungen der periodischen Überschwemmungen auf Biozönosen und Arten. Limnologie aktuell* 9: 317-336. E. Schweizerbart'sche Verlagsbuchh., Stuttgart.
- Schmude, K. L., Liber, K., Corry, T. D. and Stay, F. S. 1999a. Effects of 4-nonylphenol on benthic macroinvertebrates and insect emergence in littoral enclosures.- *Envir. Toxic. Chem* 18: 386-393.
- Schnabel, S. 1999a. *Faunistisch-ökologische Untersuchung der Chironomidae (Diptera: Nematocera) temporärer Tümpel in der Lahnaue bei Marburg.*- Dipl.-Arb., Univ. Marburg. 221 pp.
- Schönbauer, B. 1999a. Spatio-temporal patterns of macrobenthic invertebrates in a free-flowing section of the River Danube in Austria.- *Arch. Hydrobiol. Suppl.* 115: 375-397.
- Sergeeva, I. V. and Makarchenko, E. A. 1999a. Morphokaryological description of *Derotanypus sibiricus* (Kruglova et Tshernovskij, 1940) (Diptera: Chironomidae, Tanypodinae) from Siberia.- *Ent. Sci.* 2: 359-365.
- Servia, M. J., Cobo, F. y González, M. A. 1999a. Sobre la posible repercusión de la presencia de deformidades en el ciclo vital de *Chironomus riparius* Meigen, 1804 (Diptera: Chironomidae).- *Boln Asoc. esp. Ent.* 23: 105-113.
- Servia, M. J., Cobo, F. y González, M. A. 1999b. Aparición de deformidades en larvas del género *Chironomus* (Diptera, Chironomidae) recolectadas en ambientes no alterados.- *Boln Asoc. esp. Ent.* 23: 331-332.
- Shieh, S.-H. and Wang, P.-S. 1999a. Colonization patterns of aquatic insects on artificial substrates in a Taiwan stream.- *Zhonghua Kunchong* (= *Chin. J. Ent.*) 19: 27-50.
- Shieh, S.-H., Kondratieff, B. C., Ward, J. V. and Rice, D. A. 1999a. The relationship of macroinvertebrate assemblages to water chemistry in a polluted Colorado plains stream.- *Arch. Hydrobiol.* 145: 405-432.
- Shishehchian, F., Yusoff, F. M., Omar, H. and Kamarudin, M. S. 1999a. Nitrogenous excretion of *Penaeus monodon* postlarvae fed with different diets.- *Mar. Pollut. Bull.* 39: 224-227.
- Shobanov, N. A. 1999a. Kariotip *Chironomus fraternus* Wuelker (Diptera, Chironomidae) iz basseina Rybinskogo vodokhranilishcha (The karyotype of *Chironomus fraternus* Wuelker from the basin of Rybinsk Reservoir).- *Tsitologiya* 41: 641-646.

- Shobanov, N. A., Kiknadze, I. I. and Butler, M. G. 1999a. Palearctic and Nearctic *Chironomus (Camptochironomus) tentans* (Fabricius) are different species (Diptera: Chironomidae).- *Ent. scand.* 30: 311-322.
- Sibley, P. K., Benoit, D. A., Balcer, M. D., Phipps, G. L., West, C. W., Hoke, R. A. and Ankley, G. T. 1999a. In situ bioassay chamber for assessment of sediment toxicity and bioaccumulation using benthic invertebrates.- *Envir. Toxic. Chem* 18: 2325-2336.
- Simi, V. and Simi, S. 1999a. Use of the river macrozoobenthos of Serbia to formulate a biotic index.- *Hydrobiologia* 416: 51-64.
- Singh, O. P., Björkroth, B., Masich, S., Wieslander, L. and Daneholt, B. 1999a. The intranuclear movement of Balbiani ring pre-messenger ribonucleoprotein particles.- *Expl Cell Res.* 251: 135-146.
- Slaymaker, A. K., Ferrington, L. C. Jr. and Lichtwardt, R. W. (1998)1999a. Chironomidae-Trichomycete associations: a literature review.- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc.* 71: 490-500.
- Smart, A. C. and Harper, D. M. 1999a. Life after lakes: the ecology and management of the water distribution network.- *Hydrobiologia* 395/396: 379-386.
- Snyder, D. J. and Peterson, M. S. 1999a. Foraging and prey selection by bluespotted sunfish *Enneacanthus gloriosus* (Holbrook) in backwater vegetated ponds in coastal Mississippi.- *J. Freshwat. Ecol.* 14: 187-196.
- Soong, K., Chen, G. F. and Cao, J. R. 1999a. Life history studies of the flightless marine midges *Pontomyia* spp. (Diptera: Chironomidae).- *Zool. Stud.* 38: 466-473.
- Sozen, M. and Yigit, S. 1999a. (The benthic fauna and some limnological aspects of Lake Akşehir (Konya).)- *Turk. J. Zool.* 23 (Suppl. 3): 829-847.
- Spencer, M., Blaustein, L., Schwartz, S. S. and Cohen, J. E. 1999a. Species richness and the proportion of predatory animal species in temporary pools: relationships with habitat size and permanence.- *Ecol. Lett.* 2: 157-166.
- Spies, M. (1998)1999a. *Cricotopus (Isocladius) subletteorum*, a new species of Chironomidae (Diptera) from the southwestern United States.- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc.* 71: 199-207.
- Spindler, K.-D. and Spindler-Barth, M. 1999a. Inhibitors of chitinases.- In: Jollès, P. and Muzzarelli, R.A.A. (eds.): *Chitin and chitinases*, pp. 201-209. Birkhäuser, Basel.
- Stahl, J. B. (1998)1999a. Chironomid extirpation at Crooked Lake, Indiana.- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 2: Biology, ecology and natural history of Chironomidae.* - *J. Kans. ent. Soc.* 71: 383-387.
- Stanko-Mishic, S., Cooper, J. K. and Silver, P. 1999a. Manipulation of habitat quality: effects on chironomid life history traits.- *Freshwat. Biol.* 41: 719-727.
- Steinhart, M. 1999a. Die Chironomiden des Unteren Odertals - Untersuchung möglicher Adaptationen an das Überflutungsgeschehen.- In: Dohle, W., Bornkamm, R. und Weigmann, G. (eds.): *Das Untere Odertal. Auswirkungen der periodischen Überschwemmungen auf Biozöosen und Arten.* *Limnologie aktuell* 9: 337-351. E. Schweizerbart'sche Verlagsbuchh., Stuttgart.
- Stelzer, R. S. and Lamberti, G. A. 1999a. Independent and interactive effects of crayfish and darters on a stream benthic community.- *J. N. Am. benthol. Soc.* 18: 524-532.
- Stevens, M. M., Fox, K. M., Coombes, N. E. and Lewin, L. A. 1999a. Effect of fipronil seed treatments on the germination and early growth of rice.- *Pestic. Sci.* 55: 517-523.
- Stewart, T. W., Miner, J. G. and Lowe, R. L. 1999a. A field experiment to determine *Dreissena* and predator effects on zoobenthos in a nearshore, rocky habitat of western Lake Erie.- *J. N. Am. benthol. Soc.* 18: 488-498.
- Stuijffzand, S. C., Jonker, M. J., Ammelrooy, E. van and Admiraal, W. 1999a. Species-specific responses to metals in organically enriched river water, with emphasis on effects of humic acids.- *Envir. Pollut.* 106: 115-121.
- Sturm, A. and Hansen, P.-D. 1999a. Altered cholinesterase and monooxygenase levels in *Daphnia magna* and *Chironomus riparius* exposed to environmental pollutants.- *Ecotoxicol. envir. Saf.* 42: 9-15.
- Svensson, J. M., Bergman, E. and Andersson, G. 1999a. Impact of cyprinid reduction on the benthic macroinvertebrate community and implications for increased nitrogen retention.- *Hydrobiologia* 404: 99-112.
- Szító, A. and Mózes, K. 1999a. The Oligochaeta and the chironomida [sic!] fauna in the River Some /Szamos system.- In: Sárkány-Kiss, A. and Hamar, J. (eds.): *The Some /Szamos River valley. A study of the geography, hydrobiology*

- and ecology of the river system and its environment. *Tiscia Monogr. Ser.*, pp. 179-191. Szolnok, Szeged, Târgu Mure_.
- Takamura, K. 1999a. Wing length and asymmetry of male *Tokunagayusurika akamusi* chironomid midges using alternative mating tactics.- *Behav. Ecol. 10*: 498-503.
- Talcott, B. and Moore, M. S. 1999a. Getting across the nuclear pore complex.- *Trends Cell Biol. 9*: 312-318.
- Tátrai, I., Mátyás, K., Korponai, J., Paulovits, G., Pomogyi, P. and Présing, M. 1999a. Stable isotope analysis of food webs in wetland areas of Lake Balaton, Hungary.- *Arch. Hydrobiol. 146*: 117-128.
- Telesh, I. V., Alimov, A. F., Golubkov, S. M., Nikulina, V. N. and Panov, V. E. 1999a. Response of aquatic communities to anthropogenic stress: a comparative study of Neva Bay and the eastern Gulf of Finland.- *Hydrobiologia 393*: 95-105.
- Theiling, C. H. and Tucker, J. K. 1999a. Nektonic invertebrate dynamics and prolonged summer flooding on the lower Illinois River.- *J. Freshwat. Ecol. 14*: 499-510.
- Thompson, R. M. and Townsend, C. R. 1999a. The effect of seasonal variation on the community structure and food-web attributes of two streams: implications for food-web.- *Oikos 87*: 75-88.
- Tolonen, A. 1999a. Application of a bioenergetics model for analysis of growth and food consumption of subarctic whitefish *Coregonus lavaretus* (L.) in Lake Kilpisjärvi, Finnish Lapland.- *Hydrobiologia 390*: 153-169.
- Tucker, K. A. and Burton, G. A. Jr. 1999a. Assessment of nonpoint-source runoff in a stream using in situ and laboratory approaches.- *Envir. Toxic. Chem. 18*: 2797-2803.
- Ueno, R., Iwakuma, T. and Takamura, K. (1998)1999a. Seasonal and diel drift patterns of *Chironomus yoshimatsui* (Diptera: Chironomidae) in Otto River, Japan.- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc. 71*: 474-483.
- Vallenduuk, H. J. 1999a. *Key to the larvae of Glyptotendipes (Diptera, Chironomidae) in Western Europe.*- Vallenduuk, Schijndel. 46 + 10 pp.
- Vanin, S. e Turchetto, M. 1999a. (New data on the distribution of the glaciers-fly *Diamesa steinboeckii* Goetghebuer, 1933 (Diptera, Chironomidae).)- *Soc. venez. Sci. nat. Lavori 24*: 7-12.
- Venkatachalaiah, G., Prasanna-Kumar, R. and Bhanumathi, A. 1999a. Non-random association of linked inversions in an Indian midge, *Nilodorus [sic!] biroi*.- *Nucleus 42*: 66-68.
- Verneaux, V. et Aleya, L. 1999a. Comparaison des peuplements chironomidiens du lac de l'Abbaye obtenus par différentes méthodes d'échantillonnage. Intérêts de la récolte des exuvies nymphales.- *Revue Sci. Eau 12*: 45-63.
- Verneaux, V. et Aleya, L. 1999b. Diptères Chironomidés et caractérisation des lacs.- *Année biol. 35*: 220-234.
- Verschuren, D., Cocquyt, C., Tibby, J., Roberts, C. N. and Leavitt, P. R. 1999a. Long-term dynamics of algal and invertebrate communities in a small, fluctuating tropical soda lake.- *Limnol. Oceanogr. 44*: 1216-1231.
- Vignes, J.-C. 1999a. Rythmes alimentaires de jeunes Saumons Atlantiques (*Salmo salar* L.) et relations trophiques avec la dérive des invertébrés.- *Vie Milieu 49*: 293-300.
- Vinebrook, R. D. and Leavitt, P. R. 1999a. Differential responses of littoral communities to ultraviolet radiation in an alpine lake.- *Ecology 80*: 223-237.
- Vinyoles, D., Côté, I. M. and de Sostoa, A. 1999a. Egg cannibalism in river blennies: the role of natural prey availability.- *J. Fish Biol. 55*: 1223-1232.
- Vögtli, M., Imhof, M. O., Brown, N. E., Rauch, P., Spindler-Barth, M., Lezzi, M. and Henrich, V. C. 1999a. Functional characterization of two Ultraspiracle forms (CtUSP-1 and CtUSP-2) from *Chironomus tentans*.- *Insect Biochem molec. Biol. 29*: 931-942.
- Voogt, P. de, Bleeker, E. A. J., Vlaardingen, P. L. A van, Fernandez, A., Slobodnik, J., Wever, H. and Kraak, M. H. S. 1999a. Formation and identification of azarene transformation products from aquatic invertebrate and algal metabolism.- *J. Chromatogr. B 724*: 265-274.
- Vuori, K.-M., Luotonen, H. and Liljaniemi, P. 1999a. Benthic macroinvertebrates and aquatic mosses in pristine streams of the Tolvajärvi region, Russian Karelia.- *Boreal Env. Res. 4*: 187-200.
- Wallace, J. B., Eggert, S. L., Meyer, J. L. and Webster, J. R. 1999a. Effects of a source limitation on a detrital-based ecosystem.- *Ecol. Monogr. 69*: 409-442.
- Wang, X. 1999a. *Harnischia* Kieffer from China, with emendations to the diagnosis of the genus (Diptera: Chironomidae).- *Aquat. Insects 21*: 169-177.

- Watson, C. N. Jr. (1998)1999a. Description of the larva of *Apsectrotanypus algens* (Coquillett) with a review of the generic placement of the species (Diptera: Chironomidae).- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc.* 71: 241-246.
- Williams, D. D. and Williams, N. E. 1999a. Canadian springs. Postglacial development of the invertebrate fauna.- In: Batzer, D. P., Rader, R. B. and Wissinger, S. A. (eds.): *Invertebrates in freshwater wetlands of North America: ecology and management*, pp. 447-467. John Wiley & Sons, N. Y., Chichester.
- Wi_niewski, J. R., Krohn, N. M., Heyduk, E., Grasser, K. D. and Heyduk, T. 1999a. HMG1 proteins from evolutionary distant organisms distort B-DNA conformation in similar way.- *Biochim. biophys. Acta* 1447:25-34.
- Wi_niewski, J. R., Szewczuk, Z., Petry, I., Schwanbeck, R. and Renner, U. 1999a. Constitutive phosphorylation of the acidic tails of the high mobility group 1 proteins by casein kinase II alters their conformation, stability, and DNA binding specificity.- *J. biol. Chem* 274: 20116-20122.
- Wolfram, G., Donabaum, K., Schagerl, M. and Kowarc, V. A. 1999a. The zoobenthic community of shallow salt pans in Austria - preliminary results on phenology and the impact of salinity on benthic invertebrates.- *Hydrobiologia* 409: 193-202.
- Wolfram-Wais, A., Wolfram, G., Auer, B., Mikschi, E. and Hain, A. 1999a. Feeding habits of two introduced fish species (*Lepomis gibbosus*, *Pseudorasbora parva*) in Neusiedler See (Austria), with special reference to chironomid larvae (Diptera: Chironomidae).- *Hydrobiologia* 409: 123-129.
- Wotton, R. S. and Hirabayashi, K. 1999a. Midge larvae (Diptera: Chironomidae) as engineers in slow sand filter beds.- *Wat. Res.* 33: 1509-1515.
- Wrubleski, D. A. 1999a. Northern prairie marshes (Delta Marsh, Manitoba). 2. Chironomidae (Diptera) responses to changing plant communities in newly flooded habitats.- In: Batzer, D. P., Rader, R. B. and Wissinger, S. A. (eds.): *Invertebrates in freshwater wetlands of North America: ecology and management*, pp. 571-601. John Wiley & Sons, N. Y., Chichester.
- Wülker, W. (1998)1999a. A *Lobochironomus*-species with 3 chromosomes ($2n = 6$) - the true *Chironomus (Lobochironomus) mendax* Storå (Diptera, Chironomidae).- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc.* 71: 304-314.
- Wülker, W. 1999b. Fennoscandian *Chironomus* species (Dipt., Chironomidae) - identified by karyotypes and compared with the Russian and Central European fauna.- *Studia dipterol.* 6: 425-436.
- Wülker, W., Kiknadze, I. I., Kerkis, I. E. and Nevers, P. 1999a. Chromosomes, morphology, ecology and distribution of *Sergentia baueri*, spec. nov., *S. prima* Proviz & Proviz, 1997 and *S. coracina* Zett., 1824 (Insecta, Diptera, Chironomidae).- *Spixiana* 22: 69-81.
- Yiallourous, M., Storch, V. and Becker, N. 1999a. Impact of *Bacillus thuringiensis* var. *israelensis* on larvae of *Chironomus thummi thummi* and *Psectrocladius psilopterus* (Diptera: Chironomidae).- *J. Invert. Path.* 74: 39-47.
- Yamamoto, M. (1998)1999a. *Trichosmittia hikosana* n. gen. et n. sp. (Diptera: Chironomidae, Orthocladinae) from Japan.- In: Berg, M. B., Ferrington, L. C. Jr. and Hayford, B. L. (eds.): *A festschrift honoring Mary and Jim Sublette. Part 1: Taxonomy and systematics of Chironomidae.* - *J. Kans. ent. Soc.* 71: 263-271.
- Yan, Y.-j., Liang, Y.-l. and Wang, H.-zh. 1999a. Annual production of five species of Chironomidae (Diptera) in Houhu Lake, a typical algal lake (Wuhan, China).- *Chin. J. Oceanol. Limnol.* 17: 112-118.
- Yanoviak, S. P. 1999a. Effects of leaf litter species on macroinvertebrate community properties and mosquito yield in Neotropical tree hole microcosms.- *Oecologia* 120: 147-155.
- Yong, T.-S., Lee, J.-S., Lee, I.-Y., Park, S.-J., Park, G.-M., Ree, H.-I., Park, J.-W., Hong, C.-S. and Park, H.-S. 1999a. Identification of *Chironomus kiiensis* allergens, a dominant species of non-biting midges in Korea.- *Korean J. Parasit.* 37: 171-179.
- Youn, B.-J. and Chon, T.-S. 1999a. Effects of pollution on communities of Chironomidae (Diptera) in the Soktae stream a tributary of the Suyong River.- *Kor. J. Limnol.* 32: 24-34.
- Zhantiev, R. D. and Fyodorova, A. M. V. 1999a. (The fine structure of male Johnston's organ of *Chironomus plumosus* L. (Diptera, Chironomidae).)- *Ent. Obozr.* 78: 287-295.

The International Conference "Small rivers: Modern ecological state, actual problems"

April 25 - 28, 2001, Togliatti, Russia

Registration Form

Surname:

First name

Report Oral Poster

Organisation

Scientific degree

Position

Address

Phone _____ Fax _____

e-mail _____

Authors _____

The title of report _____

Signature
